

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Statesman*

EPA Reg. No. 62719-218

For Use in Crop Residue Management Systems in Soybeans (Preplant Application Only)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Authorization under FIFRA Section 3 (c)(7)(B) to use Statesman according to directions provided in this supplemental labeling expires on December 31, 1998. Use of this product according to these directions for use after December 31, 1998 will be illegal and sale of soybeans containing any residues of DMA 6 will constitute a violation of the Federal Food, Drug and Cosmetic Act.
- Read the label affixed to the container for Statesman before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Statesman according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container and this supplemental labeling.

Directions for Use

General Information

Statesman* herbicide may be used for postemergence control of many susceptible annual and perennial broadleaf weeds. This product may be applied prior to planting soybeans to provide foliar burn-down control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. Make only preplant applications to emerged weeds prior to planting soybeans grown in reduced tillage production systems. Apply only according to instructions given below.

Do not use any tillage operations between herbicide application and planting of soybeans.


Mixing Instructions

Compatible crop oil concentrates, and agricultural surfactants approved for use on growing crops may be added to spray mixtures to increase the herbicidal effectiveness of Statesman on certain weeds. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

Application Procedures

Apply using ground equipment in a spray volume sufficient to provide uniform coverage of weeds. Use 10 or more gallons per acre for ground equipment

(continued on back)



Application Timing and Use Rates

| Crop/Use/Application Timing | Statesman (pounds/acre) | Packet Factor (acres/packet) |
|---------------------------------------|-------------------------|------------------------------|
| Soybean (Preplant Only) | | |
| Not less than 7 days before planting | 0.6 | 2 |
| Not less than 30 days before planting | 1.1 | 1 |

Weeds Controlled

| | | |
|------------------------------|--------------------------|--------------------------|
| alfalfa† | geranium, Carolina | purslane, common |
| bindweed† | horseweed or mare's tail | ragweed, common |
| bullnettle | ironweed | ragweed, giant |
| bittercress, smallflowered | lambsquarters, common | shepherdspurse |
| buttercup, smallflowered | lettuce, prickly | smartweed, Pennsylvania† |
| cinquefoil, common and rough | morningglory, annual | sowthistle, annual |
| clover, red† | mousetail | speedwell |
| cocklebur, common | mustard, wild | thistle, Canada† |
| dandelion† | onion, wild† | thistle bull |
| dock, curly† | pennycress, field | velvetleaf |
| eveningprimrose, cutleaf | peppergrass† | vetch, hairy† |
| garlic, wild† | plantains | Virginia copperleaf |

†These weed species are only partially controlled.

For best weed control results, application should be made when weeds are small, actively growing and free of stress caused by temperature extremes, moisture stress, diseases, or insect damage. The control of individual weed species may be variable. Consult your local county agent or state agricultural extension specialist or crop consultant for advice.

Use Precautions and Restrictions

- **Important Notice:** Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of such injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.
- Do not use on sandy soils with less than 1.0% organic matter.
- Do not make more than one application per season regardless of the application rate used.
- Do not apply when weather conditions such as atmospheric temperature inversion or when wind direction favors drift from the treated area to susceptible plants.
- **Do not allow livestock grazing or harvest hay, forage or fodder from treated fields.** Livestock should be restricted from feeding/grazing of treated cover crops.
- In treated fields, plant soybean seed as deep as practical, but not less than 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- **Do not apply Statesman as described in this supplemental labeling unless you are prepared to accept the results of soybean injury, including possible stand loss and/or yield reduction.**
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with Statesman.

*Trademark of DowElanco

K1A/Statesman/Amend-Supp/02-19-96

Approved ___/___/___

Replaces 122-50-005.

Amendment(s):

1. Renewal of use for postemergence broadleaf weed control prior to planting soybeans under time-limited tolerance.