



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

62719-
220

Date of Issuance:

MAR 26 1993

NOTICE OF PESTICIDE:

 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: Until
Reregistration

Name of Pesticide Product:

Lorsban 4E-HF

Name and Address of Registrant (include ZIP Code):

DowElanco
Quad IV--9002 Purdue Road
Indianapolis, Indiana 46268-1189

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Submit, as you agreed in your letter dated February 25, 1993, new acute dermal and acute inhalation studies if the upgrades for these studies that are presently under review are found to be unacceptable.

3. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 62719-220."

Signature of Approving Official:

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Date:

MAR 26 1993

4. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide & Rodenticide Branch
Registration Division (H7505C)

Enclosure

3918

Contains aromatic petroleum distillates.

Lorsban 4E-HF may cause skin sensitization reactions in certain individuals.



Lorsban* 4E-HF

Insecticide

For control of various insects infesting certain field, fruit, nut and vegetable crops

Active ingredient:	
chlorpyrifos: O,O-diethyl-O-(3,5,6-trichloro-2-pyridinyl)phosphorothioate	41.7%
Inert ingredients	58.3%
Total	100.0%

Contains 4 pounds of chlorpyrifos per gallon.

EPA Reg. No. 62719-XXX
KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta se haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

May be fatal if swallowed. Harmful if absorbed through skin. Causes substantial but temporary eye injury. Causes skin irritation. Avoid breathing vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wear protective clothing, rubber gloves, and goggles. Wash thoroughly with soap and water after handling and before eating or smoking. Immediately remove contaminated clothing and wash before reuse. Destroy contaminated shoes.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Keep away from food, feedstuffs and water supplies. See page 2 for additional farm worker safety requirements under the heading "Notes to User - Reentry".

First Aid

If swallowed: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician or Poison Control Center immediately.

If in eyes: Flush with plenty of water for at least 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If inhaled: Remove to fresh air if effects occur. Get medical attention.

Note to physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Osmes, such as 2-PAM/procapsen, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of waste. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your cooperative agricultural extension service.

PHYSICAL OR CHEMICAL HAZARDS

Combustible - Do not use or store near heat or open flame. Do not cut or weld container.

NOTICE

READ THE ENTIRE LABEL. USE ONLY ACCORDING TO LABEL DIRECTIONS.

Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

IN CASE OF AN EMERGENCY

endangering health or the environment involving this product, call collect 817-438-4400

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 26 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-220

BEST AVAILABLE COPY

Lorsban* 4E-HF

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all "DIRECTIONS FOR USE" carefully before applying.

NOTE TO USER - REENTRY

Do not allow reentry into treated areas within 24 hours unless protective clothing is worn. Protective clothing consists of at least a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. These oral warnings shall inform workers of areas or fields that may not be entered without protective clothing and actions to take in case of accidental exposure as shown by the first aid section on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

WARNING: Area treated with Lorsban 4E-HF insecticide on (insert date of application). Do not enter without protective clothing for 24 hours. In case of accidental exposure.

STORAGE AND DISPOSAL

Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100°F for extended periods of time. Storage below 20°F may result in formation of crystals. If product crystallizes, store at 50 to 70°F and agitate to redissolve crystals. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For reusable portable containers, do not rinse or put any other material into container. When returning reusable portable container, seal all openings and return to owner.

Container Disposal: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Lorsban 4E-HF insecticide forms an emulsion when diluted with water and is suitable for use in all conventional spray equipment.

GENERAL USE PRECAUTIONS

Do not use on turf or for structural pest control. Do not formulate this product into other end use products. Attention: Do Not Cut or Weld Container.

MIXING DIRECTIONS

To prepare the spray, add a portion of the required amount of water to the spray tank and with agitation add the Lorsban 4E-

HF. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

Lorsban 4E-HF may also be used in tank mixtures with certain herbicides and/or with non-pressure fertilizer solutions as recommended under specific crop use directions. Note that Lorsban 4E-HF may be inactivated by sulfur containing fertilizer solutions. Prepare tank mixtures in the same manner as recommended above for use of Lorsban 4E-HF alone. When tank mixtures of Lorsban 4E-HF and herbicides are involved, add wettable powders first, flowables second, and emulsifiable concentrates last. Where a fertilizer solution is involved, it is strongly recommended that a fertilizer pesticide compatibility agent such as Unite or Complex be used. Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture. Do not allow spray mixtures to stand overnight.

Note: Test compatibility of the intended tank mixture before adding Lorsban 4E-HF to the spray or mix tank. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

SPRINKLER IRRIGATION

Lorsban 4E-HF may be applied by sprinkler irrigation for the following crop uses: alfalfa, citrus orchard floors, field corn, popcorn, sweet corn, cotton, cranberries, sorghum, and soybeans.

See the use sections for the individual crops for further application information. Do not apply this product to the above listed crops through any other type of irrigation system. Do not apply this product by chemigation to any other crop.

Special Use Directions

The following use directions are to be followed when Lorsban 4E-HF is applied through sprinkler irrigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injector with soap and water. Determine the amount of insecticide needed to cover the desired acreage. Pump the required Lorsban 4E-HF into a steel tank, start mechanical or hydraulic agitation, and add in order the non-emulsifiable oil and/or water. Continually agitate the mixture containing Lorsban 4E-HF. Set the sprinkler system to deliver the desired inches of water per acre. Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injector system according to number 14 in "Special Use Precautions" section. The mixture containing Lorsban 4E-HF must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

Special Use Precautions

The following use precautions will result in a safe and successful application of mixtures containing Lorsban 4E-HF

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
3. If you have questions about calibration, you should contact state extension service specialists, equipment manufacturers or other experts.

BEST AVAILABLE COPY

5918

4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. Refer to the American Society of Agricultural Engineers Engineering Practice 408 for more information.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. The metering pump must provide a greater pressure than that of the irrigation system at the point of injection. The pump must meet Section 675 for "Electrically Driven or Controlled Irrigation Machines" NEC 70 and must contain Viton or Teflon seals.
12. To insure uniform mixing of the insecticide into the water line, inject the mixture through a nozzle placed in the fertilizer injection port or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. It is suggested that the injection point be higher than the insecticide tank to prevent siphoning.
13. The steel tank holding the insecticide mixture should be large enough to allow the system to complete a revolution with one filling. It should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector pump.
14. In order to calibrate the irrigation system and injector to apply the mixture containing Lorsban 4E-HF, determine the following: 1) Calculate the number of acres irrigated by the system, 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area, 3) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes to cover the treatment area. This value equals the gallons per minute output that the injector must deliver. Convert the gallons per minute to milliliters or ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the injector pump be calibrated at least twice before operation, and the system should be monitored during operation.
15. Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application, if they irrigate nontarget areas.
16. Do not allow irrigation water to collect or runoff and pose a hazard to livestock, wells, or adjoining crops.
17. Allow foliage to dry before reentering the field.

18. Do not apply through sprinkler systems which deliver a low coefficient of uniformity such as certain water drive units.

Approved Crops

Alfalfa

Use Lorsban 4E-HF to control the following pests at the dosages indicated by application as a broadcast, foliar spray:

Pests	Lorsban 4E-HF
Aphids (suppression)	1/2 pint/acre
Corn rootworm adults (spotted cucumber beetle) Grasshoppers	1/2 - 1 pint/acre
Alfalfa blotch leaf miner Alfalfa looper Alfalfa weevil larvae and adults Armyworms Cutworms Egyptian alfalfa weevil larvae and adults Plant bugs Leafhoppers Spittlebugs	1 - 2 pints/acre

Note: Use higher rates to control spotted alfalfa aphid in California or Nevada. Stubble spray may be applied to control leafhopper in the Northeast.

Mix the required dosage with enough water to ensure thorough coverage of crop foliage and apply using aerial (fixed-wing or helicopter) or power-operated ground spray equipment. For aerial application use 2 to 5 gallons of water per acre. For best coverage when using ground application, a minimum of 20 gallons of water per acre with hollow cone nozzles is recommended. Control may be reduced at low spray volumes under high temperature and wind conditions. Treat when field counts or crop injury indicates that damaging pest populations are developing or present; however, do not apply more than once per crop cutting. Some reduction in insect control may be evident under excessively cool conditions. For Egyptian alfalfa weevil control in California, apply the specified dosage in a minimum of 5 gallons of water per acre when larvae are actively feeding and populations reach 15 to 20 larvae per 180" sweep with a 15-inch diameter net.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section for further information.

Lorsban 4E-HF should not be tank mixed with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination noninjurious under your current conditions of use. Some phytotoxic symptoms may be observed on young, tender rapidly growing alfalfa when treated with Lorsban 4E-HF. Alfalfa will outgrow the symptoms and no yield loss should be expected.

This product is highly toxic to bees exposed to direct treatment on alfalfa. Do not apply if nearby bees are clustered outside of hives and bees are foraging. Protective information may be obtained from your agricultural extension service.

Restrictions: Do not cut or graze treated alfalfa within 7 days after application of 1/2 pint of Lorsban 4E-HF per acre, within 14 days after application of 1 pint per acre, or within 21 days after application of rates above 1 pint per acre. Do not make more than 4 applications per year or apply more than once per crop cutting.

.6918

Lorsban® 4E-HF

Asparagus

Use Lorsban 4E-HF to control cutworms, asparagus aphids, and asparagus beetles by application at the rate of 2 pints per acre. Mix the specified dosage in sufficient water to ensure thorough coverage of treated plants and apply as a broadcast foliar spray. For cutworms, it is preferable to apply Lorsban 4E-HF when the soil is moist and worms are active on or near the soil surface. Applications may be made during the fern stage for control of asparagus beetles and asparagus aphids when field counts or crop injury indicates that damaging pest populations are developing or present.

Restrictions

Do not make more than one preharvest application per season or apply within one day of harvest. Do not make more than two postharvest applications during the fern stage. Based on available residue data, the use of Lorsban 4E-HF on asparagus is limited to the Midwest and Pacific Northwest.

Cherries

Use Lorsban 4E-HF for the control of lesser peach tree borer, greater peach tree borer, and American plum borer by application as a trunk spray. Mix 1 1/2 to 3 quarts of Lorsban 4E-HF with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks and lower limbs. Make a second application two weeks after the first one and a third application after harvest. Avoid contact with foliage in sweet cherries as premature leaf drop may result. Consult your state agricultural experiment station or extension service specialist for proper time to treat in your area.

In addition, one of the three allowable applications per year may be applied as a dormant spray for control of San Jose scale, peach twig borer, and climbing cutworms. For control of these pests, tank mix 1/2 to 1 pint of Lorsban 4E-HF with 1 to 2 gallons of a petroleum oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using ground spray equipment. For low volume (nonconcentrate) sprays (40 to 100 gallons of spray mixture per acre) use the same amounts of Lorsban 4E-HF and spray oil per acre required for application as a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of Lorsban 4E-HF for severe infestations. Use oil as recommended by your state agricultural experiment station or extension service specialist.

Restrictions

Make only three applications per year. Do not apply within 6 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

Christmas Trees (Nurseries and Plantations)

Use Lorsban 4E-HF at the rate indicated to control the following insects on the tree varieties listed.

Restrictions

Use Lorsban 4E-HF on tree plantations only in Connecticut, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oregon, Pennsylvania, South Carolina, Tennessee, Vermont, Virginia, Washington, and Wisconsin.

Do not allow livestock to graze in treated areas.

Tree Variety	Insects Controlled	Dosage Lorsban 4E-HF	Remarks
Balsam fir Blue spruce Concolor fir Douglas fir Eastern white pine Fraser fir	Ara Aphids Adelges (Coley) (Eastern spruce gall) European pine sawfly European pine shoot moth	1 qt/acre	Do not treat plants under extreme heat and drought stress. Apply to foliage in sufficient water to ensure adequate coverage.

Tree Variety	Insects Controlled	Dosage Lorsban 4E-HF	Remarks
Grand fir Noble fir Scotch pine White spruce	Grasshoppers Gypsy moth Mites ¹ (European red spider) (Two spotted spider) (except in WA & OR) Pine weevil (adult) Pine needle mege Pine sawtooth Plant bugs Spruce budworm Spruce budminer Scale ² (Pine needle) (Pine spruce) (Spruce bud) (Black pine) (Striped pine)	1 qt/acre	1. For effective control of adult spider mites if large numbers of eggs are present, apply a second spray 7 - 10 days after initial treatment to control newly hatched nymphs. 2. For scale control apply when scale crawlers are active.
	Pine weevil	3 qt/100 gal	Apply as a cut stump drench.

Citrus Fruits

Use Lorsban 4E-HF at the rates indicated according to the designated geographic area to control the following pests. Use the lower rates for light infestations and increase the dosage for heavier infestations.

A petroleum spray oil recommended for use on citrus trees may be added to dilute spray mixtures only at a rate of up to 1.8 gallons per 100 gallons of water to improve control of aphids, mealy bugs, scale insects, and thrips. Treat when insects become a problem or in accordance with the local spray schedule recommended by your state extension service specialist.

Lorsban 4E-HF may be applied in tank mixtures with ethion, dicofol, or Vendex. See "Mixing Directions" for further instructions. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with Lorsban 4E-HF.

Precautions

Observe local use directions for tank mix combinations especially in regard to applications of Lorsban 4E-HF plus spray oil. Consult with a county farm advisor, county agency, extension service personnel, agricultural commissioner, pest control advisor, or local DowElanco representative for such information regarding a given locality.

Do not apply when trees are stressed by drought or high temperatures.

Lorsban 4E-HF should not be tank mixed with Difolatan 80 Sprites as crop injury may occur.

Lorsban 4E-HF is highly toxic to bees exposed to direct treatment and should not be applied when bees are actively visiting the area. During the bloom period in California, apply from one hour after sunset until two hours before sunrise.

Restrictions

Do not apply more than 2 applications or more than 15 pints of Lorsban 4E-HF per acre per year. Do not make second foliar application within 30 days of the first application. Do not treat within 21 days of harvest for applications up to 7 pints of Lorsban 4E-HF per acre nor within 35 days for application of rates above 7 pints per acre. Do not do any work involving contact with trees within 2 days after treatment. Do not allow livestock to graze in treated areas.

Agri-Mek,

CITRUS FRUIT

Lorsban 4E-HF

Crop	Geographic Location	Pest	Dosage of Lorsban 4E-HF (pt/acre)	Spray Volume (gal/acre)	Remarks
Grapefruit Lemons Oranges and Other Citrus Fruit	California Arizona	Aphids Katydid Lepidopterous larvae Avocado leafroller Cutworms Fruit tree leafroller Orange (tortrix) Western sawtooth moth	2 - 7	ground: 100 - 750 aerial: min. of 15	Do not use a spray concentration of Lorsban 4E-HF of less than 1/2 pt/100 gal of total volume.
		Scale insects Black scale Brown soft scale California red scale	8 - 12	100 - 2400	
		Thrips (suppression) Mealy bugs	6 - 12	100 - 750	
		Additional Precautions for California and Arizona: Lorsban 4E-HF should not be used in combination with spray oil when temperatures are expected to exceed 95°F the day of application or for several consecutive days thereafter. Do not apply during the months of December, January or February.			
	Florida	Aphids Grasshoppers ^a Orange dogs Mealy bugs Scale insects Snow scale Florida red scale Purple scale Long scale Chaff scale Black scale Brown soft scale	2 - 7	ground: 100 - 1400 aerial: min. of 20	Do not use a spray concentration of Lorsban 4E-HF of less than 1/2 pt/100 gal of water per acre.
		Citrus rust mites	4 - 7	100 - 700	
		^a Lubber grasshoppers must be controlled when they are small (less than 1 inch in length) by direct contact with spray.			Do not use a spray concentration of Lorsban 4E-HF of less than 1 pt/100 gal of water per acre.
	Texas	Aphids Cutworms Katydid Mealy bugs Scale insects Brown soft scale California red scale Chaff scale	4 - 7	200 - 700	Do not use less than 2 pt of Lorsban 4E-HF per 100 gallons of water in dilute applications.
		Citrus rust mites (suppression)	4 - 7	200 - 700	
Small Transplanted Grapefruit, Orange and other Citrus Trees	Texas	Aphids Cutworms Katydid Mealy bugs Scale insects Brown soft scale California red scale Chaff scale	Max. of 7		Apply Lorsban 4E-HF at a rate of 1 fl oz/1 gal of water with a backpack sprayer. Apply to runoff.

BEST AVAILABLE COPY

BEST AVAILABLE COPY

8918

In Florida, do not apply more than 3 quarts per season.

Lorsban® 4E-HF

Citrus Orchard Floors

Use Lorsban 4E-HF to control red imported fire ants and other ant species by applying the specified dose in 25 or more gallons of water with ground application equipment that will uniformly apply the spray to the orchard floor. To control foraging ants and suppress mounds, apply Lorsban 4E-HF to the orchard floor at the rate of 3/4 to 1 quart per acre. Retreat as needed. For best insect control, uniform coverage of the orchard floor is necessary. Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Do not apply in tank mixtures with Evik herbicide. Foliar applications of Lorsban 4E-HF may be made in addition to the orchard floor treatments.

Lorsban 4E-HF may also be applied to citrus orchard floors through sprinkler irrigation systems only if the system uniformly covers the soil surface at the base of the tree. For best results, use the recommended amount of Lorsban 4E-HF per acre. See "Sprinkler Irrigation" section for further information.

Restrictions

Do not apply more than 10 quarts of Lorsban 4E-HF per acre per season. Do not apply last treatment within 28 days before harvest. Do not allow livestock to graze in treated areas. In Florida, do not make more than 3 applications per season.

Cranberries

Use Lorsban 4E-HF by application as a broadcast, foliar spray to control brown spearmoth, cranberry fruitworm, cranberry weevil, cutworms, fireworms, and Sparganothis fruitworms at the rate of 3 pints per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply no less than 5 gallons of spray per acre when using aerial equipment or no less than 100 gallons of spray per acre when using ground equipment. For weevil control, apply once at flower bud development (late May, early June) and, if weevils are present, once after 100% bloom (early to mid July). For other insects, treat when field counts indicate damaging insect populations are developing or present. Apply only after the winter flood has been removed. To avoid pesticide contamination of flood waters, make no applications while bogs are flooded.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems to control the above listed pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section for further information.

Restrictions

Do not make more than two applications per year or apply within 60 days before harvest.

Field Corn, Popcorn, Sweet Corn

For use to control cutworms, armyworms, corn earworm, corn rootworm adults, chinch bugs, grasshoppers, wireworms, flea beetle larvae and adults, aphids, billbugs, grubs, western bean cutworm, corn borers, symphylan, common stalk borer, and lesser cornstalk borer.

Preplant Incorporation Treatment

Use Lorsban 4E-HF at the following rates by application in sufficient water to the soil surface and incorporate into the soil:

Pests	Lorsban 4E-HF
Cutworms	2 - 4 pints/acre
Symphylan	
Wireworms	4 pints/acre
Billbugs	
Flea beetle larvae	
Grubs	
Seed corn maggots	
Seed corn beetle	6 pints/acre
Lesser cornstalk borer	
Corn rootworm larvae	

Note: For suppression of nematodes in North Carolina, South Carolina, and Virginia, apply 4 to 6 pints of Lorsban 4E-HF per acre.

Use recommended rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power-operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator or equivalent equipment.

Lorsban 4E-HF may also be applied in tank mixtures with non-pressure fertilizer solutions and/or with Bladex, Eradicane, Sutan, Lasso, Dual, and atrazine herbicides. See "Mixing Directions" section for further information. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with Lorsban 4E-HF.

Preplant, At-Plant or Preemergence Treatment in Conservation Tillage

Use Lorsban 4E-HF at the following rates by application in sufficient water to surface trash and exposed soil:

Pests	Lorsban 4E-HF
Cutworms	1 - 2 pints/acre
Armyworms	

Use recommended rate in not less than 20 gallons of water per acre and apply as a broadcast spray using suitable power-operated ground spray equipment. Use higher rates for residual control.

Lorsban 4E-HF may also be applied in tank mixtures with non-pressure fertilizer solutions and/or with paraquat and Roundup. See "Mixing Directions" section for further information. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with Lorsban 4E-HF.

Cultivation Time Treatment

Use Lorsban 4E-HF at the rate of 2 pints per acre to control corn rootworm larvae. Apply Lorsban 4E-HF as a water emulsion on both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. The best time to apply a basal treatment of a soil insecticide with cultivation is near the beginning of egg hatch. A cultivation application of Lorsban 4E-HF may be made in addition to an at planting application of Lorsban® 15G granular insecticide.

Postemergence Treatment

Use Lorsban 4E-HF at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Lorsban 4E-HF
Grasshoppers	1/2 - 1 pint/acre
Armyworms	1 - 2 pints/acre
Chinch bugs	
Aphids	
Corn rootworm adults	
Webworms	
Western bean cutworm	
European corn borer (see note)	1 1/2 - 2 pints/acre
Southwestern corn borer	
Corn earworm	2 - 3 pints/acre
Cutworms	
Billbugs	
Lesser cornstalk borer	
Flea beetle adults	
Common stalk borer	

Note: The recommended dosage will control silk clipping by corn rootworm adults. For European corn borer control, use

Field Corn, Popcorn, Sweetcorn (Including Grown for Seed)

BEST AVAILABLE COPY

7 of 18

1 1/2 to 2 pints per acre when application is made with power-operated ground and aerial equipment and 1 to 2 pints per acre when application is made through a sprinkler irrigation system.

Treat when field counts indicate that pests are or may become a problem. For beet bug, chinch bug, and flea beetle control, apply with sufficient water to ensure a minimum spray volume of 20 to 40 gallons per acre and 40 psi using ground spray equipment. On corn less than 6 inches tall, apply the insecticide spray in a 9 to 12 inch wide band over the row. On corn greater than 6 inches tall, apply the insecticide spray using drop nozzles directed to the base of the plant. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pressure, a second application of Lorsban 4E-HF may be needed.

For cutworm, webworm, western bean cutworm, armyworm, aphid, European and southwestern corn borer, grasshopper, lesser cornstalk borer, corn rootworm adult, corn earworm, and common stalk borer control, apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment. For aerial application use 2 to 5 gallons of spray per acre. Control may be reduced at low spray volumes under high temperature and wind conditions. For cutworms, it is preferable to apply Lorsban 4E-HF when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy or crusty at time of treatment, worms may be protected from the spray and effectiveness will be reduced. If such conditions exist, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. Consult your agricultural experiment station or extension service specialist for additional information concerning control practices in your area. For webworm control, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary. For first-generation European corn borer control, treat when 25 to 50 percent of the corn plants show pinhole feeding or leaf-feeding scars. Ground applications should be directed into the corn whorls. For second-generation European corn borer control, treat when field counts of egg masses indicate pests are or may become a problem. For southwestern corn borer control, treat when field counts of egg masses indicate pests are or may become a problem. A second application may be applied 10 to 14 days later, if needed due to reinfestation. For common stalk borer control, treat approximately 11 days after application of Roundup herbicide or after complete burndown with paraquat herbicide (3 to 5 days). Do not use Lorsban 4E-HF in combination with the burndown herbicide for control of common stalk borer.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar insects. For best results, use the recommended rate of Lorsban 4E-HF in a tank mix with 2 pints per acre of non-emulsifiable oil. Maintain vigorous tank agitation to assure uniformity of the Lorsban 4E-HF plus oil mixture throughout the injection period. Lorsban 4E-HF may also be applied through sprinkler irrigation systems at the rate of 2 to 3 pints per acre to control corn rootworm larvae. Time application to coincide with the appearance of the second instar larvae. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. Apply with enough water to wet the root zone to the depth control is needed. Under saturated soil conditions, allow enough soil drying to occur so that an application using a minimum water rate will not produce runoff. Consult university extension personnel or other experienced consultants to determine the need to treat and to aid in application timing. See "Sprinkler Irrigation" section for further information.

Restrictions

Do not apply within 35 days before harvest of grain. Do not apply more than a total of 15 pints of Lorsban 4E-HF per acre per season. Do not allow livestock to graze in treated areas nor harvest treated corn silage as feed for meat or dairy animals within 14 days after last treatment. Do not feed treated corn fodder to meat or dairy animals within 35 days after last treatment.

Sweet Corn Grown Only In Florida and Georgia

Use Lorsban 4E-HF to control infestations of beet armyworm, fall armyworm and corn earworm by application as a broadcast, foliar spray at the rate of 1 to 2 pints per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment. For aerial application, use at least 2 gallons of spray per acre. Treat when field counts indicate damaging pest populations are developing or present. Re-treat as necessary to maintain control but do not apply more than twenty-two 1-pint or eleven 2-pint treatments per season.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar insects. For best results, use the recommended rate of Lorsban 4E-HF in a tank mix with 2 pints per acre of non-emulsifiable oil. Maintain vigorous tank agitation to assure uniformity of the Lorsban 4E-HF plus oil mixture throughout the injection period. See "Sprinkler Irrigation" section for further information.

Restrictions

Do not apply more than 22 pints of Lorsban 4E-HF per acre per season. Do not harvest corn ears, allow livestock to graze in treated areas, nor feed treated silage, fodder, or grain to meat or dairy animals within 21 days after treatment. Do not use in conjunction with postplant broadcast, foliar applications of Lorsban 15G.

Cotton

Use Lorsban 4E-HF for control of the following pests in all states except Arizona and California at the dosages indicated:

Pests	Lorsban 4E-HF
Cotton fleahopper	3/8 - 1 pint/acre
Plant bugs (Lycus, Miridae)	
Cotton aphid	1/2 - 1 pint/acre
Fall armyworm	
Grasshoppers	
Thrips	
Yellow-striped armyworm	
Spider mites	1 pint/acre
Beet armyworm	1 1/2 - 2 pints/acre
Bollworms (Heliothis spp.)	
Boll weevil	
Cutworms	
Pink bollworm	
Belt marsh	
Caterpillar	

Insert 1.

Note: The recommended dosage rate of 3/8 pint per acre will not achieve the high degree of control of the higher label rate, but will minimize the damage done by plant bugs and cotton fleahopper and allow the beneficial insects to survive, build up, and be available to aid in the control of bollworms infesting cotton.

Use Lorsban 4E-HF for control of the following pests in Arizona and California at the dosages indicated:

BEST AVAILABLE COPY

Lorsban® 4E-HF

Insert 2:

Pests	Lorsban 4E-HF
Amyworms Cotton aphid Cotton fleahopper Lygus Salt marsh caterpillar Thrips	1 - 2 pints/acre
Bollworms (<i>Heliothis</i> spp.) Boll weevil Cutworms Pink bollworm	2 pints/acre

Note: The 2 pint rate will aid in the suppression of cotton leafperforator and spider mites.

Mix the required dosage with sufficient water to ensure thorough coverage of plants and apply using aerial or power-operated ground spray equipment. For aerial application, use at least 1 gallon of spray per acre. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section for further information.

For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly hatched nymphs.

For best results on bollworms, it is suggested that fields be scouted twice per week and treatments made when worms are 1/4 inch or less in length. The following table illustrates the size of worms in relation to age and stage of development (inset) as a guide to timing of treatments for best control.

From the table it can be seen that a scouting schedule of only once per week will not be satisfactory since the worms may be too big to control effectively by the seventh or eighth day.

Timing for the Best Worm Control

Age (days)	Size	Instar
1	1/16"	I
3	3/32"	II
5	5/32"	III
7	7/16"	IV
9	1 1/16"	

Proper application techniques help to ensure thorough spray coverage and correct dosage and are thus important in obtaining good control of pests. Consider these suggestions when applying Lorsban 4E-HF on cotton.

Aerial Application

Shorten boom length to avoid spray entering the vortices at the wing tips. Swath width should be reduced when wind direction is the same as direction of spraying.

The proper nozzle arrangement and swath width to avoid skips and vortices effect can be checked out by flying over a paper tape (adding machine paper) using water with or without soluble dye. (The dye gives a permanent record.)

Flying at a height of 5 to 15 feet above the target results in the best coverage.

Nozzle orientation of the boom is important. More break-up occurs when nozzles are pointed straight down versus the straight back position. Desired droplet size (100 to 200 microns) can be obtained by angling the nozzles somewhere in this range.

Marking of swath by flagging or permanent markers is essential.

Ground Application

Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom; drift spray is wasted spray so do not depend on it. Use flat fan or disc-core hollow cone nozzles with maximum spacing of 20 inches and a spray pressure of 40-60 psi with a droplet size of 100 - 200 microns.

Restrictions

Do not apply within 14 days before harvest. Do not allow livestock to graze in treated areas. Do not feed gin trash or treated forage to livestock.

Lorsban 4E-HF may also be used in tank mixtures in all states to control armyworms, bollworms (*Heliothis* spp.), boll weevil, cotton fleahopper, plant bugs (*Lygus*, *Mirids*), and spider mites as indicated in the following table.

Tank Mixture	Rate/Acre
Lorsban 4E-HF plus Ambush 2E	1 - 2 pints plus 1/5 - 2/5 pint
Lorsban 4E-HF plus Pounce 3.2E	1 - 2 pints plus 1/8 to 1/4 pint
Lorsban 4E-HF plus Pydrin 2.4E	1 - 2 pints plus 1/8 - 1/3 pint
Lorsban 4E-HF plus Bolstar 6E	1/2 - 1 pint plus 2/3 - 1 1/3 pints
Lorsban 4E-HF plus methyl parathion† plus EPN†	1 pint plus 0.6 lb a.i. plus 0.6 lb a.i.

†Use only emulsifiable formulations containing these active ingredients (a.i.).

Mix the tank mixture components in sufficient water to ensure thorough coverage of plants (at least 1 gallon of spray per acre) and apply using aerial spray equipment. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control.

In California and Arizona, use Lorsban 4E-HF at 1 to 2 pints per acre plus Monitor 4E at 1 to 2 pints per acre or Lorsban 4E-HF at 1 to 2 pints per acre plus Orthene 75S at 0.33 to 1.3 pounds per acre for control of armyworms, bollworms, cabbage looper, cotton aphids, cotton fleahopper, cutworms, grasshopper, omnivorous leafroller, pink bollworm, plant bugs (*Lygus*, *Mirids*), saltmarsh caterpillar, thrips, and whitefly (suppression). Apply in 10 to 25 gallons of water per acre with ground spray equipment or in 5 to 10 gallons of water per acre with aerial spray equipment. Treat when field counts indicate that damaging insect populations are developing or present. For tank mixtures which include Monitor 4E, apply before 65 percent of the bolls have opened. Re-treat as necessary to maintain control.

Restrictions

Make no applications of these tank mixtures closer to harvest than the longest pre-harvest interval shown for any of the products in the tank mixture. Do not allow livestock to graze in

Do not apply within 14 days before harvest or make more than 6 applications per season. Do not allow livestock to graze in treated areas. Do not feed gin trash or treated forage to livestock.

Delete

For best results on bollworms and budworm it is suggested that fields be scouted twice per week and...

trashed areas. Do not feed gin trash or trashed forage to livestock. Workers entering the field within 48 hours after application should wear protective clothing.

Figs

Use Lorsban 4E-HF at the rate of 2 quarts per acre for control of dried fruit beetle by application in sufficient water to the soil surface followed by incorporation into the top 3 inches of soil. Apply to fig orchard soil as a dormant application in late winter prior to beetle emergence and prior to leaf formation.

Restrictions

Make only one application per year. Do not apply within 7 months of harvest. Based on available residue data, use of Lorsban 4E-HF on figs is restricted to California.

Grapes

Use Lorsban 4E-HF for control of grape root borer by application just before the pest emerges from the soil. Mix 4 1/2 pints of Lorsban 4E-HF with 100 gallons of water and apply 2 quarts of the diluted spray mixture to the soil surface on a 15-square foot area around the base of each vine. Do not allow spray to contact fruit or foliage.

Restrictions

Do not make more than one application per season or apply within 35 days before harvest. Based upon available residue data, the use of Lorsban 4E-HF in grapes is restricted to states east of the Rocky Mountains.

Mint

Use Lorsban 4E-HF by application as a broadcast, foliar spray to control cutworms at the rate of 2 to 4 pints per acre and mint root borer at the rate of 4 pints per acre. Mix the specified dosage in water to give no less than 10 gallons of spray per acre and apply using ground spray equipment. For cutworm control, treat during May and June when field counts indicate damaging insect populations are developing or present. When larvae are less than 3/4 inch in length, use the 2 pint rate. When larvae are 3/4 inch or more in length, use the higher rate. Make only one application during the growing season. Do not apply within 90 days before harvest. For mint root borer control, apply post-harvest when field counts indicate damaging insect populations are developing or present. Follow treatment with approximately 1 acre inch of sprinkler irrigation immediately after application to incorporate the insecticide into the soil. Make only one post-harvest application per season.

Nectarines, Peaches

Use Lorsban 4E-HF for the control of peach tree borers by application as a trunk spray before newly hatched borers enter the trees. Mix 3 quarts of Lorsban 4E-HF with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your state agricultural experiment station's or extension service specialist's written recommendations for proper time to treat in your area.

Lorsban 4E-HF may also be used as a preplant dip application for non-bearing peach trees at the equivalent application rate of 3 quarts per 100 gallons of water for control of peach tree borer. Dip trees several inches above the grafting bud scar and plant immediately or allow to dry before returning to storage. Do not allow peach trees to remain in contact with the dip solution.

Restrictions

Make only one application per season. Do not apply within 14 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

Onions (Dry Bulb)

Use Lorsban 4E-HF to control onion maggot by application as an in-furrow drench. Apply Lorsban 4E-HF at the rate of 1 1

fluid ounce per 1000 linear feet of row at an 18-inch row spacing. Use a minimum of 40 gallons of total drench per acre. Incorporate to a depth of 1 to 2 inches.

Restrictions

Do not make more than one application per year.

Peanuts

Use Lorsban 4E-HF for control of lesser cornstalk borer by application as a directed spray to the base of the plants in an 8 to 10 inch band. Apply Lorsban 4E-HF at the rate of 2 to 4 pints per acre when field counts indicate damaging insect populations are developing or present. Mix the required dosage in enough water to ensure thorough coverage. Do not apply as a foliage application.

Restrictions

Do not make more than one application per season. Do not harvest within 21 days after treatment. Do not feed treated peanut forage or hay to meat or dairy animals.

Sorghum

Use Lorsban 4E-HF insecticide for control of the following pests at the dosages indicated:

Pests	Lorsban 4E-HF	Special Directions
Sorghum midge	1/2 pint/acre	Apply when 30 to 50% of the seed heads are in bloom, repeat at 3-day intervals if necessary.
Greenbugs Grasshoppers Yellow sugar cane aphid and other aphids	1/2 - 1 pint/acre	
Chinch bugs Lesser cornstalk borer	1 - 2 pints/acre	Apply as a directed spray toward the base of the plant using power-operated ground spray equipment with sufficient water to ensure coverage of an 8 to 12 inch band centered on the row. On plants less than 6 inches high, apply an 8 to 12 inch band over the row. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone.
Webworms	1 pint/acre	
Armyworms Corn earworm Cutworms	1 - 2 pints/acre	
European and Southwestern corn borer	1 1/2 - 2 pints/acre	

Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment. To minimize chemical injury, do not apply Lorsban 4E-HF to drought stressed grain sorghum within 3 days following irrigation or rain except where the product is applied in irrigation water.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section for further information.

Restrictions

The treated crop is not to be used for grain, forage, fodder, hay, or silage within 30 days after application of 1 pint of Lorsban 4E-HF per acre or within 60 days after application of rates above 1 pint per acre. Do not treat sweet varieties of sorghum. Do not apply more than 3 pints of Lorsban 4E-HF per acre per season.

Soybeans

For use to control armyworms, bean leaf beetle, corn earworm, cutworms, European corn borer, grasshoppers, green cloverworm, lesser cornstalk borer, Mexican bean beetle, saltmarsh caterpillar and other woolly bears, southern green stink bug, spider mites, and velvetbean caterpillar.

Soil Treatment

Use Lorsban 4E-HF at the rate of 1 to 2 pints per acre to control cutworms and lesser cornstalk borer. Mix the specified dosage in a minimum of 10 gallons of spray per acre and apply to the soil surface using suitable ground spray equipment. Equivalent rates of insecticide spray required per 100 feet of row for various row spacing are given in the accompanying table. For at-plant treatments apply the insecticide over the row in a 4 to 6-inch band in front of the planter shoe or press wheel or after the press wheel followed by a drag chain for light incorporation. Do not apply as an in-furrow treatment. For postemergence rescue treatments, apply as a directed spray in a 9 to 12-inch band at the base of the plant. To plants under 6 inches high apply over the top in a 6- to 12-inch band. Treat when field counts or conditions indicate that pests are or may become a problem.

Fluid Ounces of Spray Required Per 100 Feet of Row for Various Row Spacing				
Volume of Spray Per Acre	30"	32"	36"	40"
10 gallons	8.8	7.9	6.9	5.9
15 gallons	13.2	11.8	10.3	8.8
20 gallons	17.6	15.7	13.7	11.8

Foliar Treatment

Use Lorsban 4E-HF at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Lorsban 4E-HF
European corn borer Southern green stink bug	2 pints/acre
Bean leaf beetle Cutworms Corn earworm Saltmarsh caterpillar and other woolly bears	1 - 2 pints/acre
Mexican bean beetle Armyworms	1 - 1 1/2 pints/acre
Velvetbean caterpillar Grasshoppers Green cloverworm Spider mites	1/2 - 1 pint/acre

Apply as a broadcast spray using either aerial or ground equipment when field counts indicate damaging insect populations are developing or present; re-treat as necessary to maintain control. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly-hatched nymphs. On determinate soybeans do not apply more than one application after pod set.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to

control the above listed foliar pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section for further information.

Restrictions

Do not apply more than 6 pints of Lorsban 4E-HF per acre or 3 pounds of chlorpyrifos (active ingredient) per acre per season. Do not apply last treatment within 28 days before harvest nor apply last two treatments closer than 14 days apart. Do not allow livestock to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals.

Strawberries

Use Lorsban 4E-HF by application as a broadcast foliar spray to control strawberry bud weevil at the rate of 1 quart per acre. Apply in a minimum of 40 gallons of spray per acre when buds first appear and 10 to 14 days later. Do not apply after berries start to form or when berries are present. Lorsban 4E-HF should not be tank mixed with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination noninjurious under your current conditions of use. Phytotoxicity may occur when Lorsban 4E-HF is applied to strawberries experiencing high temperature and drought stress.

Restrictions

For pre-bloom use only. Do not make more than two applications per season or apply within 21 days before harvest.

Sunflowers

For use to control cutworms, sunflower beetle larvae and adults, stem weevil, sunflower moth, banded sunflower moth, woolly bears, seed weevil and grasshoppers.

Preplant Incorporation Treatment

Use Lorsban 4E-HF at the following rates by application in sufficient water to the soil surface and incorporate into the soil:

Pests	Lorsban 4E-HF
Cutworms	2 - 4 pints/acre

Use recommended rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power-operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator or equivalent equipment.

Postemergence Treatment

Use Lorsban 4E-HF for control of the following pests at the dosage indicated by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Lorsban 4E-HF
Cutworms	2 - 3 pints/acre
Sunflower beetle larvae and adults Stem weevil Sunflower moth Banded sunflower moth Woolly bears Seed weevil	1 - 1 1/2 pints/acre
Grasshoppers	1 pint/acre

Apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment when field counts indicate that pests are or may become a problem. For cutworm control, a second treatment may be made 7 to 10 days later, if needed. For stem weevil control, optimal treatment time is within 5 to 7 days after adult weevils begin to appear. For sunflower moth control, make first application during early 1 to 5 percent bloom stage. A second treatment may be made 7 days later, if needed. For seed weevil control,

treat when field counts indicate there are 10 to 12 adults per plant for oil crops and 1 to 3 adults per plant on confectionary crops. Additional treatments should be made at successive 7 to 10 day intervals if field counts indicate need to re-treat. For sunflower beetle larvae or adult control, treat when field counts indicate there are 10 larvae or 1 to 2 adults per seedling. Additional treatments may be made at successive 7 to 10 day intervals if field counts indicate need to re-treat.

Restrictions

Do not apply more than 9 pints of Lorsban 4E-HF per acre per season. Do not apply within 42 days before harvest. Do not allow livestock to graze in treated areas.

Sugar Beets

Use Lorsban 4E-HF by application as a broadcast, foliar spray to control beet armyworms at a rate of 1 1/2 to 2 pints per acre and cutworms at a rate of 2 pints per acre. Treat when field counts indicate that damaging insect populations are or may be a problem. Mix the specified dosage with 2 to 5 gallons of water per acre and apply using suitable aerial spray equipment or with 10 to 30 gallons of water per acre when using power-operated ground spray equipment. Re-treat as necessary to maintain control.

Restrictions

Do not apply within 30 days before harvest of beet roots and tops. Do not apply more than a total of 8 pints of Lorsban 4E-HF per acre per season. Do not allow livestock to graze in treated areas nor harvest treated beet tops as feed for meat or dairy animals within 30 days after last treatment.

Sweet Potatoes

Use Lorsban 4E-HF to reduce the feeding damage caused by populations of *Conoderus* wireworm, *Systena* flea beetle and the sweet potato flea beetle. Apply at the rate of 4 pints per acre as a broadcast (overall) spray to the soil surface followed by incorporation. Mix the specified dosage with enough water to obtain uniform coverage and apply as a coarse spray using suitable ground spray equipment. Incorporate the insecticide to a depth of 4 to 6 inches as soon as possible after application by using a rotary hoe, disc cultivator, or other suitable incorporation equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that Lorsban 4E-HF will protect against feeding damage). Lorsban 4E-HF will not control false wireworms or white fringe beetle or other grubs that attack sweet potatoes.

Restrictions

Do not make more than one application per season. Do not harvest within 125 days of treatment.

Tobacco

Use Lorsban 4E-HF for preplant treatment to control larvae of cutworms, flea beetles, mole crickets, root maggots, and wireworms. Apply 2 to 3 quarts of Lorsban 4E-HF per acre in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface one week before transplanting. Immediately following application, incorporate the insecticide into the soil to a depth of 2 to 4 inches using suitable equipment. The application of Lorsban 4E-HF will also suppress the movement of imported fire ants into treated fields.

To control the above insects and low to moderate populations of rootknot nematodes in North Carolina, South Carolina, and Virginia, use Lorsban 4E-HF at the rate of 5 quarts per acre. To control the above insects and moderate populations of rootknot nematodes in all tobacco growing regions, use Lorsban 4E-HF in a tank mix with NemaCur 3 at the rate of 2 quarts of Lorsban 4E-HF plus 4 quarts of NemaCur 3 per acre. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for NemaCur 3 used in combination with Lorsban 4E-HF. Apply the specified dosage in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment. Where the nematode species *Heterodoglyne*

arenaria or *M. javanica* are present or high populations of *M. incognita*, apply Telone II soil fumigant at the recommended label rate.

Restrictions

Do not make more than one application per season.

Tree Fruits

Use Lorsban 4E-HF as a dormant or delayed dormant spray at the rates indicated to control the following insects on the crops listed. While Lorsban 4E-HF may be used without oil, oil is recommended for additional pest control.

Crop	Insect	Dosage Lorsban 4E-HF
Apples	Rosy apple aphid San Jose scale Lygus Pandemia leafroller Climbing cutworms	1/2 - 1 pint
Pears	San Jose scale Climbing cutworms Pear psylla adults	
Plums Prunes	San Jose scale Mealy plum aphid Climbing cutworms Peach twig borer	
Almonds Peaches Nectarines	San Jose scale Peach twig borer Climbing cutworms	

Insert 4.

Tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using suitable ground spray equipment. (See "Additional Precautions Specific to California" for use in California.) For low volume (concentrate) sprays (40 to 100 gallons of spray mixture per acre), use the same amount of Lorsban 4E-HF and spray oil per acre required for application as a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of Lorsban 4E-HF for severe infestations. Use oil as recommended by your state agricultural experiment station or extension service specialist.

Insert 5.

Precautions

Because cold or dry conditions may cause Lorsban 4E-HF plus oil sprays to injure trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated.

Additional Precautions Specific to California: Use a minimum of 250 gallons of total spray volume per acre. Do not use more than 4 pints of Lorsban 4E-HF per acre. Do not use more than 4 gallons of spray oil per acre on almonds, peaches, or nectarines. Do not use any adjuvants or surfactants in addition to or as a substitute for a petroleum spray oil in a tank mix with Lorsban 4E-HF. Do not apply on almonds in the following counties in California: Butte, Colusa, Glenn, Sutter, Tehama, Yuba, and Yuba.

Restrictions

Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated orchards.

Tree Nuts

Use Lorsban 4E-HF at the dosages indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop and apply as a concentrate or dilute spray using conventional, power-operated spray equipment. For dilute sprays applied to tree nut crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of Lorsban 4E-HF per acre. Treat when pests appear or in accordance with local conditions. Insect control by aerial application may be less than control by ground application because of less coverage. Consult your state agricultural experiment station, certified pest control adviser, or extension service specialist for specific use information in your area.

Lorsban® 4E-HF

ALMONDS, FILBERTS, WALNUTS			
Use Lorsban 4E-HF at the rates indicated to control the listed pests			
Crop	Insects Controlled	Dosage Lorsban 4E-HF	Restrictions
Almonds	Naval orangeworm Peach twig borer San Jose scale	4 pints/acre	Make no more than 3 foliar applications per season on almonds and filberts and no more than 2 applications per season on walnuts. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated orchards.
Filberts	Eye-spotted bud moth Filbert aphid Filbert leafroller Filbert worm Oblique-banded leafroller Omnivorous leafminer Winter moth	3 - 4 pints/acre	
Walnuts	Codling moth Walnut scale	4 pints/acre	

PECANS		
Use Lorsban 4E-HF at the rates indicated to control the listed pests.		
Insects Controlled	Dosage of Lorsban 4E-HF (Dilute or Concentrate) Pints/Acre	Remarks and Restrictions
Pecan nut casebearer Fall webworm	1 1/2 - 4	<p>Make no more than five applications per season year</p> <p>Do not apply within 28 days of harvest.</p> <p>Do not allow livestock to graze in treated orchards.</p> <p>Make no applications of tank mixtures closer to harvest than the longest pre-harvest interval shown for any of the products in the tank mixture.</p> <p>For dilute applications with ground equipment use at least the minimum rate of Lorsban 4E-HF listed for the pest. Apply in 100 - 600 gallons of water per acre.</p> <p>For aerial applications use 5 - 15 gallons of water per acre.</p> <p>Note: With aerial application control may be reduced due to poor coverage.</p> <p>Up to 20 pints of Lorsban 4E-HF may be applied per acre per season year</p>
Phylloxera spp. ¹ Black pecan aphid Hickory shuckworm ² Pecan leaf scorch mite (suppression) ³ Fire ants and other ant species ⁴	2 - 4	
Yellow pecan aphid Black margined aphid	1 - 4 pints of Lorsban 4E-HF plus: 5.33 fl oz of Pydrin 2.4E, or 1.70 fl oz of Azena 1.9EC, or 3.00 fl oz of Amvr. 2.5 EC, or 2.56 fl oz of Cymbush 3E	

- ¹ For best *Phylloxera* spp. control, make 2 applications at a 7 to 10 day interval using a minimum of 1.0 pint of Lorsban 4E-HF per acre starting at bud swell.
- ² For best results make 2 applications, 10 to 14 days apart.
- ³ To suppress pecan leaf scorch mite, use a preventative program.
- ⁴ For ant control, apply as an orchard floor spray. Do not apply where weed growth or other obstructions prevent uniform coverage of the orchard floor.

Almond Orchard Floors

Use Lorsban 4E-HF to control southern fire ant and pavement ant by applying the specified dose with ground application equipment that will uniformly apply the spray to the orchard floor. Use when ant activity becomes evident within the orchard. Since worker ants cease most of their foraging activity at temperatures above 90°F, best results will be achieved with applications made at temperatures below 90°F at the time of application. Dosage of Lorsban 4E-HF and spray volume may vary depending on the irrigation method employed in the orchard as follows:

Ant Control in Sprinkler- or Drip-Irrigated Orchards

Apply Lorsban 4E-HF as a broadcast spray to the entire orchard floor using ground spray equipment at 4 to 8 pints per acre in 25 or more gallons of water. Use the high rate for heavy infestations and the low rate for light infestations. In orchards where ant activity is concentrated around the irrigation emitters, apply the high rate to a 6 to 8 foot band along the drip-irrigation line and the low rate to the rest of the orchard.

Ant Control in Flood-Irrigated Orchards

Apply Lorsban 4E-HF at 4 to 8 pints per acre in 25 or more gallons of water to the entire orchard floor using ground spray equipment. Apply the high rate to heavily infested areas and the low rate to lightly infested areas. Where ant colonies are abundant only in the berm areas, apply Lorsban 4E-HF at 8 pints per treated acre in 50 or more gallons of water to a 6 to 10 foot band along the tree line (berm).

Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Mow or chemically control weeds before the application of Lorsban 4E-HF. Foliar applications of Lorsban 4E-HF may be made in addition to the orchard floor treatment.

Restrictions

Do not apply more than 16 pints of Lorsban 4E-HF per ~~season~~ year to the orchard floor. Do not apply the last treatment within 14 days of harvest. Do not allow livestock to graze in treated orchards.

Vegetables

Use Lorsban 4E-HF at the dosages indicated to control the pests listed in the following table. To avoid phytotoxicity in vegetables, do not mix with other pesticide products or treat plants that are under extreme heat and drought stress.

Crop	Insects Controlled	Dosage Lorsban 4E-HF	Use Directions	Restrictions
Cauliflower	Root maggot	1.6 - 2.4 fl oz/ 1000 linear ft of row	For direct seeded crops apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the presswheel is recommended.	Do not apply more than 2 pints of Lorsban 4E-HF to cauliflower planted in 40 inch rows. Use proportional amounts for other row spacings not to exceed 4 pints of Lorsban 4E-HF per acre.
Broccoli Brussels sprouts Cabbage Chinese cabbage Collards Kale Kohlrabi Turnips	Root maggot	1.6 - 2.75 fl oz/ 1000 linear ft of row	For transplanted crops, apply Lorsban 4E-HF as a water-based spray directed to the base of the plants immediately after setting. Use a minimum of 40 gallons of total spray per acre. Do not add any additional adjuvants, surfactants or spreader stickers. Do not apply as a foliage application.	Do not apply more than 2.6 pints of Lorsban 4E-HF per acre to broccoli, brussels sprouts, cabbage, Chinese cabbage, collards, kale, kohlrabi, and turnips planted in 40-inch rows. Do not apply more than 4 1/2 pints of Lorsban 4E-HF per acre to these crops in 20-inch rows (or two rows per bed). Use proportional amounts for other row spacings not to exceed 4 1/2 pints of Lorsban 4E-HF per acre. Do not make more than one application per season or apply within 30 days before harvest.
Broccoli Cabbage	Root aphid	1.2 fl oz/1000 linear ft of row for single row plantings, and 2.4 fl oz/1000 linear ft of row for double row plantings.	Apply Lorsban 4E-HF in a water emulsion or with liquid fertilizer injected as a sidedress on each side of the row after plants are established. Avoid mechanical damage to crop roots. Use a minimum of 15 gallons of total spray volume per acre.	
Radishes	Root maggot	1.0 fl oz/1000 linear ft of row	Apply the specified dosage as a water-based drench in the seed furrows with the seed at planting time. Use a minimum of 40 gallons of total drench per acre.	Do not apply more than 5 1/2 pints of Lorsban 4E-HF per acre or make more than one application per season.
Rutabagas	Root maggot	1.6 - 3.3 fl oz/1000 linear ft of row	Apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time, behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Use a minimum of 40 gallons of total spray volume per acre.	Do not apply more than 4 1/2 pints of Lorsban 4E-HF per acre or make more than one application per season. Do not use rutabaga tops for feed or feed purposes.

except Brussels sprout,

Insert 6.

Lorsban 4E-HF

Brussels Sprouts

Use Lorsban 4E-HF to control cabbage aphid, cutworms, imported cabbageworm, and striped flea beetle adult by application as a full-coverage leaf spray at the rate of 1 quart per acre in 20 to 100 gallons, per acre using conventionally power-operated spray equipment. Apply when insects appear on the foliage and at 7 to 14 day intervals thereafter or as needed. Consult your state agricultural experiment station, extension service specialist, or registered pest control adviser for proper time to treat in your area.

To avoid phytoxicity in vegetable crops, do not mix with other pesticides or treat plants that are under extreme heat and drought stress.

See insert 8.

Label Revisions:

1. Added the phrase "Contains aromatic petroleum distillates" to ingredients section.
2. Added the phrase "Lorsban 4E-HF May cause skin sensitization reactions in certain individuals" to "Warning and Hazards" section.
3. Mint added to crops for application with sprinkler irrigation.
4. Agri-Mek added to tank mixes for citrus fruits.
5. Modified restrictions for citrus orchard floors.
4. Added "Corn grown for seed" to corn uses.
6. Cotton uses:
 - Growing areas including Arizona and California - "Bollworm" and "tobacco budworm" replace "Bollworms (Heliothis spp.)".
 - Number of applications to cotton limited to 5 per season.
 - All tank mix uses for cotton deleted.
7. Tree fruits: oblique banded leafroller added (apples); recommendations on use rates and spray volumes refined.
8. Pecans and Almond Orchard Floors: Restrictions edited.
9. Vegetables: Use directions on brussels sprout edited and put into a table format (changes based on previously approved label changes inadvertently omitted).

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unwanted consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, windless, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (excluding claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought; or
2. Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is properly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

DowElanco

Indianapolis, IN 46206, U.S.A.
 *Trademark of DowElanco

Insert 1.

Pests	Lorsban 4E-HF
Cotton fleahopper	3/8 - 1 pint/acre
Plant bugs (Lygus, Mirids)	
Cotton aphid	1/2 - 1 pint/acre
Fall armyworm	
Grasshoppers	
Thrips	
Yellow-striped armyworm	
Spider mites	1 pint/acre
Beet armyworm	1-1/2 - 2 pints/acre
Cotton bollworm	
Tobacco budworm	
Cutworms	
Pink bollworm	
Salt marsh caterpillar	

BEST AVAILABLE COPY

Insert 2.

Pests	Lorsban 4E-HF
Armyworms	1 - 2 pints/acre
Cotton aphid	
Cotton fleahopper	
Lygus	
Salt marsh caterpillar	
Thrips	2 pints/acre
Cotton bollworm	
Tobacco budworm	
Boll weevil	
Cutworms	
Pink bollworm	

Insert 3.

Lorsban 4E-HF may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed pests. For best results, use the recommended rate of Lorsban 4E-HF per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" on page 6 for further information.

Insert 4.

Use Lorsban 4E-HF as a dormant or delayed dormant spray at the rates indicated to control the following insects on the crops listed. While Lorsban 4E-HF may be used without oil, oil is recommended to control additional pests such as European red mite.

Crop	Insect	Lorsban 4E-HF per 100 gallons of Spray ¹
Apples	Rosy apple aphid San Jose scale <i>Lygus</i> Pandemis leafroller Climbing cutworms Oblique banded leafroller	1/2 - 1 pint (Use a minimum of 1.5 pints/acre)
Pears	San Jose Scale Climbing cutworms Pear psylla adults	
Plums, Prunes	San Jose scale Mealy plum aphid Climbing cutworms Peach twig borer	
Almonds, Peaches Nectarines	San Jose scale Peach twig borer Climbing cutworms	

BEST AVAILABLE COPY

¹Based on 200 to 600 gallons per acre as a dilute spray.

Insert 5.

For dilute sprays, tank mix the specified dosage with 1 to 2 gallons of petroleum spray oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using suitable ground spray equipment. (See "Additional Precautions Specific to California" (below) for use in California).

For low volume (concentrate) sprays, less than 200 gallons of spray mixture per acre, use the same amount of Lorsban 4E-HF as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of Lorsban 4E-HF for severe infestations. Use oil as recommended by your state agricultural experiment station or extension service specialist.

Precautions

Because cold or dry conditions may cause Lorsban 4E-HF oil sprays to injure trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated. Do not use more than 4 pints of Lorsban 4E-HF per acre.

Insert 6.

Crop	Insects Controlled	Dosage Lorsban 4E-HF	Use Directions	Restrictions
Brussels sprouts	Amyworms Cabbage aphid Cutworms Imported cabbageworm Striped flea beetle (adult)	1 to 2 pints per acre	Apply Lorsban 4E-HF with conventional power-operated spray equipment in 20 to 150 gallons of water per acre. Apply when insects appear on foliage and at 7 to 14 day intervals thereafter as needed. Consult your state agricultural experiment station, extension service specialist, or integrated pest control advisor for proper time to treat in your area.	Do not make more than 6 applications per season. Do not apply within 21 days before harvest.