

Dowelanco Brush & Weed Herbicide

Dow Elanco  
Quad IV  
9002 Purdue Road  
Indianapolis, IN 46268-1189

JUN 24 1992

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data or other material required for registration/ reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phrase "EPA Registration No. 62719-226."
  - b. Submit an acceptable inhalation and sensitization studies if the sensitization waiver request and inhalation rebuttal, which have not been reviewed yet, are denied. Please note that further label revisions may be required based on the review of the information you have submitted.

64682:I:WP50:Allen:WP25-5:KEVRIC. 06/19/92:07/18/92

295

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Robert J. Taylor  
Product Manager (25)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosures

395

(logo) DowElanco


### DowElanco Brush & Weed Herbicide

Controls Woody Plants and Broadleaf Weeds  
Contains Hammer® Herbicide

Active Ingredient(s):  
triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), as the triethylamine salt\*\* ..... 8.3%  
Inert Ingredients ..... 91.2%  
Contains 0.6 pounds of active ingredient per gallon.  
\*\*Equal to 6.36% Triclopyr acid.

\*Trademark of DowElanco

EPA Reg. No. 62719-XXX  
EPA Est. No.



Keep out of reach of children

### CAUTION

#### Precautionary Statements

Hazards to Humans and Domestic Animals

**Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin**

Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Do not allow children or pets onto treated areas until sprays have dried. Remove contaminated clothing and wash before reuse.

#### First Aid

**If In Eyes:** Immediately flush eyes with plenty of water for 15 minutes. Call a physician if irritation persists.

**In On Skin:** Remove contaminated clothing and wash skin with soap and water. If irritation continues, get medical attention.

**If Swallowed:** Call a physician. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL BEFORE USING.**

Read entire label before use.

Use strictly in accordance with label precautionary statements and directions.

#### STORAGE AND DISPOSAL

**Storage:** Store product in original container and keep out of reach of children. Store in a cool, dry place. Avoid contamination of food and foodstuffs.

**Container Disposal:** Do not reuse container. Rinse container thoroughly before discarding in trash. Securely wrap container in several layers of newspaper and discard in trash.

495

## General Information

DowElanco Brush and Weed Herbicide is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, buildings, trails, fences, walkways rights-of-way and other non-crop areas.

### **General Use Precautions**

Do not apply this product directly to, or allow to drift to grapes, vegetable crops, flowers, ornamentals or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift. Do not spray when the wind is blowing toward desirable plants. Do not apply with nozzles that produce a fine-droplet spray. Measuring utensils and spray equipment should not be reused for purposes other than herbicide application.

Do not use on or around edible food or food crops.

Do not graze treated areas.

### **Woody Plants and Vines Controlled Include:**

Alder	Dogwood	Raspberry
Arkansas Rose	Elderberry	Salmonberry
Arrowwood	Elm	Sassafras
Ash	Hazel	Scotch Broom
Aspen	Honeysuckle	Sumac
Beech	Hornbeam	Sweetbay Magnolia
Birch	Kudzu	Sweetgum
Blackberry	Locust	Sycamore
Blackgum	Madrone	Tanoak
Box Elder	Maples	Thimbleberry
California Rose	Mesquite	Trumper Creeper
Cascara	Mimosa	Tulip Poplar
Ceanothus	Mulberry	Virginia Creeper
Cherry	Oaks	Western Hemlock
Chinquapin	Persimmon	Willow
Choke Cherry	Pine	Winged Elm
Cottonwood	Poison Ivy	Wild Grape
Crataegus (Hawthorn)	Poison Oak	Wild Rose
Douglas Fir	Poplar	

### **Perennial Broadleaf Weeds Controlled Include:**

Bindweed	Field Bindweed	Smartweed
Burdock	Field Horsetail Rush	Spurge
Canada Thistle	Henbit	Tansy Ragwort
Chicory	Lambsquarter	Vetch
Clover	Oxalis	Wild Lettuce
Curly Dock	Plantain	Wild Violet
Dandelion	Ragweed	

## **Application Directions**

### **When to Apply**

For best results, apply as a full coverage foliar spray when brush and weeds are fully leaved out and actively growing. Wilting or other symptoms will occur in 1-6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

### **Application Methods**

#### **Foliage Spray**

Using a hose-end or tank sprayer, mix 8-16 tablespoons (4-8 fl. oz.) in enough water to make one gallon mixed spray. For mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine or winged elm), or during drought conditions, use the highest rate of application indicated. Resprouting may occur the year following treatment when lower rates are used. The addition of a surfactant (spreader/sticker) enhances herbicidal action and improves distribution of the spray on hard-to-wet plant surfaces. Apply at the rate of one gallon mixed spray to approximately 500 square feet of area. Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See "Use Precautions".

#### **Stump Treatment**

To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

#### **Vine Control**

When treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump treatment.

---

### **Warranty Information**

---

Subregistrant may include appropriate warranty and disclaimer information.

DowElanco • Indianapolis, IN 46268, U.S.A.

12/12/91