



COMPANY
NUMBER

62719

PRODUCT
SERIAL NO.

250

LABEL
APPROVAL
DATE

2.9.95

PM 23 62719-250

1 of 99



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

62719-
250

Date of Issuance:

FEB 09 1995

NOTICE OF PESTICIDE:

 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Treflan H.F.P.

Name and Address of Registrant (include ZIP Code):

DowElanco
9330 Zionsville Road
Indianapolis, IN 46268

Notes: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 62719-250".
- b. In the supplemental labeling entitled "Treflan for Weed Control in Rapeseed (Canola), Crambe, Safflower and Sunflower", add a precaution to read "Do not graze or harvest crambe for livestock forage."

Signature of Approving Official:

Date:

FEB 09 1995

204.97

c. In the supplemental labeling entitled "Special Use Program - Control of DNA-Resistant Goosegrass in Soybeans with Treflan Plus Dual Tank Mix", we have noted that the application rates for Dual have been crossed out. If this is a typographical error, the rates for Dual should be reinstated at 1.25 pints per acre on coarse soils, 1.5 pints per acre on medium soils, and 2 pints per acre on fine soils.

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

DK: 305-7546: FHB/PM23		CONCURRENCES					
SYMBOL	7505C						
SURNAME	D. KENNY						
DATE	12/7/95						

30297

[Editor's note: Added text denoted by underline and deleted text denoted by ~~strike-through~~.]

[Laminated Booklet -- Treflan H.F.P. Crop]
[Front Cover of Booklet]

ACCEPTED
with COMMENTS
in EPA Letter Dated
FEB 09 1995

(logo) DowElanco

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
62719-250

Treflan* H.F.P.

A selective herbicide for the preemergence control of annual grasses and broadleaf weeds.

Active Ingredient:

trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine.....	43.0%
Inert Ingredients	57.0%
Total	100.0%

Contains 4 pounds active ingredient per gallon.

Manufactured using the processes of U.S. Patents 4,120,905 and 4,226,789

Keep Out of Reach of Children

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), Engineering Controls Statements and User Safety Recommendations, and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Avoid Freezing -- Store Above 40°F

EPA Reg. No. 62719-ELN

EPA Est. 1471-IN-2

*Trademark of DowElanco

DowElanco • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents 2.5 gal

BEST AVAILABLE COPY

40297

[Page 2 of Booklet]

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment	-
Use Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
General Use Precautions	-
Rotation Crop Restrictions	-
Soil Texture Guide for Application Rates	-
Mixing Directions	-
Testing for Compatibility in Liquid Fertilizers	-
Application Methods	-
Ground Broadcast Application	-
Aerial Broadcast Application	-
Application with Dry Bulk Fertilizer	-
Application by Chemigation	-
Application Timing	-
Spring Application	-
Fall Application	-
Preemergence Application Immediately After Planting	-
Postemergence and Layby Application	-
Incorporation Directions	-
Soil Preparation and Incorporation	-
Incorporation Equipment	-
Weeds Controlled by Treflan H.F.P.	-
Special Use Programs	-
Approved Crops	-
Alfalfa (Established)	-
Almond (See Tree and Vine Crops section)	-
Apricot (See Tree and Vine Crops section)	-
Asparagus (Established)	-
Barley (See Small Grains section)	-
Beans	-
Beans - Guar and Mungbean	-
Beans - Lima Bean and Snap Bean	-
Broccoli (See Cole Crops section)	-
Brussels Sprouts (See Cole Crops section)	-
Cabbage (See Cole Crops section)	-
Canola (see Rapeseed)	-
Carrot	-
Castor Beans	-
Cauliflower (See Cole Crops section)	-
Celery	-
Chicory	-
Cole Crops (Broccoli, Brussels Sprouts, Cabbage and Cauliflower)	-
Collard (See Greens section)	-
Corn (Field Corn Only)	-
Cottonwood Trees Grown for Pulp	-
Cotton	-
Crambe (See Rapeseed)	-
Cucurbits	-

5897

Durum (See Small Grains section)
 Endive (See Chicory)
 Flax (Fall Application Only)
 Forage Legumes
 Grain Sorghum (Milo)
 Grape (See Tree and Vine Crops section)
 Grapefruit (See Tree and Vine Crops section)
 Greens (Turnip greens, Collard, Kale and Mustard Greens)
 Guar (see Beans - Guar and Mungbean)
 Hops
 Kale (See Greens section)
Kenaf
 Lemon (See Tree and Vine Crops section)
 Milo (See Grain Sorghum)
 Mustard (Grown for Seed or Processing for Food)
 Mustard Greens (See Greens section)
 Nectarine (See Tree and Vine Crops section)
 Okra
 Onions (Dry Bulbs Only)
 Orange (See Tree and Vine Crops section)
 Peach (See Tree and Vine Crops section)
 Peas (Dry and English Peas)
 Peas (Southern Peas)
 Peanuts
 Pecan (See Tree and Vine Crops section)
 Pepper (Transplant only)
 Plum (See Tree and Vine Crops section)
Potatoes
 Prune (See Tree and Vine Crops section)
 Radish
Rapeseed (Canola) and Crambe
 Safflower
 Small Grains (Barley, Durum and Wheat)
 Barley, Spring Seeded - Spring Application Preplant
 Incorporated for Foxtail (Pigeongrass) Control (For Use
 in Minnesota, North Dakota and South Dakota)
 Barley, Spring Seeded - Spring Application Preplant
 Incorporated for Foxtail (Pigeongrass) Control in Barley
 Used as a Cover Crop or in the Conservation Reserve Program
 Winter Wheat - Preplant Incorporated for Control of Cheatgrass
 and Other Annual Grasses and Broadleaves (For Use in
 Idaho, Oregon and Washington)
 Winter Wheat - Post Plant Incorporated (For Use in Idaho,
 Oregon and Washington)
 Winter Wheat - Fallow Soil Application Prior to Planting
 (For Use in Idaho, Oregon and Washington)
 Wheat, Durum and Barley - Spring Seeded - Fall Applied Preplant Soil
 Incorporated for Foxtail (Pigeongrass) Control (For Use in Minnesota,
 North Dakota, and South Dakota)
 Spring Wheat, Durum and Barley - Post Plant Incorporated
 for Foxtail (Pigeongrass) Control
 Sorghum (see Grain Sorghum)
 Soybeans
 Sugar Beets
 Sugarcane
 Sunflower
 Tangelo (See Tree and Vine Crops section)
 Tangerine (See Tree and Vine Crops section)

6 of 97

Tomato	-
Tree and Vine Crops - Citrus, Fruit and Nut Trees and Vineyards:	-
New Plantings of Citrus Fruit and Nut Trees	-
New Plantings of Vineyards	-
Established Non-bearing and Bearing Citrus, Fruit and Nut Crops	-
Trees and Vineyards	-
Wheat (See Small Grains section)	-
Walnut (See Tree and Vine Crops section)	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

The following new supplemental labels are being submitted for EPA approval:

1. Fall Postplant Soil Incorporated Application for Suppression of Common Windgrass in Winter Wheat (For Distribution and Use Only in the State of Michigan)
2. The Coulter Caddy as an Option for the Second Incorporation Pass for Trellan in Soybeans Grown Under Reduced or Conservation Tillage Conditions
3. Preplant Incorporated Treatment for Weed Control in Chinese Cabbage or Kohlrabi
4. Weed Control in Eggplant
5. Preplant Incorporated Treatment for Weed Control in Lentils (For Distribution and Use Only in the State of North Dakota)

The following supplemental labels were previously accepted for Trellan E.C. (62719-93) by EPA and are attached to make this labeling totally inclusive of all labeled uses for Trellan H.F.P. (Note: These labels contain minor editing to correct (one or more may apply): reference to "Trellan" as a DowElanco trademark; grammar; punctuation; and use of upper/lower case.) Because of their insignificance, these changes are not indicated by underlining and strike-through notations. The first four (4) supplemental labels in the following list contain additional significant changes (added text is underlined and deleted text denoted by ~~strike-through~~):

1. Fall Application to Fine Textured Soils for Weed Control in Cotton to be Grown the Following Year (EPA-accepted 10/23/91)
2. Spring Applied Trellan for Foxtail (Pigeongrass) Control in Spring Seeded Barley Grown Under Irrigation (For Distribution and Use Only in the State of Montana) (EPA-accepted 10/23/91)
3. Trellan plus Pursuit Combinations for Weed Control in Spanish Peanuts or Florunner and Florigiant Varieties (EPA-accepted 10/23/91)
4. Trellan for Weed Control in Rapeseed, (Canola), Crambe Safflower and Sunflower (For Distribution and Use Only in the State of Montana) Note: Crambe being added to this previously accepted supplemental label as per EPA-accepted supplemental labeling dated 03/30/93.
5. Trellan Plus Tri-Scept Tank Mix for Control of Fall Panicum, Seedling Johnsongrass and Shattercane in Soybeans (EPA-accepted 12/06/90)
6. Trellan Plus Canopy and Crop Oil Concentrate Early Preplant Application for Weed Control in No-till Soybeans (Not For Use in California) (EPA-accepted 12/06/90)
7. Trellan 48 Hour Incorporation Delay (For Distribution and Use Only in Texas, Oklahoma and New Mexico) (EPA-accepted 03/20/91)
8. Trellan for Partial Control or Suppression of Annual Brome Species (Cheatgrass, Downy Brome, Japanese Brome, Hairy Chess) and Jointed Goatgrass in Winter Wheat in Colorado, Kansas, Nebraska, Oklahoma, South Dakota, Texas and Wyoming (EPA-accepted 09/01/89)
9. Trellan Applied in Irrigation Water Rings for Weed Control in Non-bearing Citrus Trees (EPA-accepted 10/23/91)
10. Trellan plus Scepter Combinations for Weed Control in Soybeans (Not for Use in California) (EPA-accepted 10/23/91)
11. Trellan 72 Hour Incorporation Delay (For Distribution and Use Only in the States of Arizona and California) (EPA-accepted 06/29/92)
12. Trellan Application Over Standing or Shredded Cotton Stalks After September 1 for Weed Control in Cotton (For Distribution and Use Only in the State of California) (EPA-accepted 06/29/92)

7897

13. Incorporation with the Springtooth Harrow on Coarse Textured Soils to be Bedded up Prior to Planting ((For Distribution and Use Only in the State of Texas) (EPA-accepted 06/29/92)
14. Weed Control in Potatoes (Not for Use in the State of Maine) (EPA-accepted 02/03/93)
15. Use Rates for Fall Application Prior to Planting Cotton or Soybeans in the Spring (For Use Only in Arkansas, Louisiana and Mississippi) (EPA-accepted 01/25/93)
16. Treflan Plus Atrazine Tank Mix for Weed Control in Field Corn ((For Distribution and Use Only in the States of Alabama, Florida, Georgia and Texas) (EPA-accepted 06/22/93)
17. Green Foxtail (Pigeongrass) Resistance to Dinitroaniline Herbicides Including Treflan, Identified in the State of North Dakota (For Distribution and Use Only in the State of North Dakota) (EPA-accepted 04/20/94)
18. Special Use Program - Control of DNA-Resistant Goosegrass in Soybeans with Treflan Plus Dual Tank Mix (EPA-accepted 04/15/93)
19. Special Use Program - Enhanced Control of Broadleaf Signalgrass in Soybeans with Treflan Plus Dual Tank Mix (EPA-accepted 05/18/93)
20. Postemergence Soil Incorporated Application for Weed Control in Cotton (For Distribution and Use in Texas, Oklahoma and New Mexico) (EPA-accepted 08/24/94)

8097

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin • Trellan H.F.P. May Cause Skin Sensitization Reactions In Certain Individuals

Avoid breathing vapors or spray mist and contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate or Viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonsful) with a large quantity of water. Do not induce vomiting or give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation assistance, and get medical attention immediately.

If on skin: Immediately wash with plenty of soap and water. Get medical attention if irritation develops.

Note to physician: This product contains an aromatic hydrocarbon and can be extremely harmful if swallowed. Aspiration of this product may produce a severe pneumonitis. Stomach lavage with a cuffed endotracheal tube in place and immediate administration of activated charcoal, 6-8 heaping teaspoonsful with water, should be considered. Treatment is otherwise symptomatic and supportive.

9597

Environmental Hazards

This pesticide is toxic to fish. ~~For terrestrial uses, do not~~ Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use before applying this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as Barrier Laminate or Viton
- Shoes plus socks
- Protective eyewear

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. Do not store near heat or flame. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Plastic Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Bulk/Mini-bulk Tank Cleaning: Triple rinse (or equivalent) and wash with appropriate cleaners before reusing.

10897

General Information

Treflan® H.F.P. herbicide is a selective herbicide for the preemergence control of annual grasses and broadleaf weeds. Treflan H.F.P. may be applied in liquid sprays of water or liquid fertilizer, or impregnated on dry bulk fertilizer. To prevent reduce loss of herbicidal activity, Treflan H.F.P. must should be soil incorporated within 24 hours after application unless otherwise specified in specific use directions or supplemental labeling. Treflan H.F.P. may be tank mixed or followed by overlay or postemergence treatments with other herbicides to improve the spectrum of weeds controlled. Treflan H.F.P. controls weeds by disrupting growth processes during germination. Treflan H.F.P. does not control established weeds.

General Use Precautions

Applied according to directions and under normal growing conditions, Treflan H.F.P. will not harm the treated crop. Over-application may result in crop injury or rotational crop damage from soil-residue herbicide carryover. Uneven application or improper incorporation of Treflan H.F.P. can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Treflan H.F.P. Under these conditions, delayed crop development or reduced yields may result.

Do not apply Treflan H.F.P. to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.

Do not use Treflan H.F.P. on any crop grown in Pecos county or Reeves county, Texas.

In Montana, uses of Treflan H.F.P. are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Chemigation: Treflan H.F.P. may be applied by chemigation on certain crops. See instructions for chemigation in the "Application Methods" section of this label. Also, see specific instructions for certain crops in the "Approved Crops" section of this label.

Rotation Crop Restrictions

Sugar beets, Red beets and Spinach

In Arizona, Colorado, California, Idaho, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming: Sugar beets, red beets or spinach should not be planted for 12 months after a spring application or 14 months after a fall application of Treflan H.F.P. Moldboard plowing to a depth of 12 inches prior to planting these crops will reduce the possibility of crop injury. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application of Treflan H.F.P.

In all other areas: Sugar beets, red beets, and spinach should not be planted for 12 months after a spring application or 14 months after a fall application. Before planting sugar beets, moldboard plow to a depth of 12 inches to reduce the possibility of crop injury.

Proso Millet, Corn, Sorghum (Milo), Oats and Annual or Perennial Grass Crops or Grass Mixtures

In Arizona, Colorado, California, Idaho, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming: Unless crop injury is acceptable, proso millet, corn, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 12 months after a spring application or 14 months after a fall application of Treflan H.F.P. to avoid the possibility of crop injury. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or

11897

20 months after a fall application. Moldboard plowing to a depth of 12 inches before planting these crops will reduce the possibility of crop injury.

In Minnesota, North Dakota and South Dakota: Unless crop injury is acceptable, proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 18 months after a spring application or 21 months after a fall application of Treflan H.F.P.

In those portions of Kansas, Nebraska, Oklahoma and Texas that receive less than 20 inches of rainfall and irrigation to produce a crop: Unless crop injury is acceptable, do not plant proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures for 18 months after an application of Treflan H.F.P. In sorghum, cool wet weather conditions during early growth stages may increase the possibility of crop injury. In areas receiving more than 20 inches of rainfall and irrigation, these crops should not be planted for 12 months after a spring application or 14 months after a fall application of Treflan H.F.P.

All other areas receiving more than 20 inches of rainfall and irrigation: these crops should not be planted. Unless crop injury is acceptable, do not plant proso millet, sorghum (milo), oats and annual or perennial grass crops or grass mixtures for 12 months after a spring application or 14 months after a fall application of Treflan H.F.P.

Other Crops

Vegetable crops, other than those listed on this label for use with to which Treflan H.F.P. may be applied as a preplant soil incorporated treatment, application of Treflan H.F.P. should not be planted within 5 months after an application of Treflan H.F.P.

Soil Texture Guide for Application Rates

Rate recommendations for incorporated treatments of Treflan H.F.P. are based on "Soil Texture Class" (coarse, medium or fine) and soil organic matter content. A fine textured soil (e.g., clay loam) will require a higher application rate than a coarse textured soil (e.g., loamy sand). In the table below, find the "Soil Texture Class" (coarse, medium or fine) corresponding to the "Soil Texture to be Treated". Choose the proper rate for each application based on the "Soil Texture Class" and specific crop recommendations. Do not exceed recommended rates.

Soil Texture Class	Soil Texture to be Treated
Coarse (Light) Soils	Sand, loamy sand, sandy loam
Medium Soils	Loam, silty clay loam†, silt loam, silt, sandy clay loam†
Fine (Heavy) Soils	Clay, clay loam, silty clay loam†, silty clay, sandy clay, sandy clay loam†

†Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt, they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

Mixing Directions

Treflan H.F.P. - Alone

Treflan H.F.P. may be mixed with water or most liquid fertilizer materials. Prior to mixing Treflan H.F.P. in liquid fertilizer, refer to the label section entitled "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine compatibility with the liquid fertilizer product to be used. The combination of Treflan H.F.P. with solution and suspension-type fertilizers provides weed and grass control equal to water sprays.

Fill spray tank 1/3 to 1/2 full with clean water or liquid fertilizer. Start agitation. Add correct amount of Treflan H.F.P. and continue agitation while filling tank to required spray volume.

12897

Precaution: Do not allow water or spray mixture to back-siphon into a water source.

Treflan H.F.P. in Tank Mix

For broader spectrum weed control, Treflan H.F.P. may be tank mixed with other products registered for use on crops listed in this label, unless tank mixing with Treflan is prohibited by the manufacturer's label. Use application rates recommended for "Treflan H.F.P. - Alone". Consult the tank mix product label for additional weeds controlled and applicable directions for use, before applying. Follow all precautions and limitations imposed by the label for each product used in the tank mixture.

Treflan H.F.P. may be tank mixed with other products and applied with water or most liquid fertilizer materials. Prior to mixing tank mixes containing Treflan H.F.P. with liquid fertilizer, refer to label section entitled "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine tank mix compatibility with the liquid fertilizer product to be used.

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture. To prevent foaming during filling, keep end of fill pipe below the surface of the liquid in the spray tank.

Mixing Order: Fill the spray tank to 1/4 to 1/3 of the total spray volume required. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: Dry flowables (DF); wettable powders (WP); aqueous suspensions (AS), flowables (F) and liquids (L).

Maintain agitation and fill spray tank to 3/4 of total spray volume. Add Treflan H.F.P. and other emulsifiable concentrates (EC) and any solutions (S).

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Precautions:

Read and carefully follow all label instructions for each material added to the spray tank. Do not allow water or spray mixture to back-siphon into a water source.

Premixing: Dry and flowable formulations may be premixed with water (slurried) and added to the spray tank through a 20-35 mesh screen. This procedure assures good initial dispersion of these products in liquid fertilizer or water.

Line screens in the spray tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

Testing for Compatibility in Liquid Fertilizers

Treflan H.F.P. alone or in tank mix combination with dry flowables (DF), wettable powders (WP), aqueous suspensions (AS), flowables (F), liquids (L), or solutions (S) may not combine properly with some liquid fertilizer materials. Small quantities should always be tested before full scale mixing. Follow the testing procedure below to determine if a compatibility agent is needed. If required, use only a phosphate ester-type surfactant designed for use with liquid fertilizers. Such compatibility agents can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. Add the compatibility agent just before adding pesticides, and which one works best in your herbicide/fertilizer mixture. The seven compatibility agents listed at the end of this section have been thoroughly tested. Other surfactants commercially available may or may not be suitable for use with liquid fertilizers.

13891

Testing Procedure

1. Add 1 pint of the liquid fertilizer to a quart jar.
2. Add 1 to 4 teaspoon(s) of the DF, WP, AS, F, or L formulation (depending on mixing ratio required) to the liquid fertilizer. Close the jar and agitate until the materials are evenly dispersed in the liquid fertilizer. If the materials do not disperse well, it may be necessary to slurry the chemicals in water before adding to the fertilizer.
3. After dispersing the materials (step 2) add 3 to 4 teaspoons of Treflan H.F.P. and other EC formulations to the jar and shake well. Add solution herbicides to the mixture last and agitate. Observe the jar for about 10 minutes. If materials rise to the surface and form a thick layer (oily curds) that will not redisperse when agitated, a compatibility agent is needed. If the mixture is easily redispersed with slight agitation, a compatibility agent is not required. Good agitation, however, must be provided to maintain dispersion in the spray tank.
4. If the need for a compatibility agent is demonstrated (step 3) the following procedure is recommended: Using a clean quart jar repeat step one above and add 1/2 teaspoon of the compatibility agent to the liquid fertilizer. Mix well and then repeat steps 2 and 3.

An effective compatibility agent will cause the mixture to remain uniformly dispersed with little or no separation (oil rising to the surface) for one half hour or longer. If slight separation occurs, 2 to 3 inversions of the jar should be sufficient to uniformly redisperse the mixture. If oily curds form and will not redisperse, additional compatibility agent or an alternative compatibility agent should be tried.

Use a clean jar for each test. A compatible mixture will have a uniform appearance and will be relatively easy to redisperse with gentle agitation of the jar.

Compatibility Agents

The phosphate ester type surfactants listed below are designed for use with liquid fertilizers, and can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. Add the compatibility agent just before adding pesticides.

1. Sponto 168D (Witco Chemicals Co., Chicago, IL)
2. Compat (Farm Chemicals, Inc., Aberdeen, NC) (Not for use in California)
3. Unite (Hopkins Ag Chemical, Madison, WI)
4. T-Mulz 734-2 (Thompson-Hayward Chemical Co., Kansas City, MO) (Not for use in California)
5. Rigo Compatibility Agent (Rigo Company, Buckner, KY)
6. Amoco Spray Mate (Amoco Oil Co., Chicago, IL) (Not for use in California)
7. Kem-Link (Universal Coop, Minneapolis, MN)

Note: Compliance with state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer or chemical mixture for sale.

Precautions:

Do not use the compatibility agents listed above for tank mixes in plain water. Read the compatibility agent label for use directions and precautions before use.

Application Methods**General**

As spray volume decreases, the importance of accurate calibration and uniform application increases. Check calibration and uniformity of spray application daily. To avoid spray drift, do not apply when winds are gusting or when wind speed is greater than 15 mph.

Ground Broadcast Application

Apply Treflan H.F.P. in 5 to 40 gallons of liquid carrier per acre (broadcast basis), using any properly calibrated, low pressure herbicide sprayer that will apply the spray uniformly. The carrier may be water or liquid fertilizer as specified for the crop to be treated in the "Approved Crops" section of this label. For

14897

band application, adjust herbicide rate and spray volume in proportion to the band width and row width treated.

Aerial Broadcast Application

Apply Treflan H.F.P. in 5 to 10 gallons of water per acre. Adjust pump pressure, nozzle arrangements, speed and application height to provide uniform application to the soil surface. Use swath markers or flaggers to assure proper swath width interval.

Application with Dry Bulk Fertilizer

Dry bulk fertilizers impregnated or coated with Treflan H.F.P. may be applied as a preplant incorporated treatment on approved crops. All label recommendations for Treflan H.F.P. regarding application rates, incorporation directions, special instructions and precautions should be followed. Read and follow all label instructions below concerning use of Treflan H.F.P. with dry bulk fertilizer. Properly applied dry bulk fertilizers impregnated with Treflan H.F.P. provides weed and grass control equal to water sprays.

Use the following formula to calculate the amount of Treflan H.F.P. required to impregnate a ton of dry bulk fertilizer.

$$\begin{array}{rcccl} \text{Pints Treflan H.F.P.} & & 1000 & & \text{Quarts Treflan H.F.P.} \\ \text{Per Acre} & \times & \text{Pounds Fertilizer} & = & \text{Per Ton of} \\ & & \text{Per Acre} & & \text{Fertilizer} \end{array}$$

Limitations: Apply a minimum of 200 lb/acre of dry fertilizer impregnated with Treflan H.F.P. at the recommended broadcast rate per acre. Any commonly used dry fertilizer can be used for impregnation of with Treflan H.F.P. except coated ammonium nitrate and pure limestone. These materials will not absorb the herbicide. Blends containing mixtures of these materials can be impregnated.

Impregnation: Use any closed drum, belt, ribbon or other commonly used dry bulk fertilizer blender. Nozzles used to apply Treflan H.F.P. to dry bulk fertilizer should be placed to provide uniform spray coverage.

Application and Incorporation: Spread the fertilizer/chemical mixture with properly calibrated application equipment. Be certain the material is applied uniformly to the soil surface. Treflan H.F.P. should be incorporated 2 times when impregnated on dry bulk fertilizer. Dry bulk fertilizer impregnated with Treflan H.F.P. must be incorporated 2 times. The first incorporation should occur within 24 hours after application. The second incorporation should be delayed 2 to 5 a minimum of 5 days after the first and be completed prior to planting.

Compliance with State Regulations: Compliance with state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer or chemical mixture for sale.

Application by Chemigation

Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in certain crops as specified in "Approved Crops" section of this label. Read and follow all label instructions outlined below concerning chemigation before applying Treflan H.F.P. by this method.

Note: Do not apply Treflan M.T.F. or Treflan 5 through any type of irrigation system.

General Chemigation Directions:

Apply this product only through continuously moving center pivot, lateral move, or end tow sprinkler irrigation systems equipped for chemigation. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of chemigation treated water.

15897

If you have questions about calibration you should contact state extension specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Sprinkler Chemigation Directions:

The following directions must be followed for all recommended sprinkler irrigation systems (center pivot, lateral move or end tow):

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point that pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Trellan H.F.P. should be injected continuously throughout the chemigation period. The chemigation metering pump should be checked periodically during application to insure proper operation.
9. The injection metering pump must be calibrated as specified by the manufacturer.
10. Pesticide injection hoses which connect chemigation metering equipment to the sprinkler irrigation system should be of braided reinforced construction with an internal tube made of nylon, cross-linked polyethylene or high density polyethylene.
11. Trellan H.F.P. may cause staining of plastic hoses and tanks.
12. Apply Trellan H.F.P. in sprinkler irrigation equal to 1/2 to 1 inch of water.
13. During chemigation, maintain agitation in supply tank at all times.

[Editors note: Item 10, above, added from previously accepted supplemental labeling (EPA-accepted 02/24/92).]

Chemigation System Calibration:

Sample calculation for use of Trellan H.F.P. in a chemigation system:

- Assume, in this example, 133 acres are to be covered by a chemigation treatment.
- Product required, assuming 1.5 pints per acre is 199.5 pints
(133 acres X 1.5 pt/acre = 199.5 pt = 25 gallons)
- Add 25 gallons of product directly to the injection supply tank.
- Adjust the injection system to deliver 25 gallons during the time required to apply 1 inch of water to 133 acres.

If the irrigation system requires 20 hours to apply 1 inch of water to 133 acres, the injection rate is 1.25 gal/hr and is calculated as follows:

- 25 gal ÷ 20 hr = 1.25 gal/hr
- 1.25 gal/hr = 160 fl oz/hr

16897

Proper calibration requires the injection pump to be adjusted to deliver 2.7 fl oz/min. and is calculated as follows:

$$\bullet 160 \text{ fl oz/hr} \div 60 \text{ min/hr} = 2.7 \text{ fl oz per min.}$$

Chemigation Mixing Directions:

Undiluted Treflan H.F.P.: When used alone, the injection of undiluted Treflan H.F.P. is recommended in chemigation systems. For undiluted use, the metering pump, supply tank, and any associated equipment must be thoroughly clean and dry before Treflan H.F.P. is added to the system for injection. When injecting undiluted Treflan H.F.P., maintain continuous agitation in the supply tank.

Diluted Treflan H.F.P.: Treflan H.F.P. may be diluted if required to achieve accurate calibration for existing equipment. Partially fill the injection supply tank with a volume of water equal to the amount of Treflan H.F.P. required (Do not add water to Treflan H.F.P.). Start agitation. Add the required amount of Treflan H.F.P. to water in the supply tank and continue mixing while filling the tank to the final volume required by the injection pump calibration. When injecting diluted Treflan H.F.P., maintain continuous agitation in supply tank.

Application Timing

Spring Application

Apply and incorporate Treflan H.F.P. any time after January 1 when soil can be worked and is in a condition which allows thorough mixing to insure uniform incorporation. See "Approved Crops" section for application timing recommendations for specific crops.

Fall Application

Fall application can be used for all crops for which Treflan H.F.P. is recommended as a preplant incorporated treatment. Refer to "Approved Crops" section for any crop specific fall application instructions.

In the states of California, North Dakota, South Dakota and Minnesota, apply and incorporate Treflan H.F.P. any time between September 1 and December 31. In all other states, fall apply Treflan H.F.P. between October 15 and December 31.

Ground may be bedded up over winter. On bedded ground, reduce beds to desired height before planting, by moving some treated soil from beds into furrows. Where soil is left flat over winter, care should be taken not to turn up untreated soil during spring bedding operations. Destroy established weeds during seedbed preparation. Weeds established in furrows as a result of exposing untreated soil should be destroyed before planting. Fall application of Treflan H.F.P. is not recommended on fields which remain wet or are subject to periods of flooding.

Preemergence Application Immediately After Planting

Apply and incorporate Treflan H.F.P. immediately after planting and prior to crop germination. Adjust incorporation equipment so as to not disturb planted seed. Refer to the "Approved Crops" section of this label for crop specific instructions.

Postemergence and Layby Application

Apply and incorporate Treflan H.F.P. at the recommended rate to the established crop at or before the last cultivation. Required preharvest intervals for treatments with Treflan H.F.P. for certain crops are specified in the "Approved Crops" section of this label. Crop cover may prevent uniform soil coverage from over-the-top sprays. To avoid this problem, use drop nozzles or directed sprays to achieve uniform soil coverage.

17897

Incorporation Directions

Soil Preparation and Incorporation

Ground cover, such as crop residues or existing weeds, can interfere with uniform soil incorporation of Treflan H.F.P. A manageable level of ground cover will allow uniform incorporation into the top 2 to 3 inches of soil the final seedbed. Ground cover and crop residues, if excessive, should be reduced by appropriate soil tillage prior to application.

Break up clods using tillage equipment prior to application of Treflan H.F.P. Treflan H.F.P. must be incorporated within 24 hours after application unless otherwise specified on supplemental labeling. Non-uniform application may result in erratic weed control or crop injury. With most equipment and methods of application, a second incorporation is required and may occur any time before planting. The second incorporation should be in a different direction, and to avoid bringing untreated soil to the surface, should not be deeper than the first. Note: Two-pass incorporation is required for all special use programs unless otherwise specified.

General Soil Conditions: The soil surface should be smooth enough to allow for uniform application and efficient incorporation of Treflan H.F.P. Break up clods using tillage equipment prior to application of Treflan H.F.P. Apply when soil moisture is sufficient to allow the breakup of large clods and uniform mixing during the incorporation process. Soil compaction and/or non-uniform incorporation may occur if soil is excessively moist.

Incorporation in Bedded Culture: In bedded culture, Treflan H.F.P. should be incorporated to a depth of 2 to 3 inches in the final seedbed.

Application Prior to Bedding: Apply Treflan H.F.P. and incorporate one time with recommended equipment. The bedding operation serves as the second incorporation. Do not expose untreated soil during post-bedding operations such as planting since removal of treated soil during planting can allow weed germination and establishment in the drill row.

Application After Bedding: Knock off beds to planting height before applying Treflan H.F.P. Apply and incorporate Treflan H.F.P. with recommended equipment that will conform to the shape of the bed. Do not expose untreated soil.

Cultivation After Planting: Treated crops may be shallowly cultivated without reducing the weed control activity of Treflan H.F.P. Limit depth of cultivation to the zone of treated soil (2 to 3 inches) to avoid moving untreated soil to the surface. Exposure of untreated soil may cause loss of weed control.

Incorporation Equipment

Use incorporation equipment capable of mixing Treflan H.F.P. uniformly into the top 2 to 3 inches of the final seedbed. Use of inappropriate equipment or improper use of recommended equipment may result in erratic weed control and/or crop injury. Incorporation equipment such as a tandem disc will mix Treflan H.F.P. approximately half as deep as the equipment is set to operate. For example, a disc set to cut four inches deep will mix most of the Treflan H.F.P. within the top 2 inches of soil. Any recommended incorporation implement may be used alone or in combination with any other recommended implement. Two incorporation passes are required unless, otherwise specified as described below, soil conditions and available equipment are adequate to provide thorough soil mixing in a single pass.

Single Pass Incorporation Option

[Editor's note: This use added from previously accepted supplemental labeling (ERA, accepted 02/03/93)]

Treflan H.F.P. may be incorporated in a single pass if application conditions allow for thorough and uniform mixing into the top 2-3 inches of the final seedbed. Thorough and uniform incorporation may be achieved if the soil at the time of application is of good tilth with moderate moisture, and is relatively free of clods, and crop residue. Examples of types of equipment that can be used to obtain thorough and uniform soil mixing from a single pass are indicated by (†) in the following list:

18877

†**Finishing Disc** with disc blades no greater than 22 inches in diameter, spaced no more than 7 1/2 inches apart. Operate at 4 to 6 mph. Best results are obtained when the disc is equipped with harrow, reel or basket attachments.

Tandem Disc: Set equipment to cut 4 to 6 inches deep and operate at 4 to 6 mph.

†**Field Cultivator:** Set equipment to cut 3 to 4 inches deep and operate at a minimum of 5 mph. A field cultivator is defined as an implement with 3 to 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered with sweeps on successive rows staggered so that no soil is left unturned. Chisel points should not be used. Best results are obtained when the field cultivator is equipped with harrow, reel or basket attachments.

†**Combination Seedbed-Conditioners** implements: These implements are defined as three two or more tillage devices combined to operate as a single tillage unit. For example, 2 to 3 rows of field cultivator C- or S-shaped shanks with an effective sweep spacing of 6 to 9 inches successive rows of sweeps staggered so that no soil is left unturned, followed by a spike-tooth or flexline harrow, followed by a ground driven reel, or basket or incorporator wheels. Combination implements should be set to cut 3 to 4 inches deep and operated at a minimum of 6 mph. Treflan H.F.P. can be incorporated with one pass when using a combination seedbed conditioner. Two incorporations are required recommended under conditions which prevent optimum soil mixing such as excessive surface residue, roughness, high clay content or soil moisture soil is too wet or too dry. Combination tools can also be composed of 2 rows of wide crown sweeps that overlap so that the roots of all weeds and plants are severed. This should be followed by 2 gangs of rotating spoked wheels that thoroughly mix Treflan H.F.P. into the top 2-3 inches of the final seedbed.

Rolling Cultivator: Set equipment to cut 2 to 4 inches deep and operate at 6 to 8 mph. Generally, rolling cultivators are adequate for use only on coarse and medium soils. In sugarcane the rolling cultivator may be used on fine textured soils.

Bed Conditioner (Do-All): Set equipment to cut 2 to 4 inches deep and operate at 4 to 6 mph. One incorporation pass is adequate in bedded culture, while two incorporation passes are required in flat planted culture. The Do-All should be used only on coarse and medium textured soils.

Mulch Treader and other similar disc-type implements: Set equipment to cut 3 to 4 inches deep and operate at 5 to 8 mph.

†**P.T.O.-Driven Equipment (tillers, cultivators, hoes):** Adjust equipment to incorporate Treflan H.F.P. into the top 2 to 3 inches of the final seedbed with rotors spaced to provide a clean sweep of the soil. Only one incorporation is necessary. P.T.O. equipment should not be operated more than 4 mph.

Other Equipment: Other implements including the flexible tine-tooth harrow (Flexline or Melroe), are recommended, but only for certain uses defined in the "Approved Crops" section of this label.

Conservation Tillage Practices: In reduced or minimum tillage situations, fall or spring application and incorporation of Treflan H.F.P. may be combined with tillage operations. The first incorporation may utilize equipment such as a tandem disc, combination implement or bedding equipment that provides good soil mixing but leaves a maximum amount of crop residue on the soil surface. The second incorporation may be accomplished with tillage equipment that provides uniform soil mixing used in conjunction with no-till planters (See specific recommendations for reduced or conservation tillage situations for cotton and soybeans in the Approved Crops section.

19887

Weeds Controlled by Treflan H.F.P.**Grass Weeds****Common name****Scientific Name**

annual bluegrass

Poa annua

barnyardgrass

Echinochloa crus-galli

(watergrass)

brachiaria

Brachiaria spp.

(signalgrass)

bromegrass

Bromus tectorum

(cheatgrass)

(downy brome)

cheat

Bromus secalinus

(chess)

crabgrass

Digitaria spp.

(large crabgrass)

(smooth crabgrass)

foxtail

Setaria spp.

(bottlegrass)

(bristlegrass)

(giant foxtail)

(green foxtail)

(foxtail millet)

(pigeongrass)

(robust foxtail)

(yellow foxtail)

guineagrass

Panicum maximum

(See special instructions for control in sugarcane in the "Approved Crops" section.)

itchgrass

Rottboellia exaltata

(raoulgrass)

(See special instructions for control in sugarcane in the "Approved Crops" section.)

johnsongrass (from seed)

Sorghum halepense

(rhizome - see special instructions for control in cotton, soybeans, fruit and nut crops and vineyards in the "Approved Crops" section.)

junglerice

Echinochloa colonum

oats, wild†

Avena fatua

panicum

fall panicum

Panicum dichotomiflorum

(spreading panicgrass - see special instructions for control in cotton and soybeans in the "Approved Crops" section.)

ryegrass, Italian

Lolium multiflorum

(annual ryegrass)

Texas panicum

Panicum texanum

(buffalograss)

(Coloradograss)

red rice

Oryza sativa

(See special instructions for suppression or partial control in soybeans in the "Approved Crops" section.)

sandbur

Cenchrus incertus

(burgrass)

sprangletop

Leptochloa filiformis

stinkgrass

Eragrostis ciliaris

(lovegrass)

shattercane

Sorghum bicolor

(wild cane)

(See special instructions for control in soybeans in the "Approved Crops" section.)

woolly cupgrass

Eriochloa villosa

20897

[†]When applied as a preplant incorporated treatment, Trellan H.F.P. controls wild oats that germinate in the treated zone. Wild oat control is not claimed for incorporated uses in small grains.

Broadleaf Weeds

Common Name

Scientific Name

carpetweed

Mollugo verticillata

chickweed

Stellaria media

field bindweed

Convolvulus arvensis

(See special instructions for control in fruit and nut crops and vineyards in the "Approved Crops" section.)

goosefoot

Chenopodium hybridum

henbit

Lamium amplexicaule

knotweed

Polygonum aviculare

kochia

Kochia scoparia

(fireweed)

(Mexican fireweed)

lambsquarters, common

Chenopodium album

pigweed

Amaranthus spp.

(carelessweed)

(Palmer amaranth)††

(prostrate pigweed)

(redroot)

(rough pigweed)

(spiny pigweed)

(See special instructions for control in soybeans in "Approved Crops" section.)

puncturevine

Tribulus terrestris

(Western U.S. only)

(caltrop)

(goatweed)

purslane, common

Portulaca oleracea

pusley, Florida

Richardia scabra

(Florida purslane)

(Mexican clover)

(pusley)

Russian thistle

Salsola iberica

(tumbleweed)

stinging nettle

Urtica dioica

(nettle)

††Suppression only in areas of the Southwest U.S. where reduced sensitivity to trifluralin has been observed. Consult your local extension service or DowElanco representative for information regarding alternative weed control practices.

Special Use Programs

Trellan H.F.P. is approved for the following special use programs. Refer to "Approved Crops" section of this label for details on soil preparation, use rates, application, soil incorporation, and precautions for each type or program.

Cotton

• Chemigation

• Weed Control in Conservation Tillage

- Fall Panicum Control
- Pigweed and Seedling Johnsongrass Control
- Additional Weed and Grass Control (Gulf Coast Counties of Texas)
- Rhizome Johnsongrass Control

21897

Soybeans

• Chemigation

• Weed Control Under Reduced or Conservation Tillage

- Fall Panicum Control
- Pigweed and Seedling Johnsongrass Control
- Additional Weed and Grass Control (Gulf Coast Counties of Texas)
- Itchgrass (Beauvergrass) Suppression
- Charcoal Soils in Arkansas, Louisiana and Mississippi
- Red Rice Control in Arkansas, Louisiana, Mississippi and Texas
- Rhizome Johnsongrass Control in Eastern United States and the State of Texas
- Wild Cane (Shattercane) Control
- Treflan H.F.P. plus Sencor or Lexone for Rhizome Johnsongrass Control

Fruit and Nut Crops and Vineyards

- Rhizome Johnsongrass Control
- Field Bindweed Control

Approved Crops

ALFALFA - ESTABLISHED

Mechanically Incorporated

Apply Treflan H.F.P. with ground or aerial equipment and mechanically incorporate prior to weed emergence to control weeds listed in the "General Information" section of this label. Use mechanical incorporation equipment that will insure thorough soil mixing with minimal damage to crop stand.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5
Medium	2.0
Fine	2.0

Surface Applications (Chemigation or Water Incorporated)

Treflan H.F.P. may be surface applied for annual grass control in established alfalfa by chemigation, or ground or aerial broadcast application equipment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
All Soil Textures	4.0

Chemigation

Refer to "Application by Chemigation" section in the "General Information" section of this label for use directions for chemigation.

Surface Applications Activated by Rainfall or Irrigation

Broadcast surface applications of Treflan H.F.P. to established alfalfa may be activated by rainfall, sprinkler, flood or furrow irrigation. Rainfall or a single overhead sprinkler irrigation of 0.5 acre inch or more is required to activate Treflan H.F.P. If activated by furrow irrigation, care should be taken to thoroughly wet beds between furrows. If rainfall or irrigation has not occurred within 3 days after application, Treflan H.F.P. may be mechanically incorporated. If mechanically incorporated, use equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

22897

Application Timing and Weeds Controlled

Applications to established alfalfa for annual grass control can be made during dormancy or semi-dormancy, or during the growing season immediately after a cutting. Because Treflan H.F.P. does not control established weeds, application must be made prior to the expected time of weed germination. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather. To control these weeds, apply Treflan H.F.P. immediately after a cutting between August 1 and October 1, but prior to weed germination. When fall applied, Treflan H.F.P. controls bromegrass and cheat in addition to other labeled weeds that germinate after application.

The following weeds are controlled when Treflan H.F.P. is applied by chemigation or surface applied and incorporated by rainfall or irrigation:

barnyardgrass	crabgrass
bromegrass	cupgrass
(cheatgrass)	foxtail
(downy brome)	jungerice
(cheat)	sandbur
(chess)	wildbarley
canarygrass	

Precautions:

- Do not cut or graze alfalfa within 21 days after application of Treflan H.F.P.
- Apply no more than 4 pt of Treflan H.F.P. during any growing season. In the growing season following application of 4 pt of Treflan H.F.P. to alfalfa, plant only those crops for which Treflan H.F.P. is registered as a preplant treatment or crop injury may occur.

Tank Mix Combinations

Other products registered for use on established alfalfa may be applied ground-broadcast in tank mix combination with Treflan H.F.P. or applied as sequential treatments following application of Treflan H.F.P. Tank mixes containing Treflan H.F.P. must be applied by ground broadcast when alfalfa is dormant or semi-dormant, or immediately after a cutting.

Precaution: Refer to the tank mix product label for application rates, weeds controlled, additional use directions, precautions and limitations before use.

ASPARAGUS - ESTABLISHED

Apply Treflan H.F.P. to established asparagus as a single or split application. Treflan H.F.P. will suppress volunteer seedling asparagus and field bindweed when applied as directed. Follow recommended soil preparation, application and incorporation procedures for Treflan H.F.P.

Application Timing

Make applications to dormant asparagus in winter or early spring after mature ferns have been removed. Do not apply after new spears begin to emerge. Apply post-harvest applications immediately after harvest in late spring or early summer just before ferns are allowed to develop.

23997

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	
	Split Application	Single Application
	Before and After Harvest	Before or After Harvest
	(pints)	(pints)
Coarse	1.0 + 1.0	2.0
Medium	1.5 + 1.5	3.0
Fine	2.0 + 2.0	4.0

- Do not apply more than 2.0 pt/acre on coarse soils, 3.0 pt/acre on medium soils or 4 pt/acre on fine soils during any calendar year.

BEANS - ALL DRY AND FRESH BEANS/PEAS (EXCEPT BEANS/PEAS LISTED ELSEWHERE ON THIS LABEL)

Treflan H.F.P. - Alone

Apply and incorporate Treflan H.F.P. in the spring before planting or in the fall in advance of spring planting. See instructions for fall application of Treflan H.F.P. under the heading "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.25 1.0-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Treflan H.F.P. plus Eptam Tank Mix

Treflan H.F.P. may be tank mixed with Eptam 7E and applied as a preplant incorporated treatment to control additional weeds. Use application rates recommended for dry beans "Treflan H.F.P. - Alone", above. Refer to the label for Eptam for application rates, additional use directions, precautions and limitations before use.

Tank Mixing

For broader spectrum weed control, other products registered for use in dry and fresh beans/peas may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

24897

BEANS - GUAR AND MUNGBEAN

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.5
Fine	1.5

• All soils with 2-5% organic matter - 1.5 pints

BEANS - LIMA BEAN AND SNAP BEAN

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.0
Fine	1.5

• All soils with 2-5% organic matter - 1.5 pints


CARROT

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.



25 of 97

CASTOR BEAN

Apply Treflan H.F.P. as a soil incorporated treatment, before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CELERY

Apply Treflan H.F.P. as a soil incorporated treatment. Treflan H.F.P. may be applied to direct seeded or transplant celery before planting, at planting or immediately after planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

— CHICORY / ENDIVE

Treflan H.F.P. may be applied as a preplant incorporated treatment.

Broadcast Application Rates per Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints

26897

CHICORY

(*Cichorium intybus* or *Cichorium endiva*)

[Editor's note: This use revised as per previously accepted supplemental labeling (EPA-accepted 06/29/92)]

Treflan H.F.P. may be applied as a preplant incorporated treatment to chicory grown either as a root crop or leafy vegetable as indicated below:

Cichorium intybus, considered to be a root crop, may yield the following:

- Chicory - the dried and processed root used as a coffee substitute.
- Radicchio - green leaves harvested from field grown plantings.
- Belgian Endive - white leaves grown in the dark; growth from field grown rootstalks.

Cichorium endiva, considered to be a leafy vegetable, may yield the following:

- Escarole - curly green leaves from field grown plantings.
- Endive - very curly green leaves from field grown plantings.

Apply Treflan H.F.P. as a soil incorporated treatment in spring or early summer prior to planting.

Broadcast Application Rates per Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints

COLE CROPS - BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER

Direct Seeded Cole Crops

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.0
Fine	1.5

- Soils with 2-5% organic matter - 1.5 pints

Precaution: Direct seeded cole crops exhibit marginal tolerance to higher than recommended rates of Treflan H.F.P. Stunting or reduced stands may occur.

27897

Transplanted Cole Crops

Apply and incorporate Treflan H.F.P. prior to transplanting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

CORN - FIELD CORN ONLY**Postemergence Incorporated Treatment**

Apply Treflan H.F.P. as a postemergence treatment following cultivation and/or use of a preemergence herbicide. Treflan H.F.P. does not control established weeds. Apply when crop is well established (2 true leaf stage or taller). Apply as an over-the-top spray or as a directed spray using drop nozzles if foliage prevents uniform coverage of the soil surface.

Incorporation Directions

Applications of Treflan H.F.P. must be mechanically incorporated within 24 hours. Mechanical incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at a speed that will provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Water In Option for Coarse and Medium Textured Soils: On coarse and medium textured soils, Treflan H.F.P. may be incorporated by continuous rainfall or sprinkler irrigation amounting to at least one-half to one-inch of water. Best results are obtained if application is made immediately after a cultivation when the soil surface is open and porous. Rainfall or sprinkler irrigation prior to application will tend to consolidate and seal the soil surface and prevent the downward movement of Treflan H.F.P. that is expected under porous, open, recently tilled conditions. Supplemental irrigation can be applied through a center pivot, solid set or hand moved sprinkler system. Do not use furrow irrigation. Mechanically incorporate as described above if the required amount of rainfall or sprinkler irrigation does not occur within 24 hours after application.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	0.75-1.0†
Medium	1.25-1.5
Fine	1.5-2.0

† Apply 1.0 to 1.5 pt/acre on coarse soils in Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia to control fall panicum and Texas panicum.

- Apply lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Precautions:

- Do not apply to sweet corn, popcorn or corn grown for seed.
- Do not apply Treflan H.F.P. to corn as a preplant or preemergence treatment or crop injury may occur.
- Where corn is planted in a furrow, Treflan H.F.P. should be applied only after a cultivation to move soil into the row.

28 997

Restriction: Do not apply Treflan H.F.P. within 6 weeks prior to harvesting forage, fodder or silage or after corn is 30 inches tall.

Chemigation

Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in field corn. Refer to "Application by Chemigation" section in the "General Information" section of this label for chemigation use directions. Do not apply Treflan H.F.P. through any type of irrigation system unless these directions are carefully followed.

Application Timing

Apply Treflan H.F.P. in 0.5 to 1 acre inch of sprinkler irrigation when field corn is at the 2 true leaf stage of growth or taller. Apply Treflan H.F.P. prior to weed emergence or after existing weeds have been controlled with herbicides or cultivation. Treflan H.F.P. does not control established weeds.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5-2.0
Medium	1.5-2.0
Fine	Do not apply Treflan H.F.P. by chemigation to fine textured soils.

Precautions:

- Do not apply Treflan H.F.P. by chemigation to sweet corn, popcorn or corn grown for seed.
- Where corn is planted in a furrow, Treflan H.F.P. should be applied only after a cultivation to move soil into the row.
- Do not apply Treflan H.F.P. to corn as a preplant or preemergence treatment as crop injury may occur.

Restriction: Do not apply Treflan H.F.P. within 6 weeks prior to harvesting forage, fodder or silage or after corn is 30 inches tall.

~~Treflan H.F.P. plus Atrazine Tank Mix~~

~~Treflan H.F.P. may be applied in tank mix combination with atrazine plus an emulsifiable oil or oil concentrate when corn is at the 2-leaf stage of growth or taller and weeds are no more than 1 1/2 inches in height. A period of 24 to 48 hours is required to obtain atrazine postemergence activity after which the preemergence activity of the Treflan H.F.P. plus atrazine combination may be activated by 0.5 inch or more of rainfall or overhead sprinkler irrigation or mechanical incorporation. Use the application rates and incorporation methods for Treflan H.F.P. recommended under "Postemergence Incorporated Treatment" in the "Corn - Field Corn Only" section of this label.~~

Precautions:

- ~~• Where corn is planted in a furrow, Treflan H.F.P. should be applied only after a cultivation to move soil into the row.~~
- ~~• Refer to the product label for atrazine for application rates, additional use directions, precautions and limitations before use.~~

[Editor's note: Tank mix with atrazine removed from product label to attached supplemental labeling.]

29 3 97

COTTON**Treflan H.F.P. - Alone**

Apply Treflan H.F.P. to cotton as a soil incorporated treatment. Treflan H.F.P. may be applied before planting, immediately after planting, to the established crop up to layby (See supplemental labeling for postemergence applications), or in the fall in advance of spring planting. Refer to instructions for fall application under "Application Timing" in the "General Information" section of this label. Follow recommended soil preparation, application and incorporation procedures in the "General Information" section of this label. When incorporating Treflan H.F.P. after planting, but prior to crop emergence, set equipment so as to not disturb planted seed. Postemergence application of Treflan H.F.P. may be made from the 4 true leaf stage of growth up to layby, but not less than 90 days before harvest. Apply postemergence treatments as a directed spray beneath cotton plants to soil between the rows. Use the same application rates for preplant, preemergence and layby treatments.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.		
	Spring Application†	Fall Application	
		Eastern U.S.††	Western U.S.†††
	(pints)	(pints)	(pints)
Coarse	1.0	2.0	1.5
Medium	1.25-1.5	2.0	2.0
Fine	1.5-2.0	2.5	2.5

†Spring Application:

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 to 2.5 pints
- Use lower rate in rate range for areas receiving less than 20 inches of total annual rainfall and irrigation.

††Fall application rates for eastern cotton producing areas, including: Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri (Bootheel), North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas.

†††Fall application rates for western cotton producing areas, including: Arizona and California and Nevada. For cotton grown in states other than those listed above, fall apply at the highest broadcast rates for each soil texture under spring application, recommended for areas receiving greater than 20 inches average annual rainfall.

Precautions: Cotton should be planted after early season adverse weather conditions have passed, especially when using higher rate programs. Cool, wet weather early in the growth cycle causes additional stress to the cotton plant. This may result in reduced stand, delayed maturity and reduced yields.

Layby Treatment

Apply and incorporate Treflan H.F.P. in established cotton from the 4 true leaf stage of growth up to layby, but not less than 90 days before harvest. Apply uniformly to the soil surface, using drop nozzles if necessary. Use the application rates recommended above for preplant incorporated treatments. Set incorporate using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop.

Tank Mixes, Overlay and Postemergence Treatments

[Editor's note: This section moved to this location from end of "Special Use Programs section".]

30 of 97

Treflan H.F.P. in Tank Mix

~~Treflan H.F.P. may be tank mixed with Caparol, Cotoran, Zorial and other products registered for use on cotton as a preplant incorporated treatment to control additional weeds. Use the application rates for Treflan H.F.P. recommended for cotton "Treflan H.F.P. - Alone".~~ **Precaution:** ~~Refer to the tank mix product label for additional weeds controlled, application rates, additional use directions, precautions and limitations before use.~~

For broader spectrum weed control, other products registered for use in cotton may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

~~Treflan H.F.P. - Preplant Incorporated Followed by Overlay Treatments~~

~~Apply Treflan H.F.P. as a preplant incorporated treatment. Additional weeds tolerant to Treflan H.F.P. may be controlled using overlay preemergence applications of Cotoran, Karmex, Zorial or other products registered for use on cotton, unless use following Treflan H.F.P. is specifically prohibited by the manufacturer. Consult the manufacturer's labels for additional weeds controlled, use directions, precautions and limitations before use.~~

~~Treflan H.F.P. - Preplant Incorporated Followed by Postemergence Treatments~~

~~Apply Treflan H.F.P. as a preplant incorporated treatment. Additional weeds tolerant to Treflan H.F.P. may be controlled using postemergence treatments of products registered for use on cotton, unless use following Treflan H.F.P. is specifically prohibited by the manufacturer. Consult the manufacturer's labels for additional weeds controlled, use directions, precautions and limitations before use.~~

Special Use Programs

[Editor's note: Uses within this section have been numbered for sake of clarity.]

1. Cotton - Chemigation

[Editor's note: This use now included as part of Special Use Programs section.]

Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in cotton. Refer to "Application by Chemigation" in the "General Information" section of this label for use directions for chemigation. Do not apply Treflan H.F.P. through any type of irrigation system unless these directions are carefully followed.

Apply Treflan H.F.P. in overhead sprinkler irrigation equal to 1/2 to 1 inch of water. Planting and application should occur as soon as possible after the last tillage operation. Treflan H.F.P. must be applied within 2 days after planting prior to crop emergence. Treflan H.F.P. does not control established weeds. Soil incorporation is not required when Treflan H.F.P. is applied through chemigation systems.

[Editor's note: The chemigation instructions which follow have been revised to incorporate uses from previously accepted supplemental labeling entitled "Treflan, Applied Through Chemigation Systems for Weed Control in Conventional and Minimum Tilled Cotton" (EPA-accepted 07/30/90) and "Treflan for Weed Control in Conservation Tillage Cotton" (EPA-accepted 10/23/91).

Broadcast Application Rates/Acre for Chemigation Application Where Conventional Tillage Practices are Used: See rates for cotton "Treflan H.F.P. - Alone" above. Apply at the maximum recommended rate for spring application for each soil texture class to be treated.

Cultivation: Soil treated by chemigation with Treflan H.F.P. may be shallow cultivated without reducing weed control activity.

31 07 97

Broadcast Application Rates/Acre for Chemigation Application Where Minimum Tillage Practices are Used:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0 - 3.0
Medium	1.5 - 4.0
Fine	2.0 - 4.0

Use the lower rate in the rate range when additional sequential applications of Treflan are anticipated. Use the higher rate in the rate range when high crop residue levels are present, where dense weed populations are anticipated, or where no additional sequential applications of Treflan are to be made.

Rotational Crop Restrictions

- **Conventional Tillage:** Refer to the rotational crop restrictions in the General Use Precautions section of this label.
- **Minimum Tillage:** In addition to the rotational crop restrictions listed in the General Use Precautions section of this label, do not plant grain sorghum in the year following the application of Treflan.

2. Treflan for Cotton - Weed Control in Conservation Tillage in Cotton

[Editor's note: This section contains uses from previously accepted supplemental labeling entitled "Treflan for Weed Control in Conservation Tillage Cotton" (EPA-accepted 10/23/91). Proposed changes to previously accepted text are indicated by underline for added text and ~~strike-through~~ for deletions.]

This supplemental labeling section describes application methods and techniques for weed control with Treflan H.F.P. in conservation tillage cotton. Treflan H.F.P. may be applied and incorporated in the fall in advance of spring planting, in the spring before planting, after planting prior to crop emergence, or at layby. Single or multiple applications may be made so long as maximum application rates are not exceeded and rotational crop restrictions are followed.

Broadcast Application Rates for Conservation Tillage:

Soil Texture	Treflan H.F.P. (pints/A)
Coarse	1.0 - 2.0
Medium	1.5 - 2.0
Fine	2.0 - 4.0

Strip Planting into Small Grain Cover Crops

Fall planted cover crops may be utilized to control wind erosion and protect developing crop seedlings from wind damage. Prior to planting cotton, the cover crop may be treated with a contact herbicide to prevent continued growth and development and prevent competition with crop seedlings for water and soil nutrients. The standing cover crop (now dead) continues to control wind erosion and provide protection to the developing crop until it is well established.

Strip Planting: In strip planting, cotton is seeded into competition-free bands established in the cover crop. Competition-free bands may be established by leaving unseeded drill rows when seeding the cover crop, by tillage, or by use of a contact herbicide to prepare competition-free bands prior to planting.

Fall Application Before Establishing a Cover Crop

Small grain cover crops (wheat, barley or rye) may be established following a preplant incorporated application of Treflan H.F.P. Apply Treflan H.F.P. to flat ground at a broadcast rate of 2 to 3 pints per acre and incorporate once within 24 hours using incorporation implements that can be set to cut no more than 2 to 3 inches deep, such as a springtooth harrow. Do not incorporate with a tandem disc. Form beds with disc bedders or other bedding implements that will mix and move most of the treated soil (top) the furrows to the beds. Phosphate and other fertilizer may be applied as appropriate during incorporation

32 of 97

operations. Plant 2 to 4 rows of the small grain cover crop 2 inches deep in the furrows between the beds. To avoid injury to small grain seedlings, place seed below the treated layer of soil. Barley is usually less susceptible to injury than wheat or rye. Soil moisture must be adequate to establish and maintain the cover crop. In late winter (February), apply 2,4-D if necessary for broadleaf weed control.

Spring Application Before or After Planting (Within Competition-Free Bands)

Apply Treflan H.F.P. using low pressure ground equipment as a band (within the weed free zone) or as a broadcast treatment. Application and incorporation may occur before planting or after planting prior to crop emergence. If applied after planting, set incorporation equipment so as to not disturb the planted seed (see incorporation instructions).

Incorporation: Equipment should be adapted to the width of the competition-free band. Use equipment that will uniformly mix Treflan H.F.P. into the weed germination zone. Weed control resulting from single pass incorporation or with incorporation equipment that does not result in thorough mixing of Treflan H.F.P. treated soil treated with Treflan H.F.P. may be reduced compared to conventional double pass incorporation. Implements used to incorporate Treflan H.F.P. after planting should be operated so that they do not disturb the planted seed or growing crop.

Use the lower rate in the rate range when additional sequential applications of Treflan are anticipated. Use the higher rate in the rate range where high crop residues are present, and where dense weed populations are anticipated.

For band treatments, reduce the application rate in proportion to the row spacing and band width treated. For example, treating a 12-inch band where the row spacing is 36 inches would require 1/3 of the recommended broadcast rate per acre (12 inches divided by 36 inches = 1/3).

Layby Applications

Layby applications may be made in established cotton from the 4 true leaf stage of growth up to layby, but not less than 90 days before harvest. Apply Treflan H.F.P. uniformly to the soil surface using drop nozzles if necessary. Soil incorporate using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop. Cumulative layby application rate may not exceed the layby application rate shown for each soil texture.

Broadcast Application Rates:

Soil Texture	Treflan H.F.P. (pints/A)
Coarse	1.0
Medium	1.5
Fine	2.0

Chemigation

[Editor's note: Instructions from this section incorporated into the Cotton Chemigation section above.]

~~Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in conservation tillage cotton. Apply the recommended rate of Treflan H.F.P. in 0.5 to 1.0 acre inch of sprinkler irrigation. Apply Treflan before planting or after planting, but prior to weed germination. Treflan does not control established weeds. Soil incorporation is not required when Treflan is applied by chemigation. Soil treated with Treflan may be shallow cultivated without reducing weed control activity. Refer to supplemental labeling entitled "Treflan H.F.P. Chemigation Instructions" for detailed use directions for chemigation. Do not apply Treflan H.F.P. through any chemigation system unless supplemental labeling on chemigation is followed.~~

~~Do not apply Treflan 5 or Treflan M.T.F. through any type of irrigation system.~~

33 of 97

Broadcast Application Rates:

Soil Texture	Treflan H.F.P. (pints/A)
Coarse	1.0 - 3.0
Medium	1.5 - 4.0
Fine	2.0 - 4.0

Use the lower rate in the rate range when additional sequential applications of Treflan are anticipated. Use the higher rate in the rate range when high crop residue levels are present, where dense weed populations are anticipated, or where no additional sequential applications of Treflan are to be made.

Repeat, Sequential Applications

Treflan may be applied one or more times sequentially during the growing season using the rates and methods of application described above for full season weed control. The maximum dosage that can be used for a single application cannot exceed the rates shown for each application method. The maximum cumulative application rate that may be applied within the same growing season (including fall applications) cannot exceed 4 pints per acre for Treflan H.F.P. (2 pounds active ingredient per acre).

Contact, overlay or Postemergence Herbicides

Contact herbicides approved for use in cotton may be used to control existing weeds prior to planting cotton. To control additional weeds, tolerant to Treflan may be controlled using overlay, preemergence or postemergence applications of other products registered for use on cotton may be applied. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

Rotation Crop Restrictions

Refer to the General Information section of this label for specific rotational crop restrictions. When the cumulative application rate exceeds the application rates in the table below, plant only those crops for which Treflan H.F.P. can be applied as a preplant incorporated treatment in the season following the application of Treflan H.F.P. or crop injury may result.

Soil Texture	Cumulative Application Rate
	Treflan H.F.P. (pints/acre)
Coarse	1.5
Medium	1.5
Fine	2.0

Small grain cover crops that will not be grazed or harvested and are intended for prevention of wind erosion in conservation tillage cotton may be planted in the fall following spring applications of up to 4 pints per acre of Treflan H.F.P. Injury in the form of reduced stands or delayed emergence and development may result when small grains are planted under these conditions.

3. Cotton - Fall Panicum Control

Apply and incorporate a broadcast rate of 2.0 p/acre on both coarse and medium soils.

34 of 97

4. Cotton - Pigweed and Seedling Johnsongrass Control

Apply Treflan H.F.P. as a preplant incorporated treatment.

Broadcast Application Rates/Acre: In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, southeastern Missouri (Bootheel), North Carolina, South Carolina, Tennessee and southern Virginia, apply Treflan H.F.P. at the following broadcast rates:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0-1.5
Medium	1.5-2.0
Fine	2.0

(Exception: Louisiana, where 3.0 pt/acre can be applied to fine soils).

- Use higher rates in the rate range where high weed populations are anticipated.

5. Cotton - Additional Weed and Grass Control in (Gulf Coast Counties of Texas)

Apply Treflan H.F.P. as a preplant incorporated treatment up to 2 weeks before planting.

Broadcast Application Rates/Acre: For cotton grown in Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton counties of the Texas Gulf Coast, apply Treflan H.F.P. at the following broadcast rates:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5
Medium	2.0
Fine	3.0

6. Cotton - Rhizome Johnsongrass Control

(For use in all cotton producing states except Arizona and California.)

Rhizome johnsongrass control with Treflan H.F.P. requires double application rates for 2 consecutive years. Commercially acceptable control cannot be obtained with only one year of double rate use of Treflan H.F.P. Carefully follow all special use directions.

Soil Preparation: Satisfactory results are dependent upon proper preparation of soil prior to application. Chisel plow to bring rhizomes to the soil surface. Disc twice before application to chop rhizomes into small (2 to 3 inch) pieces and destroy any recently emerged johnsongrass plants.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	2.0
Medium	3.0
Fine	4.0

Spring Application: Apply Treflan H.F.P. any time before planting in the spring for 2 years in succession.

Fall Application: Apply Treflan H.F.P. between October 15 and December 31 for 2 years in succession.

Incorporation: Deep incorporation with a tandem disc is essential for good results. Set disc to operate 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary and the second should be in a different direction than the first.

Cultivation: Some johnsongrass plants will not be controlled. Timely cultivation during the crop season is necessary to remove escaped plants and maintain commercially acceptable control.

Precautions:

In the season following a double rate treatment, plant only rice or those crops for which Treflan H.F.P. can be applied as a preplant treatment or crop injury may occur.

COTTONWOOD TREES GROWN FOR PULP

[Editor's note: This use added from previously accepted supplemental labeling (EPA-accepted 10/23/91).]

Apply as a soil incorporated treatment to control weeds susceptible to Treflan H.F.P. in new and established plantings of cottonwood trees grown for pulp.

Application Before Planting

Apply and incorporate Treflan H.F.P. before planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25 - 1.5
Fine	1.5 - 2.0

- All soils with 2-5% organic matter - 1.5-2.0 pints
- All soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total rainfall and irrigation.

Application to Established Plantings

In established plantings, apply Treflan H.F.P. as a directed spray to the soil and use incorporation methods not injurious to the crop.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
All Soil Textures	2.0-4.0

Application rate within the rate range may be adjusted according to weed pressure.

Johnsongrass Suppression in Established Plantings

Proper soil preparation before application is necessary for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the soil surface. Then work the soil twice using a tandem disc to cut rhizomes into small (2 to 3 inch) pieces and to destroy emerged johnsongrass.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
All Soil Textures	4.0

Incorporation: Incorporate twice with tandem disc set to cut 4 to 6 inches deep and operated at 4 to 6 mph.

Cultivation: Some johnsongrass plants will escape. Timely cultivation with tillage implements or spot spraying with effective postemergence herbicides will improve the level of johnsongrass control.

36 of 97

CUCURBITS — CANTALOUPE, CUCUMBER AND WATERMELON

Apply Treflan H.F.P. after emergence when plants have reached the 3 to 4 true leaf stage of growth. Apply as a directed spray to soil between the rows. Avoid foliage contact as slight crop injury may occur. Set incorporation equipment to move treated soil around the base of plants.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Restriction: Do not apply within 30 days of harvest, except for Watermelon which has a 60 day pre-harvest interval.

FLAX (Fall Application Only)

Apply and incorporate Treflan H.F.P. in the fall for weed control in spring seeded flax. Incorporate once within 24 hours after application. The second incorporation may be performed in the spring prior to planting.

Special Instructions for Flax

1. Incorporation operations or other tillage practices performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed, and the seedbed should be packed prior to seeding.
2. Seeding should be done with a press drill or hoe drill. Seed into moist seedbed and plant no more than 1.5 inches deep.
3. Delay seeding until soil has warmed sufficiently to allow rapid germination and establishment.
4. Refer to "General Use Precautions" in the "General Information" section of this label for information on growing conditions that can lead to crop injury or yield reduction.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	2.0

37 of 97

FORAGE LEGUMES

Forage Legumes Used as Cover Crops or in the Acreage Conservation Reserve Program

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.0-1.5
Fine	1.5

- Use the lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Precautions:

- If used under the Acreage Conservation Reserve Program, follow the most severe grazing restrictions imposed by the USDA Conservation Use Program. Consult the local ASCS committee or other state agency to determine the period of the USDA grazing restriction.
- Some crop stand reduction may occur with this use; however, reduced weed competition will allow establishment of a quality stand.

GRAIN SORGHUM (MILO)

Postemergence Incorporated Treatment

Apply Treflan H.F.P. as a directed or over-the-top spray when grain sorghum is 8 to 24 inches tall or taller. Drop nozzles should be used if foliage prevents uniform soil coverage.

Soil Preparation: Cultivate before application of Treflan H.F.P. to remove established weeds and to cover the base of grain sorghum plants with soil. Cultivation equipment should be set to add approximately 1 inch of soil to the base of sorghum plants.

Incorporation Directions: Applications of Treflan H.F.P. must be mechanically incorporated within 24 hours after application. Mechanical incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Sweep-type cultivators should have 3 to 5 sweeps per row middle and be operated at a speed that will provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	0.75-1.0
Medium	1.0-1.5
Fine	1.5-2.0

- Apply Treflan H.F.P. at lower rate in rate range in areas receiving less than 20 inches total rainfall and irrigation.

Precautions:

- Do not apply Treflan H.F.P. to grain sorghum as a preplant or preemergence treatment or crop injury will occur.
- Over-application may result in injury to grain sorghum.

Restriction: Do not apply after grain sorghum is 24 inches tall.

Chemigation

Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in grain sorghum 8 to 24 inches tall or taller. Refer to "Application by Chemigation" section in the "General Information" section of this label for chemigation use directions. Do not apply Treflan H.F.P. through any irrigation system unless these directions are carefully followed.

Soil Preparation: Cultivate before application of Treflan H.F.P. to destroy existing weeds and cover the base of the grain sorghum plants with soil. Cultivation equipment should be set to add approximately 1 inch of soil to the base of sorghum plants.

Application Timing: Apply Treflan H.F.P. to grain sorghum in 0.5 to 1 acre inch of overhead sprinkler irrigation as soon as possible after a cultivation when grain sorghum is at least 8 to 24 inches tall. Treflan H.F.P. must be applied prior to weed emergence or after existing weeds are controlled. Treflan H.F.P. does not control established weeds.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	0.75-1.0
Medium	1.0-1.5
Fine	Do not apply Treflan H.F.P. by chemigation to fine textured soils.

Restriction: Do not apply after grain sorghum is 24 inches tall.

~~Treflan H.F.P. plus Atrazine Tank Mix~~

~~Treflan H.F.P. may be applied in tank mix combination with atrazine plus an emulsifiable oil or oil concentrate when grain sorghum is 8 inches tall or taller and weeds are no more than 1 1/2 inches in height. A period of 24 to 48 hours is required to obtain postemergence activity of atrazine after which the preemergence activity of the Treflan H.F.P. plus atrazine combination may be activated by 0.5 inch or more of sprinkler irrigation or mechanical incorporation. Use application rates and incorporation methods for Treflan H.F.P. recommended under "Postemergence Incorporated Treatment" in the "Grain Sorghum (Milo)" section of this label.~~

Precautions:

- Where grain sorghum is planted in a furrow, Treflan H.F.P. should be applied only after a cultivation to move soil into the row.
- Refer to the product label for atrazine for application rates, additional use directions, precautions and limitations before use.

GREENS - TURNIP GREENS GROWN FOR PROCESSING: COLLARD, KALE AND MUSTARD GREENS

Apply Treflan H.F.P. to greens as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	1.5

- Soils with 2-10% organic matter - 1.5 pints

39 of 97

HOPS

Apply and incorporate Treflan H.F.P. to established crop during dormancy. Use incorporation equipment that will insure thorough soil mixing with minimal damage to crop stand.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25 - 1.5
Fine	1.5

- Soils with 2-10% organic matter - 1.5 pints

KENAF

[Editor's note: This use added from previously accepted supplemental labeling (EPA-accepted 02/04/92).]

Apply as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse†	1
Medium	1 - 1.5
Fine	1.5

†Coarse soils with 2-5% organic matter - 1.5 pints

- Use higher rate in rate range where high weed populations are anticipated.

Precaution: Do not graze or harvest treated crop for livestock forage.

MUSTARD - GROWN FOR SEED OR PROCESSED FOR FOOD

Apply Treflan H.F.P. to mustard as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	1.5

- Soils with 2-10% organic matter - 1.5 pints

40097

OKRA

Apply Treflan H.F.P. as a soil incorporated treatment, before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

~~ONIONS GROWN FOR DRY BULBS ONLY~~

~~Apply Treflan H.F.P. to established onions as a soil incorporated treatment. Apply as a directed spray to soil between onion rows. Spray shields should be used to avoid injury to foliage or exposed bulbs. Do not apply within 60 days of harvest.~~

~~Broadcast Application Rates/Acre:~~

Soil Texture	Treflan H.F.P. (pints)
Coarse	0.75-1.0
Medium	1.0-1.25

- ~~• Use lower rate in rate range in areas receiving less than 20 inches total rainfall and irrigation or where light weed pressure is anticipated~~

Incorporation

Incorporate with 1 pass of a sweep-type or rolling cultivator. Set equipment to cut 2 to 4 inches deep and operate at 6 to 8 mph. Avoid covering exposed onion bulbs with treated soil during incorporation as crop injury may occur. Avoid injury to crop roots during incorporation.

Precautions: When applied according to directions under normal growing conditions, Treflan H.F.P. will not adversely affect onions. Diseases, improper incorporation depth, excessive moisture, high salt concentration or drought may weaken the crop and increase the possibility of damage from Treflan H.F.P. Under these conditions, delayed crop development or reduced yields may result.

41 of 97

ONIONS (DRY BULBS ONLY)

Postemergence Layby Application: Apply at layby to the soil between onion rows. Avoid applying directly to the tops or exposed bulbs of onion plants. Emerged weeds should be removed prior to application of Treflan H.F.P. Treflan H.F.P. will not control established weeds.

Broadcast rates per acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	0.75 - 1.0
Medium	1.0 - 1.25

- Apply only to soils containing 3.5% or less organic matter
- Note: Use the lower rate in rate range where light weed pressure is anticipated.

Incorporation: Treflan H.F.P. should be uniformly incorporated into the soil between the onion rows. Incorporation may be accomplished by operating a sweep-type or rolling cultivator 2 to 4 inches deep at 6 to 8 mph. Two incorporation passes are required with the first occurring within 24 hours after application or erratic weed control may result. Avoid covering onions with treated soil during incorporation as injury to the crop may occur. Care should be taken to avoid mechanical injury to onion roots during incorporation.

Precautions

- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Do not apply as a preplant or preemergence treatment.
- Do not apply to muck soils.
- Note: Reduced yields may result from use of Treflan H.F.P. on onion crops weakened by diseases. Improper incorporation depth, excessive moisture, high salt concentration or drought may weaken the crop and increase the possibility of damage from Treflan H.F.P. Under these conditions reduced yields may result.

PEAS - DRY PEAS AND ENGLISH PEAS

Treflan H.F.P. - Alone

Apply and incorporate Treflan H.F.P. in the spring before planting or in the fall in advance of spring planting. Refer to instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	
	Spring Application (pints)	Fall Application† (pints)
Coarse	1.0	1.0
Medium	1.0 - 1.5††	1.25-1.5
Fine	1.5	1.5

- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.
- †Treflan H.F.P. may be fall applied to dry and English peas in the states of Idaho, Oregon and Washington.
- ††Medium soils with 3% or greater organic matter - 1.5 pints

42 of 97

Treflan H.F.P. plus Far-Go Tank Mix (For Use in Idaho, Oregon and Washington)

Treflan H.F.P. may be tank mixed with Far-Go and applied as a preplant soil incorporated treatment to control wild oats in dry and English peas. Use application rates recommended for dry and English peas "Treflan H.F.P. - Alone", above. Refer to the label for Far-Go for application rates, additional use directions, precautions and limitations before use.

Tank Mixing

For broader spectrum weed control, other products registered for use in dry and English peas may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

PEAS - SOUTHERN PEAS

Apply as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- All soils with 5-10% organic matter - 2.0 pints
- Use the lower rate in the rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

PEANUTS**Treflan H.F.P. - Alone**

Spanish Peanuts, Florunner and Florigiant Varieties (For Use in Texas, Oklahoma and New Mexico Only)

Apply and incorporate Treflan H.F.P. before planting, at planting or immediately after planting. When incorporating after planting, adjust equipment so as to not disturb planted seed.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5

Treflan H.F.P. plus Vernam Tank Mix

Spanish Peanuts, Florunner and Florigiant Varieties (For Use in Texas, Oklahoma and New Mexico)

Treflan H.F.P. may be tank mixed with Vernam and applied as a preplant incorporated treatment to control additional weeds. Use application rates recommended for peanuts "Treflan H.F.P. - Alone" above. Refer to the labels for Vernam for application rates, additional use directions, cautions and limitations before use.

430/97

Tank Mixing

For broader spectrum weed control, other products registered for use in peanuts may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

PEPPER (Transplant Only)

Apply and incorporate Treflan H.F.P. prior to transplanting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

~~POTATOES (Not for Use in the State of Maine)~~

[Editor's note: Use on potatoes removed from this main label and placed on attached supplemental labeling.]

~~Treflan H.F.P. - Alone~~

~~Apply and incorporate Treflan H.F.P. after planting prior to crop emergence, immediately following dragoff, or after potato plants have fully emerged.~~

~~**Incorporation:** Set incorporation equipment so that the bed and furrow are uniformly covered with a layer of treated soil. If the layer of treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. When applying and incorporating Treflan H.F.P. after potato plants have fully emerged, do not completely cover the plants with treated soil. Likewise, do not completely cover plants during subsequent cultivations. Be careful that incorporation equipment does not damage potato seed pieces or elongating sprouts.~~

~~Broadcast Application Rates/Acre:~~

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- ~~• Coarse and medium soils with 2-5% organic matter - 1.5 pints~~
- ~~• Fine soils with 2-5% organic matter - 2.0 pints~~
- ~~• Soils with 5-10% organic matter - 2.0 pints~~
- ~~• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.~~

~~Chemigation~~

~~Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in potatoes. Refer to "Application by Chemigation" section in the "General Information" section of this label.~~

44 of 99

Do not apply Treflan H.F.P. through any type of irrigation system unless these directions are carefully followed.

Apply Treflan H.F.P. to potatoes in 0.5 to 1 acre inch of overhead sprinkler irrigation after planting, before emergence, or immediately following dragoff or after the potato plants have fully emerged. Existing weeds must be destroyed by tillage or cultivation prior to application of Treflan H.F.P. Treflan H.F.P. does not control established weeds. Incorporation is not necessary when Treflan H.F.P. is applied by chemigation.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.5
Fine	Do not apply Treflan H.F.P. by chemigation to fine textured soils.

Precautions:

- If cultivation is required after treatment with Treflan H.F.P., avoid completely covering potato plants with treated soil.
- Erratic weed control may result if cultivation exposes untreated soil between rows.

Treflan H.F.P. in Tank Mix

Treflan H.F.P. plus Eptam Tank Mix - Post Plant Preemergence Treatment
Treflan H.F.P. may be tank mixed with Eptam and applied as a soil incorporated treatment to control additional weeds. Apply after planting, but before crop emergence. In areas where potatoes are normally dragged off, apply and incorporate up to or immediately following drag off. Use application rate recommended for potatoes "Treflan H.F.P. Alone". Refer to the label for Eptam for application rates, additional use directions, precautions and limitations before use.

RADISH

Apply Treflan H.F.P. as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.5
Fine	1.5

RAPSEED (CANOLA)

Apply and incorporate Treflan H.F.P. in the spring before planting or in fall. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.5
Fine	2.0

Precaution: Do not apply Treflan H.F.P. to rapeseed (canola) grown in the state of Alaska.

RAPESEED (CANOLA) AND CRAMBE

[Editor's note: This use revised/added (crambe) as per previously accepted supplemental labeling, EPA-accepted 03/30/93.]

Apply as a soil incorporated treatment in the spring before planting or in late summer or early fall before a fall planting. Follow soil preparation, application and incorporation directions for Treflan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	2.0

Precautions

- Do not apply to rapeseed (canola) grown in the state of Alaska.
- Where applications are made in late summer or fall, plant as rotation crops in the season following application only those crops to which Treflan H.F.P. may be applied as a preplant incorporated treatment or crop injury may occur.
- Do not graze or harvest crambe for livestock forage.

SAFFLOWER

Apply and incorporate Treflan H.F.P. in the spring before planting or in fall in advance of spring planting. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	
	Spring Application (pints)	Fall Application† (pints)
Coarse	1.0	1.5
Medium	1.25-1.5	2.0
Fine	1.5-2.0	2.5

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.5 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.
- † Treflan H.F.P. may be fall applied to safflower in the states of Arizona, California, Idaho, Nevada, Oregon, Utah, Washington and Wyoming.

4/6 of 97

SMALL GRAINS - BARLEY, DURUM AND WHEAT

Special Precautions for Use of Treflan H.F.P. on Small Grains

Carefully follow directions for use of Treflan H.F.P. on small grains to minimize potential crop stress. Under certain conditions, delayed crop emergence and or stand reduction may occur when Treflan H.F.P. is applied to barley, durum or wheat. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield. For best results, observe the following cultural practices or precautions:

Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.

Irrigate prior to planting or after germination and emergence. Moisture received between planting and emergence may cause crusting, especially on loose, friable seedbeds.

Do not exceed recommended application rates for Treflan H.F.P. This is particularly important on coarse textured or low organic matter soils.

Carefully follow incorporation directions. When applying preplant incorporated treatments, operate equipment at recommended depth and speed to place Treflan H.F.P. into the upper 1 to 1.5 inches of soil. If applied after planting, set equipment so as to not disturb planted seed.

Set drills to place seed at a the depth specified in use directions. A planting depth greater than 2.5 inches for spring wheat or durum will result in increased seedling stress and decreased emergence.

Use only high quality seed where Treflan H.F.P. is to be applied (avoid use of small seed with low starch reserves).

If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.

Avoid use of seed varieties known to have poor seedling (emergence) vigor.

~~Do not fall apply Treflan H.F.P. in combination with any other preplant incorporated herbicide.~~

Soil characteristics and environmental conditions which may contribute to crop seedling stress that may be accentuated by use of Treflan H.F.P. include:

Soil related: High salinity, eroded knolls/hilltops, loose dry soils and compaction.

Weather related: Cold and/or wet soils, excessively hot soils, excessive moisture, drought, and soil crusting from heavy rainfall.

Note: Do not apply Treflan H.F.P. on small grains where a dinitroaniline herbicide such as Treflan-H.F.P. or Sonalan® herbicide was applied at a rate recommended for row, crops (oil seeds) during the previous growing season greater than 0.5 lb ai per acre the previous growing season.

470297

Application Directions for Small Grains

Barley, Spring Seeded -- Spring Application Preplant Incorporated for Foxtail (Pigeongrass) Control (For Use in Minnesota, North Dakota and South Dakota)
Apply Treflan H.F.P. as a preplant incorporated treatment prior to planting spring seeded barley. Treflan H.F.P. may be applied to ground that has a manageable trash level of crop residue or has been fallowed or pre-tilled. The first incorporation is required within 24 hours after application. The second incorporation is required prior to planting to destroy emerged weeds and to insure even distribution of Treflan H.F.P. in the soil surface.

Broadcast Application Rates/Acre: Apply at a rate of 1.0 pint per acre for all soil textures regardless of organic matter content.

Incorporation: Recommended incorporation tools include the chisel plow (first incorporation pass only), tandem disc and field cultivator. Refer to "Incorporation Equipment" in "General Information" section of this label for details on operation of incorporation equipment.

Planting Directions: Barley should be seeded approximately 2 1.5 inches deep.

Precautions:

- Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.
- While use of this weed control practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Barley, Spring Seeded -- Spring Application Preplant Incorporated for Foxtail (Pigeongrass) Control in Barley Used as a Cover Crop or in the Conservation Reserve Program

Apply Treflan H.F.P. as a preplant incorporated treatment prior to planting spring seeded barley on land enrolled in acreage conservation reserve programs. Follow recommended soil preparation, application and incorporation procedures for Treflan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.5
Fine	1.5

Planting Directions: Barley should be seeded approximately 2 1.5 inches deep.

Precautions: Use of this weed control practice may result in slight stand reduction. Follow the most severe grazing restrictions imposed either by the label for Treflan H.F.P. or by the USDA Acreage Conservation Reserve Program, whichever is longest. Consult the local ASCS office or other state agency to determine the period of USDA grazing restriction.

Winter Wheat -- Preplant Incorporated for Control of Cheatgrass and Other Annual Grasses and Broadleaves (For Use in Idaho, Oregon and Washington)

Apply Treflan H.F.P. as a preplant incorporated treatment for control of downy brome (cheatgrass), annual ryegrass, annual bluegrass, pacific meadow foxtail (blackgrass), henbit and fiddleneck (tarweed). The growth, development and yield of winter wheat will not be adversely affected, provided the seed is placed below the zone of soil treated with Treflan H.F.P. Treflan H.F.P. may be applied for up to 3 weeks before planting.

48 of 97

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5
Medium	1.5
Fine	2.0

Incorporation Directions: Incorporate Treflan H.F.P. with a flexible tine-tooth harrow (Flextine or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Incorporate once within 24 hours after application and a second time in a different direction from the first prior to planting. Do not till the soil with a disc after Treflan H.F.P. has been incorporated with a flexible tine harrow.

Planting Directions: Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil treated with Treflan H.F.P.

Precautions:

- Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.
- Wheat planted in direct contact with treated soil may suffer crop injury in the form of delayed emergence and development.

Winter Wheat -- Post Plant Incorporated Treatment

(For Use in Idaho, Oregon and Washington)

Apply and incorporate Treflan H.F.P. after planting, but before emergence, to control the following weeds susceptible to Treflan H.F.P. in winter wheat: annual ryegrass, annual bluegrass, downy brome (cheatgrass), pacific meadow foxtail (blackgrass), fiddleneck (tarweed) and henbit.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0-1.5
Medium	1.5

Planting Directions: Plant wheat 2 to 3 inches deep in a well-tilled seedbed. Do not use a deep or semi-deep furrow drill.

Incorporation Directions: Incorporate Treflan H.F.P. using 2 passes with a flex-tine or spike-tooth harrow operated at least 5 mph. The second incorporation pass should be in a different direction than the first. Set equipment to cut 1 to 1 1/2 inches deep and avoid disturbing seed. Application and first incorporation should be done in the same operation if possible. Both incorporations must be done within 24 hours.

Precautions:

- Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.
- Wheat seed in direct contact with treated soil may suffer crop injury in the form of delayed emergence and development.
- If less than 20 inches of rainfall plus irrigation was received between planting and harvest, refer to rotation crop restrictions before planting sorghum or oats.

Winter Wheat -- Fallow Soil Application Prior to Planting (For Use in Idaho, Oregon and Washington)

Treflan H.F.P. may be applied and shallowly incorporated into fallow soil up to 4 months before planting wheat to control cheatgrass and certain annual grasses and broadleaf weeds. Apply Treflan H.F.P. any time from May to September prior to fall planting of winter wheat. Wheat growth, development and yield will not be adversely affected so long as the seed is placed below the zone of soil treated with Treflan H.F.P.

49 of 97

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5
Medium	1.5
Fine	2.0

Incorporation Directions: Incorporate Treflan H.F.P. with a flexible tine-tooth harrow (Flextine or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Incorporate once within 24 hours after application and a second time in a different direction from the first prior to planting. Do not till the soil with a disc after Treflan H.F.P. has been incorporated with a flexible tine harrow.

Planting Directions: Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil treated with Treflan H.F.P.

Precautions:

- Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.
- Wheat planted in direct contact with treated soil may suffer crop injury in the form of delayed emergence and development.

Wheat, Durum and Barley, Spring Seeded - Fall Applied
Preplant Soil Incorporated for Foxtail (Pigeongrass) Control
 (For Use in Minnesota, North Dakota and South Dakota)

[Editor's note: This use added from previously accepted supplemental labeling (EPA-Accepted 02/24/92).]

Apply Treflan H.F.P. herbicide in the fall for foxtail (pigeongrass) control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure a uniform distribution of Treflan H.F.P. in treated soil. Treflan H.F.P. may be applied to ground that has a manageable trash level of crop residue, or has been fallowed or pre-tilled.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse and Medium	1
Fine	1.5

Incorporation: Recommended incorporation tools include the chisel plow (first incorporation pass only), tandem disc and field cultivator. Refer to "Incorporation Equipment" in "General Information" section of this label for details on operation of incorporation equipment.

Planting Directions

Set equipment to place seed approximately 1.5 inches deep.

Precautions

Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.

While use of this control practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Spring Wheat, Durum and Barley - Postplant Incorporated for Foxtail (Pigeongrass) Control

Apply and incorporate Treflan H.F.P. after planting, but before emergence, to control foxtail (pigeongrass) in spring wheat, durum and barley. Treflan H.F.P. may be tank mixed with Far-Go to control wild oats and other weeds. Refer to the label for Far-Go for application rates, additional use directions, precautions and limitations before use.

50 of 97

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.0
Fine	1.5

Planting Directions: Plant wheat 2 to 3 inches deep in a well-tilled seedbed.

Incorporation Directions: Incorporate Treflan H.F.P. using 2 passes with a flex-tine or diamond harrow operated at least 5 mph. The second incorporation pass should be in a different direction than the first. Set equipment to cut 1 to 1.5 inches deep and avoid disturbing seed. Application and first incorporation should be done in the same operation if possible. Both incorporations must be done within 24 hours.

Precautions:

- Carefully read and follow "Special Precautions for Use of Treflan H.F.P. in Small Grains" before application of Treflan H.F.P.
- Wheat seed in direct contact with treated soil may suffer crop injury in the form of delayed emergence and development.

SOYBEANS

Treflan H.F.P. - Alone

Apply and incorporate Treflan H.F.P. in the spring before planting or in the fall in advance of spring planting, as a preplant soil incorporated treatment. Treflan H.F.P. may also be applied in the fall. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	
	Spring Application (pints)	Fall Application† (pints)
Coarse	1.0	2.0
Medium	1.5	2.0
Fine	2.0	2.5

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 to 2.5 pints

†Fall Application Rates for States including: Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri (Bootheel), North Carolina, Oklahoma, South Carolina, Tennessee and Texas.

For soybeans grown in states other than those listed above, fall apply Treflan H.F.P. at broadcast rates recommended for normal spring preplant incorporated treatment.

Precautions: Soybeans should be planted after early season adverse weather conditions have passed, especially when using higher rate programs. Cool, wet weather early in the growth cycle causes additional stress on soybean plants which may result in reduced stand, delayed maturity and reduced yield.

Tank Mix, Overlay and Postemergence Recommendations

Treflan H.F.P. In Tank Mix

Treflan H.F.P. may be tank mixed with Sencor, Lexone, Canopy, Lasso, Dual, Preview or Vamam and applied as a preplant soil incorporated treatment to control additional weeds in soybeans. Refer to the tank mix product label for weeds controlled, application rates, additional use directions, precautions and limitations before use.

~~Treflan H.F.P. plus Command (Reduced Rate) and Treflan H.F.P. plus Command and Lexone or Treflan H.F.P. plus Command and Sencor Tank Mixes (Not For Use in California):~~ Treflan H.F.P. may be tank mixed with Command, Command plus Lexone or Command plus Sencor. Apply the tank mix as a preplant incorporated treatment up to 3 weeks before planting.

Note: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10-15 gal/acre. Use nozzles suitable for broadcast boom application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays. Application to overly moist or wet soils will increase the potential for off-site movement of Command vapors and may result in poor soil incorporation and unsatisfactory weed control. These directions must be followed to reduce the potential for off-site movement of Command vapors and potential injury to desirable vegetation including adjacent crops, trees and ornamentals.

Incorporation: Tank mixes containing Command must be incorporated immediately after application. Follow other soil preparation, application and incorporation procedures for Treflan H.F.P.

Treflan H.F.P. plus Command: Use the Treflan H.F.P. plus Command tank mix to control velvetleaf and weeds susceptible to Treflan H.F.P.

Control of Jimsonweed, annual morningglory, prickly sida, common ragweed, smartweed and Venice mallow may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	Command 4 H.F.P.
	(pints)	(pints)
Coarse	1.0	0.75
Medium	1.5	1.12
Fine	2.0	1.5

~~Treflan H.F.P. plus Command and Lexone or Treflan H.F.P. plus Command and Sencor:~~ Use the Treflan H.F.P. plus Command and Lexone or Sencor tank mix to control weeds susceptible to Treflan H.F.P. plus additional weeds listed on the labels for Command and Lexone or Sencor.

Treflan H.F.P. plus Command and Lexone or Sencor also provides partial control or suppression of cocklebur, annual morningglory and giant ragweed. Control of these weeds may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.	Command 4 H.F.P.	Lexone 4L or Sencor 4	Lexone or Sencor DF
	(pints)	(pints)	(pints)	(pounds)
Coarse	1.0	0.5	0.33-0.5†	0.25-0.33†
Medium	1.5	0.75	0.5-0.75	0.33-0.5
Fine	2.0	1.12	0.75-1.0	0.5-0.67

52 of 97

† Use the higher rate range in areas where weed populations are dense, for control of Venice mallow and wild mustard, and for best control of common cocklebur, annual morningglory and giant ragweed.

Precautions:

- Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of adjacent crops, trees and ornamental plants which is usually temporary in nature but can result in permanent injury or death of the plants if the exposure is excessive. Prior to making application of this product, read and strictly follow all precautions, rotational crop guidelines and application instructions on the label for Command.
- Refer to the labels for Lexone and Sencor for additional use directions, precautions and limitations before applying Trellan H.F.P. plus Lexone or Trellan H.F.P. plus Sencor tank mix.

~~Preplant Incorporated Followed by Overlay Treatments (Not For Use in California)~~

Apply Trellan H.F.P. as a preplant soil incorporated treatment. Additional weeds tolerant to Trellan H.F.P. may be controlled using overlay preemergence applications of Canopy, Dual, Gemini, Lasso, Lexone, Lorox, Lorox plus, Preview, Pursuit†, Sceptor†† or Sencor or other products registered for preemergence use on soybeans, unless use following Trellan H.F.P. is specifically prohibited by the manufacturer. Consult the manufacturer's labels for application rates, additional weeds controlled, additional use directions and precautions before use.

† The use of Pursuit is limited to certain states. Use Pursuit as an overlay treatment following Trellan H.F.P. only in states specified on the label for Pursuit.

†† Use of Sceptor is limited to certain states. Do not use the overlay preemergence application with Sceptor following a preplant incorporated treatment with Trellan H.F.P. in the "Northern Use Area" as defined by the label for Sceptor.

~~Preplant Incorporated Followed by Postemergence Treatments (Not for Use in California)~~

Apply Trellan H.F.P. as a preplant soil incorporated treatment. Additional weeds tolerant to Trellan H.F.P. may be controlled using postemergence applications of Basagran, Blazer, Classic, Cobra, Galaxy, Pinnacle, Pursuit†, Reflex, Sceptor††, Storm, or Tackle or other products registered for postemergence use on Soybeans, unless use following Trellan H.F.P. is prohibited by the manufacturer. Consult the manufacturer's labels for application rates, additional weeds controlled additional use directions, precautions and limitations before use.

† Use of Pursuit is limited to certain states. Use Pursuit as a postemergence treatment following Trellan H.F.P. only in states specified in the label for Pursuit.

†† Use of Sceptor is limited to certain states. Do not use Sceptor as a postemergence application following a preplant incorporated treatment with Trellan H.F.P. in the "Northern Use Area" as defined by the label for Sceptor.

For broader spectrum weed control, other products registered for use in soybeans may be applied in tank mix combination with Trellan H.F.P. or as a sequential treatment following application of Trellan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

[Editor's note: The above section on tank mixing, overlay and postemergence application was moved to this location from end of "Special Use Programs section" for soybeans.]

530297

Special Use Programs

1. Soybeans - Chemigation

[Editor's note: Chemigation moved to Special Use Programs sections. Also, uses within this section are numbered for sake of clarity.]

Treflan H.F.P. may be applied through properly equipped chemigation systems for weed control in soybeans. Refer to "Application by Chemigation" in the "General Information" section of this label for use directions for chemigation. Do not apply Treflan H.F.P. through any irrigation system unless these directions are carefully followed.

Apply Treflan H.F.P. in sprinkler irrigation equal to 0.5 to 1 inch of water. Planting and application should occur as soon as possible after the last tillage operation. Treflan H.F.P. must be applied within 2 days after planting and prior to crop emergence. Treflan H.F.P. does not control established weeds. Soil incorporation is not required when Treflan H.F.P. is applied through chemigation systems.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5-2.0
Medium	1.5-2.0
Fine	2.0-2.5

- Soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 to 2.5 pints

Cultivation: Soil treated by chemigation with Treflan H.F.P. may be shallow cultivated without reducing weed control activity.

2. Soybeans - Weed Control Under Reduced or Conservation Tillage

[Editor's note: This use added from previously accepted supplemental labeling (EPA-accepted 02/08/93). Proposed changes to previously accepted text are indicated by underline for added text and ~~strike-through~~ for deletions.]

Treflan H.F.P. herbicide can be applied either in the fall or in the spring as a preplant incorporated treatment for weed control in soybeans grown under reduced or conservation tillage conditions. Make only one application per crop cycle.

Make applications to tilled land or standing or chopped stubble from the previous season's crop. Use incorporation equipment that thoroughly and uniformly mixes Treflan H.F.P. into the top 2-3 inches of the final seedbed and leaves a maximum amount of crop residue on the soil surface. Incorporation must occur within 24 hours after application. For fall applications, incorporate once in the fall and a second time in the spring before planting. For spring applications, incorporate twice for best results. The first incorporation of Treflan H.F.P. must occur within 24 hours. For the first incorporation, a tandem disc or combination tool that can thoroughly mix Treflan H.F.P. into the top 2 to 3 inches of the final seedbed while leaving the desired amount of plant residue on the soil surface is recommended. For fall or spring applications, the second incorporation can occur anytime prior to planting or at planting with tillage equipment that provides uniform soil mixing used in conjunction with no-till planters.

Application With Dry Bulk Fertilizers

Dry bulk fertilizers impregnated or coated with Treflan H.F.P. may be applied as preplant incorporated treatments. See instructions for "Application with Dry Bulk Fertilizer" in the General Information section of this label. Under reduced or conservation tillage conditions, uniformly applied dry bulk fertilizers impregnated with Treflan H.F.P. provide weed and grass control equal to or better than Treflan H.F.P. applied in liquid sprays. Two incorporation passes are required when Treflan H.F.P. is applied with dry bulk fertilizer. For best results with spring applications, incorporate once within 24 hours after application and a second time 3 to at least 5 days later.

54 of 97

Application Rates/Acre:

Soil Texture	Treflan H.F.P.	
	Spring Applied	Fall Applied
	(pints)	(pints)
Coarse	1.0 - 1.5	1.5 - 2.0
Medium	1.5 - 2.0	2.0 - 2.5
Fine	2.0 - 2.5	2.5 - 3.0

Use the higher rate in the rate range where higher crop residues are present or where dense weed populations are anticipated.

Precautions

To be effective, Treflan H.F.P. must be mixed thoroughly in the top 2-3 inches of soil in the final seedbed. Weed control may be poor or erratic where soil conditions or heavy crop residues do not permit thorough soil mixing.

3. Soybeans - Fall Panicum Control

Apply Treflan H.F.P. as a preplant incorporated treatment at a broadcast rate of 2.0 pt/acre on coarse and medium soils.

4. Soybeans - Pigweed and Seedling Johnsongrass Control

Apply Treflan H.F.P. as a preplant incorporated treatment.

Broadcast Application Rates/Acre: In Alabama, Arkansas, Florida, Georgia, Kansas, Louisiana, Mississippi, Missouri, Nebraska, North Carolina, Oklahoma, South Carolina, Tennessee, and southern Virginia, apply Treflan H.F.P. at the following broadcast rates:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0-1.5
Medium	1.5-2.0
Fine	2.0-2.5

(exception: Louisiana, 3.0 pt/acre on fine soils).

5. Soybeans - Additional Weed and Grass Control In-Soybeans-(Gulf Coast Counties of Texas)

Apply Treflan H.F.P. as a preplant incorporated treatment up to 2 weeks before planting.

Broadcast Application Rates/Acre: For soybeans grown in Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton counties of the Texas Gulf Coast, apply Treflan H.F.P. at the following broadcast rates:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.5
Medium	2.0
Fine	3.0

6. Soybeans - Itchgrass (Raouigrass) Suppression

Apply Trellan H.F.P. as a preplant incorporated treatment or at layby.

Layby Treatment: Cultivate to remove existing weeds and treat when soybeans are well established (10 inches tall). Apply as a directed spray to the soil surface and incorporate using a rolling cultivator set to cut 2-4 inches deep or sweep-type cultivator with 3 to 5 sweeps per row middle operated 2 to 3 inches deep. Set incorporation equipment to throw treated soil to the row.

Broadcast Application Rates/Acre:

Soil Texture	Trellan H.F.P.	
	Preplant Incorporated	Layby Application
	(pints)	(pints)
Medium	3.0	1.0
Fine	3.0	2.0

7. Soybeans - Charcoal Soils in Arkansas, Louisiana and Mississippi

Newly cleared land often contains high organic matter (5-10%) and charcoal from burning debris. Charcoal and organic matter tends to bind Trellan H.F.P. and reduce weed control activity. Under these conditions, higher rates of Trellan H.F.P. are necessary for weed control. Increased rates, however, can cause crop injury if charcoal or organic matter is not present to bind some of the Trellan H.F.P. In the burn row a high level of charcoal is usually present. Consequently, poor weed control may result, even if an increased rate of Trellan H.F.P. is used. Follow recommended application and incorporation procedures for Trellan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Trellan H.F.P.
	(pints)
Coarse	1.5-2.5
Medium	2.5
Fine	3.0

8. Soybeans - Red Rice Control in Arkansas, Louisiana, Mississippi, and Texas Only

Suppression or partial control of red rice can be obtained from a 2-year treatment program which consists of a double rate application the first year followed by application in the second year at normal rates indicated for soil texture, organic matter or charcoal content. Apply and incorporate Trellan H.F.P. in the spring before planting. Follow recommended soil preparation and incorporation procedures for Trellan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Trellan H.F.P.	
	Application Year 1	Application Year 2
	(pints)	(pints)
Coarse	2.0	1.0
Medium	3.0	1.5
Fine	4.0	2.0
Coarse Soils with 2-5% organic matter	3.0	1.5
Soils with 5-10% organic matter	4.0	2.0-2.5

In Arkansas, Louisiana and Mississippi, if a combination of high soil organic matter (5-10%) and charcoal are present, apply Trellan H.F.P. at the following broadcast rates:

56 of 97

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.5-2.5
Medium	2.5
Fine	3.0

For more information on charcoal soils see discussion in preceding section.

Precaution: Crop Rotation: The recommendation for red rice control in soybeans is a 2-year program. In the first year following a double rate application, plant only soybeans. During the second year, after applying Treflan H.F.P. at the normal rate indicated for soil texture and charcoal level, plant only those crops for which Treflan H.F.P. is registered as a preplant treatment or crop injury may result. Rice may be planted during the third year following application of normal use rates in year two.

9. Soybeans - Rhizome Johnsongrass Control in Eastern United States and the State of Texas

Rhizome johnsongrass control with Treflan H.F.P. requires double rate application for two consecutive years. Commercially acceptable control cannot be obtained with only one year of double rate use of Treflan H.F.P. Carefully follow the special use directions which follow.

Soil Preparation: Satisfactory results are dependent upon proper soil preparation prior to application. Use implements such as a chisel plow to bring rhizomes to the soil surface. Disc twice before application to chop rhizomes into small (2 to 3 inch) pieces and destroy any recently emerged johnsongrass plants.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	2.0
Medium	3.0
Fine	4.0

- Coarse soils with 2-5% organic matter - 3.0 pints
- Soils with 5-10% organic matter - 4.0 pints

Spring Application: Apply Treflan H.F.P. any time before planting in the spring for two consecutive years.

Fall Application: Apply Treflan H.F.P. after October 15 for two consecutive years.

Split Application: Apply Treflan H.F.P. at the broadcast rates indicated in the following table both spring and fall for 2 consecutive years.

Soil Texture	Treflan H.F.P. Spring + Fall (pints)
Coarse	1.0 + 1.0
Medium	1.5 + 1.5
Fine	2.0 + 2.0
Coarse Soils with 2-5% organic matter	1.5 + 1.5
Soils with 5-10% organic matter	2.0 + 2.0

Incorporation: Deep incorporation with a tandem disc is essential for good results. Set disc to operate 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary and the second should be in a different direction than the first.

Cultivation: Some johnsongrass plants will not be controlled. Timely cultivation during the crop season is necessary to remove escaped plants and maintain commercially acceptable control.

57 of 97

Precautions: In the season following a double rate treatment, plant only rice and those crops to which Treflan H.F.P. can be applied as a preplant treatment or crop injury may result.

~~Rhizome Johnsongrass Control with Treflan H.F.P. plus Sencor or Treflan H.F.P. plus Loxone In Tank Mix~~

~~Rhizome Johnsongrass control with Treflan H.F.P. plus Sencor or Loxone requires application for two consecutive years. Apply Treflan H.F.P. plus Sencor or Loxone as a preplant incorporated treatment up to two weeks before planting. This tank mix controls weeds susceptible to Treflan H.F.P. plus additional weeds listed on the label for Sencor or Loxone.~~

Application Rates: ~~See rate recommendations above for "Rhizome Johnsongrass Control in Eastern United States and the State of Texas". Use application rates for soybeans in the label for Sencor or Loxone.~~

Precaution: ~~Refer to the label for Sencor or Loxone for application rates, additional use directions, precautions and limitations prior to applying Treflan H.F.P. plus Sencor or Loxone tank mix. Carefully follow all use precautions on the labels for Sencor or Loxone.~~

10. Soybeans - Wild Cane (Shattercane) Control

Follow recommended soil preparation and application procedures for Treflan H.F.P. Wild cane (shattercane) can germinate throughout the growing season and from greater soil depth than most other weed seeds. Commercially acceptable control of wild cane can be obtained by using increased rates of Treflan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	2.0
Fine	2.5

Incorporation: Deep incorporation with a tandem disc is essential for good wild cane control. Incorporate Treflan H.F.P. thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary with the second in a different direction than the first.

Cultivation: Cultivation during the growing season will improve shattercane control.

SUGAR BEETS

Treflan H.F.P. - Alone

Apply Treflan H.F.P. as an over-the-top spray and incorporate. Apply from the time the first true leaves have formed until plants are 6 inches tall.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.25-1.5

Incorporation: Set incorporation equipment to move treated soil around the plants in the row. Avoid damage to the sugar beet tap root from incorporation equipment.

Precaution: Exposed beet roots should be covered with soil before application of Treflan H.F.P. to reduce the possibility of girdling.

Incorporation with a Tine-Tooth Harrow (For Use in California, Colorado, Idaho, Nebraska, Oregon, Texas, Utah, Washington and Wyoming)

A tine-tooth harrow (Flexline or Melroe) can be used to incorporate Treflan H.F.P. in sugar beets. Incorporation with tine-tooth harrow requires 2 passes in opposite directions over the same set of rows. Set the harrow to cut 1 to 2 inches deep and operate at 3 to 6 mph. Set incorporation equipment carefully to avoid damage to sugar beet tap root. Use application procedures and broadcast application rates recommended in preceding section.

~~Treflan H.F.P. plus Eptam Tank Mix~~

~~Treflan H.F.P. may be tank mixed with Eptam and applied as an over the top spray followed by incorporation to control additional weeds. Use application rates recommended for sugar beets "Treflan H.F.P. - Alone", above. Refer to the label for Eptam for weeds controlled, application rates, additional use directions, precautions and limitations before use.~~

Tank Mixing

For broader spectrum weed control, other products registered for use in sugar beets may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

SUGARCANE

Treflan H.F.P. - Alone

Apply and incorporate Treflan H.F.P. twice a year. Make the first application of Treflan H.F.P. in the fall on firmly packed beds immediately after the seed pieces are planted. Make the second application of Treflan H.F.P. in the spring before or shortly after the cane emerges. Loosen rain-packed beds 2-3 inches deep before the spring application. Take care that incorporation equipment does not damage the seed pieces or emerging shoots.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
All Textures	2.0-4.0†

†Application rate within rate range may be adjusted according to weed pressure.

Postplant Application for Control of Most Annual Grasses, Including Guineagrass (For Use in Hawaii)

Surface apply Treflan H.F.P. after planting (for plant cane) or after harvesting (for ratoon cane). For best results in plant cane, the soil surface should be smooth and finely tilled. Apply Treflan H.F.P. as soon as possible after tillage and planting before germination and emergence of grass weeds. For optimum efficacy in ratoon cane, minimize surface residue from previous crop before applying. Apply Treflan H.F.P. just before anticipated rainfall in non-irrigated and furrow-irrigated sugarcane. Irrigate as soon as possible after applying in drip-irrigated or sprinkler-irrigated sugarcane to activate Treflan H.F.P. Apply 0.5 inch or more irrigation in drip-irrigated or sprinkler-irrigated sugarcane as soon as possible after applying Treflan H.F.P.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
All Textures	6.0-8.0

59 of 97

Repeat Applications:

[Editor's note: From previously accepted supplemental labeling (EPA-accepted 10/23/91.)

Subsequent germination of grass weeds may occur prior to the development of a full dense canopy of sugarcane. If Once this occurs, additional grass weed establishment is strongly suppressed. One or two additional applications of Treflan H.F.P. can be applied to maintain weed control during the early crop development period. For repeat applications, direct the spray to the soil surface to minimize interception of the herbicide by the crop.

Restrictions:

- Do not apply Treflan H.F.P. as a postplant surface applied treatment within 180 days of harvest.

Applications Up to Layby for Plant Cane or Ratoon Cane (For Use in Louisiana and Texas)

Apply and incorporate Treflan H.F.P. in spring from shortly before or after cane emergence until layby. Apply after beds have been shaved or false shaved. Loosen rain-packed beds 2 to 3 inches deep before application. Avoid incorporation equipment damage to seed pieces or emerging shoots. Incorporate with a rolling cultivator or bed chopper for all soil textures. Set rolling cultivator to cut 2 to 4 inches deep and operate at 6 to 8 mph. Set bed chopper to cut 3 to 4 inches deep and operate 4 to 6 mph. Two incorporation passes are necessary.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
All Textures	2.0-4.0†

†Application rate within rate range may be adjusted according to weed pressure.

Itchgrass (Raoulgrass) Control (For Use in Louisiana)

Apply and incorporate Treflan H.F.P. on plant or ratoon cane. Follow use directions in preceding section for layby application.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
All Textures	2.0-4.0

SUNFLOWERS

Treflan H.F.P. - Alone

Apply and incorporate Treflan H.F.P. in the spring before planting or in the fall in advance of spring planting. See instructions for fall application under "Application Timing" in the "General Information" section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 to 2.0 pints
- Fine soils with 2-5% organic matter - 2.0 pints.
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

60097

Treflan H.F.P. plus Eptam Tank Mix

Treflan H.F.P. may be tank mixed with Eptam and applied as a preplant incorporated treatment to control additional weeds in sunflowers in Minnesota, North Dakota and South Dakota. Refer to the label for Eptam for application rates, additional use directions, precautions and limitations before use.

Tank Mixing

For broader spectrum weed control, other products registered for use in sunflowers may be applied in tank mix combination with Treflan H.F.P. or as a sequential treatment following application of Treflan H.F.P. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

TOMATOES

Apply to direct-seeded tomato as a directed spray between rows and beneath plants and incorporate at the time of blocking or thinning. For transplant tomatoes, apply and incorporate before transplanting. Do not apply after transplanting or apply post-plant as a directed spray to the soil between the rows and beneath plants and incorporate.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- Coarse and medium soils with 2-5% organic matter - 1.5 pints
- Fine soils with 2-5% organic matter - 2.0 pints
- Soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

TREE AND VINE CROPS - CITRUS, FRUIT AND NUT CROPS TREES AND VINEYARDS

Application to New Plantings of Citrus, Fruit and Nut Crops Trees

For new plantings of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, plum, prune, tangelo, tangerine and walnut trees, a Apply and incorporate Treflan H.F.P. before transplanting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
Coarse	1.0
Medium	1.25-1.5
Fine	1.5-2.0

- All soils with 2-5% organic matter - 1.5-2.0 pints
- All soils with 5-10% organic matter - 2.0 pints
- Use lower rate in rate range in areas receiving less than 20 inches total rainfall and irrigation.

Application to New Plantings of Vineyards

For new plantings of vineyards, a- Apply and incorporate Treflan H.F.P. before transplanting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
Coarse	1.0-1.5
Medium	1.5-3.0
Fine	3.0-4.0

- Soils with 2-10% organic matter - 4.0 pints
 - Use lower rate in rate range in areas receiving less than 20 inches total rainfall and irrigation.
- Note: Do not use more than 2.0 pt/acre on heat-treated mist propagated grape rootings.

Application to Established Non-bearing and Bearing Citrus, Fruit and Nut Crops Trees and Vineyards

Treflan H.F.P. may be applied in established non-bearing and bearing vineyards and plantings of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, plum, prune, tangelo, tangerine and walnut trees. In established plantings, apply Treflan H.F.P. as a directed spray to the soil and use incorporation incorporated using methods not injurious to the crop. Do not apply to vineyards within 60 days of harvest.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.
	(pints)
All Soil Textures	2.0-4.0

Application rate within the rate range may be adjusted according to weed pressure.

Rhizome Johnsongrass Control - Special Two-year Use Program

Treflan H.F.P. may be applied for two consecutive years in a special use program to control rhizome johnsongrass in established vineyards and citrus, tree fruit and nut tree plantings. of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, plum, prune, tangerine and walnut trees. Do not apply to vineyards within 60 days of harvest.

Soil Preparation: Work the soil thoroughly to move rhizomes near the soil surface and cut them into smaller pieces.

Broadcast Application Rates/Acre:

The following application rate must be applied for two consecutive years:

Soil Texture	Treflan H.F.P.
	(pints)
All Soil Textures	4.0

Incorporation: Incorporate Treflan H.F.P. thoroughly with a disc set to cut 4 to 6 inches deep and operate 4 to 6 mph. Two incorporation passes are necessary, with a second pass in a different direction from the first.

Cultivation: Some johnsongrass plants will escape. Timely cultivations are necessary to obtain commercially acceptable control. Commercially acceptable control cannot be obtained with only a single year use of Treflan H.F.P.

Precautions: Do not use the 4 pint rate on new plantings or crop injury may result. Do not interplant orchards or vineyards with other crops. If treated vineyards and orchards are diverted to other crop uses, then in the next cropping season plant only those crops for which Treflan H.F.P. has been registered as a preplant incorporated treatment.

62 of 97

Bindweed Control in California

Treflan H.F.P. can be applied using a specially equipped spray blade for the control of field bindweed in vineyards and citrus, tree fruit and nut tree plantings, of almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees.

Soil Preparation: Destroy existing weeds with soil tillage before applying Treflan H.F.P. Thorough tillage is necessary to prevent trash from interfering interference with operation of the spray blade.

Equipment: Application requires a spray blade capable of operation at 4 to 6 inches below the soil surface. The blade should be equipped with nozzles located under the blade and directed so as to allow spray to be trapped in a thin layer as the blade is pulled through the soil. Use a nozzle spacing sufficient to insure application of a uniform horizontal layer.

Application: Apply Treflan H.F.P. in 40 to 80 gallons of water per acre. Operate blade at a depth of 4 to 6 inches.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. (pints)
All Soil Textures	4.0

Precautions: Some soils may develop cracks as they dry after rainfall or irrigation. Field bindweed may emerge if the cracks extend through the layer of Treflan H.F.P. Prevent or eliminate cracks by shallow discing or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

*Trademark of DowElanco

63 of 97

[Base Label of Laminated Booklet -- Treflan H.F.P.]

(logo) DowElanco

Treflan* H.F.P.

A selective herbicide for the preemergence control of annual grasses and broadleaf weeds.

Active Ingredient:

trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine.....	43.0%
Inert Ingredients	57.0%
Total	100.0%

Contains 4 pounds active ingredient per gallon.

Manufactured using the processes of U.S. Patents 4,120,905 and 4,226,789

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin • Treflan H.F.P. May Cause Skin Sensitization Reactions In Certain Individuals

Avoid breathing vapors or spray mist and contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate or Viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-C)], the handler PPE requirements may be reduced or modified as specified in the WPS.

64 of 97

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not induce vomiting or give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation assistance, and get medical attention immediately.

If on skin: Immediately wash with plenty of soap and water. Get medical attention if irritation develops.

Note to physician: This product contains an aromatic hydrocarbon and can be extremely harmful if swallowed. Aspiration of this product may produce a severe pneumonitis. Stomach lavage with a cuffed endotracheal tube in place and immediate administration of activated charcoal, 6-8 heaping teaspoonfuls with water, should be considered. Treatment is otherwise symptomatic and supportive.

Environmental Hazards

This pesticide is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE), Engineering Controls Statements and User Safety Recommendations, and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Avoid Freezing -- Store Above 40°F

EPA Reg. No. 62719-ELN

EPA Est. 1471-IN-2

*Trademark of DowElanco

DowElanco • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents 2.5 gal

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054, USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

For Distribution and Use Only in the State of Texas

Fall Application to Fine Textured Soils for Weed Control in Cotton to be Grown the Following Year

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply Treflan® herbicide to fine textured soils in the fall between after September 1 and October 15 for weed control in Cotton to be grown the following year. Fine textured soils include clay, clay loam, silty clay loam†, silty clay, sandy clay, and sandy clay loam† soil textures. Follow application and incorporation directions on the label for Treflan.

Broadcast Application Rates:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan TR-10
	(pints/acre)	(pints/acre)	(pounds/acre)
Fine Soils Only	4.0	3.2	20.0

†Silty clay loam and sandy clay loam soils are transitional and may be classified as either medium or fine textured soils if they contain predominantly silt or sand, respectively. For details, refer "Soil Texture Guide for Application Rates" in the "General Information" section of the label for Treflan.

*Trademark of DowElanco

122-T2EAG040

Approved: Treflan E.C. 10/23/91

Treflan M.T.F. 9/25/91

Treflan 5 10/11/91

Treflan TR-10 9/23/91

Replaces EA 7141

Revisions:

1. Fall application period extended from October 15 to October 31.
2. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trade mark;
grammar; punctuation; or use of u/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan* H.F.P. Herbicide (EPA Reg. No. 62719-ELN)~~
~~Treflan E.C. Herbicide (EPA Reg. No. 62719-93)~~
~~Treflan M.T.F. Herbicide (EPA Reg. No. 62719-116)~~
~~Treflan 5 Herbicide (EPA Reg. No. 62719-118)~~
(For Distribution and Use Only in the State of Montana)

Spring Applied Treflan for Foxtail (Pigeongrass) Control in Spring Seeded Barley Grown Under Irrigation

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan* herbicide may be spring applied as a preplant incorporated treatment for foxtail (pigeongrass) control in spring seeded barley grown under irrigated conditions in Montana. Treflan may be applied to ground that has a manageable trash level or has been fallowed or pretilled. The first incorporation is required within 24 hours after application. The second incorporation is required prior to planting to destroy emerged weeds and to ensure even distribution of Treflan in treated soil.

Broadcast Application Rate: Apply Treflan ~~H.F.P., E.C., and M.T.F.~~ at a rate of 1 pint per acre and ~~Treflan 5~~ at a rate of 0.8 pint per acre regardless of soil texture or soil organic matter content. Do not exceed this application rate as crop injury may occur.

Incorporation Directions

The following tools are recommended for soil incorporation:

1. **Chisel Plow alone or Chisel Plow with a Rod Weeder attached:** A chisel plow alone should be used for the first incorporation pass only. With rod weeder attached, the chisel plow may be used for both incorporation passes. Operate 4 to 5 inches deep and at 4 to 6 mph. A chisel plow is defined as having three rows of up to 18 inch sweeps on no greater than 12 inch centers. Stagger successive rows of sweeps to ensure that no soil is left unturned.
2. **Tandem Disc:** Operate 3 to 4 inches deep and at 4 to 6 mph.
3. **Field Cultivator:** Operate 3 to 4 inches deep and at 5 or more mph. A field cultivator is defined as having 3 to 4 rows of sweeps with "C" or "S" shaped shanks spaced at intervals of 7 inches or less. Stagger successive rows of sweeps to ensure that no soil is left unturned.

Planting Directions

Plant barley 1 to 2 inches deep. Planting greater than 2 inches deep will result in increased seedling stress and decreased emergence.

Irrigation Directions

Irrigate prior to planting, or after crop emergence only. Irrigation between planting and emergence may cause reduced crop stands or delayed emergence because of soil crusting, especially on loose friable seedbeds.

Use Precautions: Carefully follow Special Use Precautions for Small Grains in main product label. Carefully follow use directions to minimize potential crop stress. Under certain conditions, delayed crop emergence and/or stand reduction may occur when Treflan is applied to barley. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield. For best results, observe the following cultural practices or precautions:

- Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.
- Do not exceed recommended application rates for Treflan. This is particularly important on coarse textured or low organic matter soils.
- Use only high quality seed where Treflan is to be applied (avoid use of small seed with low starch reserves).
- If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.
- Avoid use of seed varieties known to have poor seedling (emergence) vigor.
- Do not fall apply Treflan in combination with any other preplant incorporated herbicide.

Soil characteristics and environmental conditions which may contribute to crop seedling stress that may be accentuated by use of Treflan E.C. include:

- Soil related: High salinity, eroded knolls/hilltops, loose dry soils and compaction.
- Weather related: Cold and/or wet soils, excessively hot soils, excessive moisture, drought, and soil crusting from heavy rainfall.

Note: Do not apply Treflan on small grains where a dinitroaniline herbicide such as Treflan was applied at a rate recommended for row crops (oil seeds) during the previous growing season.

Rotational Crop Planting Restrictions

Plant only barley (grown under irrigated conditions), rapeseed, safflower or sunflower as a rotational crop in the year following the crop treated with Treflan. If one of the specified rotational crops is not planted, the land should be left idle or fallow for the entire crop year following the crop treated with Treflan.

*Trademark of DowElanco

122-T2EAG048 Approved: Treflan E.C. 10/23/91;
Treflan M.T.F. 07/5/91; Treflan 5 6/18/91
Initial printing.

Amendment(s):

1) Special Use Precautions for Small Grains deleted
from supplemental labeling. Users now referred to
main product label for this information.

68097

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Treflan plus Pursuit Combinations for Weed Control in Spanish Peanuts or Florunner and Florigiant Varieties

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide may be tank-mixed with Pursuit and applied as a preplant incorporated treatment to control additional weeds. Follow application and incorporation directions provided in the label for Treflan.

Broadcast Application Rates/Acre†:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5
	(pints)	(pints)
Coarse	1.0	0.8
Medium	1.5	1.2

†Refer to the labeling for use of Pursuit on peanuts for application rates.

Pursuit may also be used as a preemergence, "at cracking", postemergence or sequential (split) application following preplant soil incorporated application of Treflan. Refer to the labeling for use of Pursuit on peanuts for application rates, use directions, cautions and limitations before use.

*Trademark of DowElanco

122-T2EAG049

Approved: Treflan E.C. 10/23/91

Treflan M.T.F. 9/25/91

Treflan 5 10/11/91

Initial printing.

Amendment(s):

1) Removed limitation to use on certain varieties of peanuts.

2) Minor editing to correct (one or more may apply): reference to "Treflan" as a DowElanco trademark; grammar; punctuation; or use of w/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan® E.C. (EPA Reg. No. 62719-93)

Treflan® M.T.F. (EPA Reg. No. 62719-116)

Treflan® 5 (EPA Reg. No. 62719-118)

Treflan® TR-10 (EPA Reg. No. 62719-131)

(For Distribution and Use Only in the State of Montana)

Treflan for Weed Control in Rapeseed (Canola), Crambe Safflower and Sunflower

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply and incorporate Treflan® herbicide in the fall after September 1 or in the spring before planting. Make only one application of Treflan per crop cycle. Follow soil preparation, application and incorporation instructions in the product label for Treflan.

Broadcast Application Rates Per Acre:

Soil Texture	<u>Treflan H.F.P., Treflan E.C. and Treflan M.T.F.</u>	<u>Treflan 5</u>	<u>Treflan TR-10</u>
	(pints)	(pints)	(pounds)
Coarse	1.0	0.8	5.0
Medium	1.5	1.2	7.5
Fine	2.0	1.6	10.0

Use 1.5 to 2.0 pints of Treflan H.F.P., E.C. and M.T.F., 1.2 to 1.6 pints of Treflan 5, or 7.5 to 10 pounds of Treflan TR-10 per acre on coarse and medium soils with 2-5% organic matter.

Precautions

Rotational Crop Planting Restriction: Plant only spring seeded barley (grown under irrigated conditions), rapeseed, safflower or sunflower as rotational crops in the crop year following the crop treated with Treflan. If one of these specified crops is not planted, the land should be left idle or fallow for the entire crop year following the crop treated with Treflan.

Amendment(s)

- 1) Use on crambe, approved by EPA on 03/30/93, added to supplemental labeling.
2. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark; grammar; punctuation; or use of w/ case.

*Trademark of DowElanco

122-33-061

Replaces 122-T2EAGO44

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Treflan Plus Tri-Scept Tank Mix for Control of Fall Panicum, Seedling Johnsongrass and Shattercane in Soybeans

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide should be tank mixed with Tri-Scept herbicide and applied as a preplant incorporated treatment where infestations of fall panicum, shattercane and seedling johnsongrass are anticipated. Use application and incorporation methods for Treflan applied as a preplant incorporated treatment.

Broadcast Application Rates per Acre:

In addition to the recommended rate per acre of Tri-Scept (2.33 pt/acre), add the indicated amount of Treflan to control the weed species indicated.

Weed Species	Soil Textures	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5
		(pints)	(pints)
Fall Panicum	All	0.5	0.4
Johnsongrass (seedling only)	Medium	0.5	0.4
	Fine	0.5	0.4
Shattercane	Medium	0.5	0.4
	Fine	1.0	0.8

General Use Precautions:

- The use of Tri-Scept is limited to those states or portions of states listed on the label for Tri-Scept. Do not use the tank mix of Treflan plus Tri-Scept in the "Northern Use Area" as defined by the label for Tri-Scept.
- Do not graze or feed treated soybean forage, hay, or straw to livestock.
- Do not apply Tri-Scept postemergence to soybeans as crop injury may occur.
- Crops other than soybeans may be injured by spray, drift or other indirect contact with Tri-Scept.
- To avoid injury to sensitive crops, spray equipment used for applications of Tri-Scept must be drained and thoroughly cleaned with water before being used to apply other products to the following crops: cotton, corn, grain sorghum, rice and vegetables.
- A preharvest interval of 90 days is required between the last application of Tri-Scept and the soybean harvest.

7/10/97

Replanting: If replanting is necessary in a field previously treated with Tri-Scept, replant only soybeans. Rework soil no deeper than the treated zone. Do not reapply Tri-Scept.

All additional use directions, precautions and limitations applicable to the use of Tri-Scept, apply to the use of this tank-mix.

Rotational Crop Restrictions for Tri-Scept:

Use of Tri-Scept herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of Tri-Scept and therefore, rotational crop injury is always possible.

Certain severe restrictions apply to corn, wheat and other rotational crops following an application of Tri-Scept. Refer to the label for Tri-Scept for complete details of these and other restrictions.

1. Do not plant wheat within 4 months following an application of Tri-Scept in soybeans; and, do not plant rice until the spring of the year following an application of Tri-Scept.
2. Do not plant barley, field corn, edible beans, grain sorghum, oats, peanuts or tobacco within 11 months after an application of Tri-Scept.
3. Do not plant sugar beets for 26 months after an application of Tri-Scept.
4. Field corn planting restrictions: For the "Southern Use Area" as defined by the label for Tri-Scept; field corn may be planted as a rotational crop in the spring of the year following an application of Tri-Scept unless extreme drought conditions develop or less than 15 inches of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the Bootheel of Missouri, Tennessee, North Carolina and states south: field corn may be planted in the spring of the year following an application of Tri-Scept unless extreme drought conditions develop or less than 15 inches of rainfall or irrigation is received within 6 months following the date of application.
5. Do not plant cotton or rotational crops other than those listed above within 18 months following an application of Tri-Scept.
6. Do not apply products containing chlorimuron ethyl (e.g., Classic, Canopy, Gemini, Lorox Plus, Preview, etc.) or imazethapyr (e.g., Pursuit, or Pursuit Plus) the same year as Tri-Scept or injury to following crops may occur.

Refer to the labels for Trellan and Tri-Scept for additional use directions, precautions, and limitations before use.

*Trademark of DowElanco

T2EAG017 Approved 12/06/90

Revisions:

1. Added text box containing required FIFRA and DowElanco standard statements and limitations.
2. Minor editing to correct (one or more may apply):
reference to "Trellan" as a DowElanco
trademark; grammar; punctuation; or use of w/i
case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Treflan Plus Canopy and Crop Oil Concentrate Early Preplant Application for Weed Control in No-till Soybeans (Not For Use In California)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow use precautions and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all precautions and limitations imposed by the label affixed to the container for Treflan.

Directions For Use

Apply Treflan® herbicide plus Canopy herbicide and crop oil concentrate as an early preplant surface applied treatment to control existing weeds and provide preemergence control of susceptible weeds during the cropping season in full season, narrow-row no-till soybeans. The tank mix of Treflan plus Canopy with crop oil concentrate should be surface applied as a broadcast spray at least three weeks before planting. For best results, soybean row spacing should be 15 inches or less and planting should occur before May 15. Rainfall of at least 0.5 inches is required within 3 to 5 days after application to activate Treflan.

Broadcast Application Rates per Acre:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Canopy 75DF¹
	(pints/A)	(pints/A)	(ounces/A)
Coarse	2.0	1.6	6
Medium	3.0	2.4	8
Fine	4.0	3.2	10

¹Use crop oil concentrate at a rate of 1 quart per acre regardless of application rate.

Use of Additional Postemergence Herbicides:

In addition to Canopy plus crop oil concentrate, other postemergence herbicides, may be added to the tank mix to enhance control of existing weeds. These include 2,4-D, Gramoxone Extra and Roundup. Refer to the labels of these additional tank-mix products for use rates.

Precautions:

- The user must comply with all applicable use directions, precautions and limitations imposed by the labels for Treflan and the tank mix product.
- Read and understand these additional use directions, precautions and limitations before use.

*Trademark of DowElanco

122-T2EAG018 Approved 12/06/90

Revisions:

1. Added text box containing required FIFRA and DowElanco standard statements and limitations.
2. Minor editing to correct (one or more may apply): reference to "Treflan" as a DowElanco trademark; grammar; punctuation; or use of u/l case.

Supplemental Labeling



73 of 97

DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

(For Distribution and Use Only in Texas, Oklahoma and New Mexico)

Treflan 48 Hour Incorporation Delay

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan, except as described below.

Treflan® herbicide may be applied as a preplant incorporated treatment for weed control in registered crops. The incorporation delay for Treflan is extended from 24 to 48 hours when applied in the states of Texas, Oklahoma and New Mexico. If Treflan is applied to a warm wet soil surface or the wind velocity is 10 mph or greater, variable weed control may result if the first incorporation is delayed more than 24 hours.

Follow soil incorporation procedures recommended on the label for Treflan. Where two incorporation passes are required, the first pass must be accomplished within 48 hours after application. The second incorporation pass may occur anytime before planting.

*Trademark of DowElanco

122-33-025 Approved 03/20/91
Replaces T2EAG024 and T2EAG025.

Revisions:

1. Added boxed text including EPA and DowElanco required statements and limitations.
2. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of u/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

**Treflan for Partial Control or Suppression of Annual Brome Species
(Cheatgrass, Downy Brome, Japanese Brome, Hairy Chess)
and Jointed Goatgrass in Winter Wheat in Colorado, Kansas, Nebraska,
Oklahoma, South Dakota, Texas and Wyoming**

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide may be applied as a preplant incorporated treatment for partial control or suppression of annual Brome species (cheatgrass, downy brome, Japanese brome, hairy chess) and jointed goatgrass in winter wheat in Colorado, Kansas, Nebraska, and Wyoming. Apply Treflan anytime during a period from three (3) weeks before planting up to immediately prior to planting.

Broadcast Rates Per Acre¹

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F. (pints)	Treflan 5 (pints)	Treflan TR-10 (pounds)
Course	1 - 1 1/2	0.8 - 1.2	5.0 - 7.5
Medium	1 - 1 1/2	0.8 - 1.2	5.0 - 7.5
Fine	1 1/2	1.2	7.5

¹Use the higher rate in the rate range where heavy weed populations are anticipated or where medium to high crop residues are present.

Incorporation and Planting Directions: Treflan should be incorporated with tillage equipment such as a flexible tine-tooth harrow or springtooth harrow that mixes the soil no more than 1-2 inches deep. The grain drill (disc drill or hoe drill) can serve as the incorporation implement. Do not use discs, undercutters or field cultivators for incorporation. Incorporate one time within 24 hours after application. Use a grain drill that will place the seed below the zone of soil into which Treflan has been incorporated.

One pass incorporation is adequate; however, where the grain drill is used as the incorporation tool, mounting a springtooth harrow in front of the drill can enhance performance. Where a tillage tool is used to incorporate prior to planting, the wheat must be seeded below the soil treated with Treflan or crop injury may result. The wheat seed should be placed at least 1 1/2 - 2 inches deep.

(continued on back)

750497

Precautions:

- Crop injury in the form of delayed emergence and development may result from planting wheat in direct contact with treated soil.
- Do not incorporate with undercutters, field cultivators, chisel plows or discs. Any implement that incorporates Treflan deeper than the planting depth of wheat will contribute to crop injury.
- Use of seeding equipment that does not place the seed below the Treflan treated soil layer will result in crop injury.
- Use of Treflan in accordance with this label may result in stand reduction.
- Heavy rainfall prior to wheat emergence can cause soil compaction and soil crusting resulting in delayed emergence, stand reduction, stunting and yield loss.

*Trademark of DowElanco

122-T2EAG032 Approved 09/01/89
Not distributed.

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of u/l case.

Supplemental Labeling



76 of 97

DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan E.C. (EPA Reg. No. 62719-93)

Treflan 5 (EPA Reg. No. 62719-118)

(For Distribution and Use Only in the State of Florida)

Treflan Applied in Irrigation Water Rings for Weed Control in Non-bearing Citrus Trees

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow use precautions and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all precautions and limitations imposed by the label affixed to the container for Treflan.

Treflan® herbicide may be applied to non-bearing citrus trees through irrigation water rings to provide preemergence control of labeled weeds.

Directions for Use

Mixing: Mix at a rate of 12 fluid ounces of Treflan H.F.P., Treflan E.C. or 10 fluid ounces of Treflan 5 per 500 gallons of water. Agitate until uniformly dispersed in tank.

Application: Apply 10 gallons of the Treflan herbicide mixture per four foot diameter water ring per tree. Application should be made at the second or third watering and should not be applied in combination with any other pesticide.

*Trademark of DowElanco

122-T2EAG041

Approved: Treflan E.C. 10/23/91

~~Treflan 5 9/16/91~~

Replaces EA1134 6/84

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of w/ case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Treflan Plus Scepter Combinations for Weed Control in Soybeans (Not For Use in California)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide plus Scepter herbicide may be applied as a preplant incorporated tank mix treatment. Scepter may also be applied as a preemergence overlay treatment or as an early postemergence treatment following a preplant incorporated application of Treflan.

Special Use Precautions

- Use of Scepter in combination with Treflan or as a sequential treatment following application of Treflan is subject to a number of special precautions and limitations required by the label for Scepter. Carefully read, understand and follow all use precautions, rotational crop restrictions and other limitations in the label for Scepter.
- Use of Scepter is limited to certain states. The following additional restriction is imposed by this supplemental labeling: Do not use the Treflan plus Scepter tank mix applied preplant incorporated, or Scepter applied as a preemergence overlay or postemergence treatment following Treflan preplant incorporated, in the "Northern Use Area" as defined by the Scepter product label.
- The user must comply with all applicable use directions, precautions and limitations imposed by the Treflan and Scepter product labels.

Tank Mix of Treflan plus Scepter - Preplant Incorporated

Apply the tank mix of Treflan plus Scepter as a preplant incorporated treatment up to 45 days before planting soybeans. Apply and shallowly incorporate into the soil within 24 hours. Follow soil preparation, mixing and application procedures described in the label for Treflan. Use incorporation equipment that provides uniform incorporation into the top 2 inches of soil. When using a disc, field cultivator, or rolling cultivator, a second pass must be made at an angle to the first pass to ensure thorough incorporation. If soybeans are planted on beds, apply and incorporate after bed formation.

Treflan Preplant Incorporated Followed by Scepter Preemergence Overlay Application

Apply Treflan as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures described in the label for Treflan. Apply Scepter to the soil surface after planting but before the crop emerges. Rainfall or overhead sprinkler irrigation is necessary to move the Scepter overlay treatment into the weed germination zone. The amount of rainfall or continued or back sprinkler irrigation required depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after a surface applied treatment, a cultivation is recommended to control escaped weeds.

(Continued on back)

Weeds Controlled (Tank Mix or Overlay)

The tank mix of Treflan plus Scepter or Treflan followed by overlay treatments of Scepter control the weeds listed on the label for Treflan alone plus these additional weeds:

cocklebur, common	poinsetta, wild
jimsonweed	prickly sida
mallow, venice	(teaweed)
morningglory	ragweed
(pitted)	(common)
(smallflower)	(giant) ¹
mustard species	smartweed
nightshade, eastern black ¹	(ladysthumb)
pigweed	(pennsylvania)
Palmer)	sunflower, common
(smooth)	velvetleaf ¹
(tall waterhemp)	

¹ Eastern black nightshade, giant ragweed, and velvetleaf are controlled by preplant incorporated treatments only.

Treflan plus Scepter tank mix or Scepter overlay treatments will aid in the control and reduce competition from weeds in the following list. Control of these weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of weed germination, depth of weed seed in soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

morningglory ²	nutsedge, yellow ³
(entireleaf)	panicum, fall ⁴
(ivyleaf)	shattercane ⁴
(tall)	

²For best activity on morningglory species, use tank mix preplant incorporated treatments.

³Use tank mix preplant incorporated treatments only to aid in control of yellow nutsedge.

⁴Treflan alone control fall panicum and shattercane at increased rates. See the label for Treflan for special instructions.

Broadcast Application Rates for Preplant Incorporated or Preemergence Overlay Treatments:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan-S	Scepter Tank Mix or Overlay
	(pints)	(pints)	(pints)
Coarse	1.0	0.8	0.67
Medium	1.5	1.2	0.67
Fine	2.0	1.6	0.67

Treflan Preplant Incorporated Followed by Scepter Postemergence Application

Apply Treflan as a preplant incorporated treatment. Additional weeds tolerant to Treflan may be controlled using a post emergence application of Scepter. Consult the Scepter product label for application rates, additional weeds controlled, application directions and precautions before use.

*Trademark of DowElanco

122-T2EAG048

Approved: Treflan E.C. 10/23/91

Treflan M.T.F. 9/25/91

Treflan-S 10/11/91

Replaces EA 9276

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammatical punctuation or use of w/case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan® E.C. (EPA Reg. No. 62719-93)

Treflan M.T.F. (EPA Reg. No. 62719-116)

Treflan 5 (EPA Reg. No. 62719-118)

(For distribution and use only in the states of Arizona and California)

Treflan 72 Hour Incorporation Delay

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

When Treflan® herbicide is applied as a preplant incorporated treatment, the first incorporation pass must be accomplished within 24 hours after application. In Arizona and California the incorporation delay has been extended from 24 to 72 hours when applied to dry soils. However, when Treflan is applied to warm soil or if wind velocity is 10 mph or higher, variable weed control may result from delaying the first incorporation beyond 24 hours.

Where two incorporation passes are required, the second incorporation may occur anytime prior to planting. Follow other recommended incorporation directions on the label for Treflan.

*Trademark of DowElanco

122-33-063 Approved 08/29/92
Replace EA5059

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco
trademark; grammar; punctuation; or use of "I/
case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan E.C. (EPA Reg. No. 62719-93)

Treflan M.T.F. (EPA Reg. No. 62719-116)

Treflan 5 (EPA Reg. No. 62719-118)

(For distribution and use only in the state of California)

Treflan Application Over Standing or Shredded Cotton Stalks After September 1 for Weed Control in Cotton

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide may be broadcast applied, after September 1, over the top of standing cotton stalks or after shredding using ground or aerial equipment. Treflan must be incorporated once within 24 hours after application using a disc operated at 4 to 6 mph. The second incorporation may be delayed until spring prior to bedding. To avoid dilution of the herbicide or bringing untreated soil to the surface, the treatment should not be chiseled, ripped or deep plowed following incorporation.

Refer to the label of the Treflan product used for details on fall application, soil preparation, application rates and weeds controlled.

*Trademark of DowElanco

122-33-064 Approved 06/29/92
Replaces EA2387

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of u/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~
~~Treflan E.C. (EPA Reg. No. 62719-93)~~
~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~
~~Treflan 5 (EPA Reg. No. 62719-118)~~

(For distribution and use only in the state of Texas)

Incorporation with the Springtooth Harrow on Coarse Textured Soils to be Bedded up Prior to Planting

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Implement Definition: A springtooth harrow is defined as an implement with 3 to 4 rows of shanks equipped with chisel points spaced at intervals of 7 inches or less and staggered so that no soil is left unturned.

Directions For Use: The springtooth harrow may be used to effectively incorporate (mix) Treflan® herbicide into coarse textured (sandy) soils to be bedded up prior to planting.

Soil Preparation: Destroy existing weeds before an application of Treflan. Chop and thoroughly mix crop residues into the soil to a depth of at least 4 to 6 inches by deep plowing or discing prior to an application of Treflan. Use machinery that breaks up large clods before an application of Treflan.

Springtooth Harrow Use Directions: Treflan must be incorporated the first time within 24 hours after application. Set the springtooth harrow to cut 3 to 4 inches deep and operate at a speed of 5 mph or greater. Two passes over the field are required with the second pass in a different direction than the first. The springtooth harrow also may be used as the first or second incorporation tool in combination with other recommended equipment for the other incorporation.

When Treflan is applied and incorporated before bedding, do not furrow out deeper than the depth to which Treflan was incorporated. Furrowing too deep will expose untreated soil and allow weeds to germinate in the bottom of the furrow.

Precautions

- Do not incorporate with springtooth harrow if soil is too wet for good mixing.
- Avoid removal of treated soil from seedbed during planting operation since exposure of untreated soil will allow weeds to grow.

*Trademark of DowElanco

122-33-065 Approved 06/29/92
Replaces EA1154

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of w/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan E.C. (EPA Reg. No. 62719-93)

Treflan M.T.F. (EPA Reg. No. 62719-116)

Treflan 5 (EPA Reg. No. 62719-118)

Weed Control in Potatoes (Not for Use in the State of Maine)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Application After Planting

Apply and incorporate Treflan® herbicide after planting but before emergence, immediately following dragoff, or after potato plants have fully emerged.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5
	(pints)	(pints)
Coarse	1.0	0.8
Medium	1.25-1.5	1.0-1.2
Fine	1.5-2.0	1.2-1.6

- Coarse and medium soils with 2-5% organic matter - 1.5 pints of Treflan ~~H.F.P., E.C. or M.T.F.~~, or 1.2 pints of Treflan 5
- Fine soils with 2-5% organic matter - 2.0 pints of Treflan ~~H.F.P., E.C. or M.T.F.~~, or 1.6 pints Treflan 5
- Soils with 5-10% organic matter - 2.0 pints of Treflan ~~H.F.P., E.C. or M.T.F.~~, or 1.6 pints of Treflan 5
- Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Incorporation Directions: Set incorporation equipment so that the bed and furrow will be uniformly covered with a layer of treated soil. If the layer of treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. When applying and incorporating Treflan after potato plants have fully emerged, do not completely cover the foliage with treated soil. Likewise, do not completely cover foliage at subsequent cultivations. Be careful that incorporation machinery does not damage potato seed pieces or elongating sprouts.

Split Applications Before and After Planting (For use in Idaho, Oregon and Washington)

On all soils apply and incorporate Treflan at the rates shown below as split applications before planting and after planting when potato plants have fully emerged. Do not apply to soils containing 2% or more organic matter. Follow incorporation directions provided above for application to potatoes after planting.

(continued on back)

83 of 97

Broadcast Application Rates/Acre

	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan-5
	(pints)	(pints)
Before Planting	0.75	0.6
After Planting	0.75	0.6

**Treflan Plus Eptam herbicide Tank-Mix - Post Plant Preemergence Treatment
(For Use in Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas)**

Treflan may be tank-mixed with Eptam herbicide and applied as a soil incorporated treatment to control additional weeds. Apply after planting, but before crop emergence. In areas where potatoes are normally dragged off, apply and incorporate up to or immediately following drag off. Use application rates for Treflan recommended for "Applications After Planting", above. Incorporate immediately.

Precautions: Refer to the label for Eptam for application rates, additional use directions, precautions and limitations before use. Do not graze for feed forage to livestock from fields treated with the Treflan plus Eptam tank mix.

**Treflan Plus Eptam Tank-Mix - Preplant Treatment
(For Use in Idaho, Oregon and Washington)**

Treflan may be tank-mixed with Eptam and applied as a soil incorporated treatment to control additional weeds. Apply before planting and incorporate immediately.

Broadcast Application Rates/Acre

	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan-5
	(pints)	(pints)
All soil textures	0.75	0.6

Precautions: Do not use this tank mix both before and after planting in the same season. Do not graze for feed forage to livestock from fields treated with the Treflan plus Eptam tank mix. Refer to the label for Eptam for application rates, additional use directions, precautions and limitations before use.

Chemigation (Treflan H.F.P.-E.C. Only)

Treflan E.C. may be applied through properly equipped chemigation systems for weed control in potatoes. Refer to "Chemigation" section in the "General Information" section of the label for Treflan E.C. Do not apply Treflan E.C. through any type of irrigation system unless these directions are carefully followed.

Apply Treflan E.C. to potatoes in 0.5 to 1 acre inch of overhead sprinkler irrigation after planting, before emergence, or immediately following dragoff or after the potato plants have fully emerged. Existing weeds must be destroyed by tillage or cultivation prior to application of Treflan E.C.. Treflan does not control established weeds. Incorporation is not necessary when Treflan is applied by chemigation.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P.-E.C.
	(pints)
Coarse	1.0
Medium	1.5

• Do not apply by chemigation to fine textured soils.

Precautions: If cultivation is required after treatment with Treflan E.C., avoid completely covering potato plants with treated soil. Erratic weed control may result if cultivation exposes untreated soil between rows.

*Trademark of DowElanco

122-33-071 Approved 02/03/93
Initial printing.

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar, punctuation; or use of w/ vs. a.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

Use Rates for Fall Application Prior to Planting Cotton or Soybeans in the Spring

(For Use Only in Arkansas, Louisiana and Mississippi)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply Treflan® herbicide as a preplant soil incorporated treatment. Treflan may be applied in the fall prior to planting cotton or soybeans in the spring. For details on fall application, refer to instructions under "Application Timing" in the "General Information" section of the label for Treflan. Follow recommended soil preparation, application and incorporation procedures in the label for Treflan.

Broadcast application rates for fall application in cotton and soybean producing areas of Arkansas, Louisiana and Mississippi:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan TR-10
	(pints)	(pints)	(pounds)
Coarse	2.0	1.6	10.0
Medium	2.0 - 3.0	1.6 - 2.4	10.0 - 15.0
Fine	2.5 - 4.0	2.0 - 3.2	12.5 - 20.0

Note: Use the higher rate in the rate range under conditions of abundant rainfall and mild winter temperatures.

In U.S. cotton and soybean producing areas other than Arkansas, Louisiana and Mississippi, use application rates specified in the label for Treflan for fall application.

Restriction: In the season following this treatment, plant only those crops for which Treflan can be applied as a preplant incorporated treatment.

*Trademark of DowElanco

122-33-083

Approved: Treflan E.C., M.T.F. & 5 - 01/25/93

~~Treflan TR-10 - 03/24/93~~

Replaces: 122-22-076

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar, punctuation; or use of w/l case.

85 of 97

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.E.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan-5 (EPA Reg. No. 62719-118)~~

For Distribution and Use Only in the States of Alabama, Florida, Georgia, and Texas

Treflan Plus Atrazine Tank Mix for Weed Control in Field Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan Plus Atrazine Tank Mix

Treflan® herbicide may be applied in tank mix combination with atrazine herbicide plus an emulsifiable oil or oil concentrate when corn is from the 2-leaf stage of growth up to 12 inches tall and weeds are no more than 1 1/2 inches in height. A period of 24 to 48 hours is required to obtain atrazine postemergence activity after which the preemergence activity of the Treflan plus atrazine combination may be activated by 0.5 inch or more of rainfall or overhead sprinkler irrigation or mechanical incorporation. Note: In Texas, the tank mix of Treflan plus atrazine may be applied only to coarse textured soils.

Incorporation Directions: Mechanical incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at a speed that will provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.E.P., Treflan E.C. or Treflan M.T.F.	Treflan-5
	(pints)	(pints)
Coarse†	1.0††	0.8
Medium	1.0 - 1.5	0.8 - 1.2
Fine	1.5 - 2.0	1.2 - 1.6

†In Texas, the tank mix of Treflan plus atrazine may be applied only to coarse textured soils.

††Apply 1.5 pints/acre of Treflan H.E.P., E.C. or M.T.F. or 1.2 pints/acre of Treflan-5 on coarse soils to control fall panicum, pigweed and Texas panicum.

Precautions

- Do not apply to sweet corn or corn grown for seed.
- Do not apply Treflan to corn as a preplant or preemergence treatment or crop injury may occur.
- Where corn is planted in a furrow, apply Treflan only after a cultivation to move soil into the row.
- Refer to the atrazine product label for application rates, additional use directions, precautions and limitations before use.

*Trademark of DowElanco

122-33-084 Approved 06/22/93
Initial printing.

Revisions:

1. Minor editing to correct one or more may apply; reference to "Treflan" as a DowElanco trademark; grammar; punctuation; or use of w/ case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

(For Distribution and Use in the State of North Dakota)

Green Foxtail (Pigeongrass) Resistance to Dinitroaniline Herbicides Including Treflan, Identified in the State of North Dakota

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations stated below or imposed by the product labels for Treflan, except as described below.

Crop Production Precaution

Populations of green foxtail (pigeongrass) resistant to the dinitroaniline (DNA) class of herbicides have been identified in the state of North Dakota in fields which have a long history of dinitroaniline herbicide use. Treflan will not control green foxtail which has developed DNA resistance. Therefore, the grower assumes the risk of nonperformance due to DNA resistance if Treflan® herbicide is used to control green foxtail in the state of North Dakota. Alternative green foxtail control practices should be utilized in these fields.

DowElanco strongly recommends utilizing the following management practices to prevent or delay the development or spread of DNA-resistant green foxtail in spring cereal production areas:

1. Rotate herbicides so that the same product or same class of herbicide is not used repeatedly year after year. Treflan and/or other dinitroaniline herbicides should not be applied in consecutive years and preferably should be used only once in a three year period. Consult your local extension service or DowElanco representative for information regarding herbicides with alternative modes of action.
2. Rotate crops and use alternative weed control methods, including tillage, fallow periods and/or other herbicides with different modes of action.
3. Thoroughly clean small grains harvested from fields with confirmed resistance before using as seed, or avoid using grain from DNA-resistant fields for seed.
4. Thoroughly clean all crop residues from tillage and harvesting equipment before moving out of fields with confirmed resistance.

For further information or consultation, contact your DowElanco representative.

*Trademark of DowElanco

122-33-085 Approved 04/20/94
Replaces: 122-33-57

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco trademark;
grammar; punctuation; or use of w/l case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Special Use Program - Control of DNA-Resistant Goosegrass in Soybeans with Treflan Plus Dual Tank Mix

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply Treflan® herbicide plus Dual® herbicide tank-mix as a preplant incorporated treatment to control DNA-resistant goosegrass in soybeans. Follow application and incorporation directions provided in the label for Treflan.

Broadcast Application Rates/Acre

Soil Texture	Soils with 0 - 2% Organic Matter		
	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Dual 8E
	(pints)	(pints)	(pints)
Coarse	1.0	0.8	1.25
Medium	1.5	1.2	1.5
Fine	2.0	1.6	2

- Coarse and medium soils with 2-5% organic matter - 1.5 pints of Treflan H.F.P., E.C. or M.T.F. or 1.2 pints of Treflan 5
- Fine soils with 2-5% organic matter - 2.0 pints of Treflan H.F.P., E.C. or M.T.F. or 1.6 pints of Treflan 5

Note: Follow all applicable use directions, precautions and limitations in the product label for Dual herbicide.

*Trademark of DowElanco

®Dual is a registered trademark of Ciba-Geigy Corporation.

122-33-089

Approved: 04/15/93 (Treflan E.C.)

—10/06/93 (Treflan M.T.F.)

—10/14/94 (Treflan 5)

Initial printing.

Revisions:

1. Minor editing to correct (one or more may apply):
reference to "Treflan" as a DowElanco
trademark; grammar; punctuation; or use of u/
case.

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Special Use Program - Enhanced Control of Broadleaf Signalgrass In Soybeans with Treflan Plus Dual Tank Mix

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply Treflan® herbicide plus Dual® herbicide tank-mix as a preplant incorporated treatment for enhanced control of broadleaf signalgrass in soybeans. Follow application and incorporation directions provided in the label for Treflan.

Broadcast Application Rates/Acre

Soil Texture	Soils with 0 - 2% Organic Matter		
	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Dual 8E
	(pints)	(pints)	(pints)
Coarse	1.0	0.8	1.25 - 1.5
Medium	1.5	1.2	1.5 - 2
Fine	2.0	1.6	2 - 2.5

- Coarse and medium soils with 2-5% organic matter - 1.5 pints of Treflan H.F.P., E.C. or M.T.F. or 1.2 pints of ~~Treflan 5~~
- Fine soils with 2-5% organic matter - 2.0 pints of Treflan H.F.P., E.C. or M.T.F. or 1.6 pints of ~~Treflan 5~~

Note: Follow all applicable use directions, precautions and limitations in the product label for Dual herbicide.

*Trademark of DowElanco

®Dual is a registered trademark of Ciba-Geigy Corporation.

122-33-088

Approved: 05/18/93 (Treflan E.C.)

~~04/28/93 (Treflan M.T.F.)~~

~~10/14/94 (Treflan 5)~~

Initial printing.

Amendments:

- 1) Adds use of Treflan plus Dual tank mix for enhanced control of broadleaf signalgrass in soybeans.

Supplemental Labeling

89 of 97



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

Postemergence Soil Incorporated Application for Weed Control in Cotton

(For Distribution and Use in Texas, Oklahoma and New Mexico)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Treflan® herbicide may be applied to cotton as a postemergence incorporated treatment from the 4-true leaf stage up to layby. Apply as a broadcast spray using ground or aerial equipment. A ground applied directed spray is recommended if cotton foliage prevents uniform coverage of the soil surface.

Incorporation Directions: Treflan must be mechanically incorporated within 24 hours. Mechanical incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at a speed that will provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5
	(pints)	(pints)
Coarse	0.75 - 1.0	0.6 - 0.8
Medium	1.0 - 1.5	0.8 - 1.2
Fine	1.5 - 2.0	1.2 - 1.6

- Coarse and medium soils with 2-5% organic matter - 1.5 pints of Treflan ~~H.F.P., E.C. or M.T.F.~~ or 1.2 pints of Treflan 5
- Fine soils with 2-5% organic matter - 2.0 pints of Treflan ~~H.F.P., E.C. or M.T.F.~~ or 1.6 pints of Treflan 5

Precautions

- Do not apply within 90 days of harvest.
- Treated soil may be shallow cultivated or rotary hoed without loss of herbicidal activity. To avoid bringing untreated soil to the surface and loss of weed control, do not cultivate deeper than the depth of incorporation.

*Trademark of DowElanco

122-33-086 Approved 08/24/94
Initial printing.

Amendments:

- 1) Added broadcast treatment from 4-true leaf stage of growth to layby to directions for postemergence incorporated treatment.

90 of 97

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan* H.F.P. (EPA Reg. No. 62719-ELN)~~
~~Treflan E. C. (EPA Reg. No. 62719-93)~~
~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~
~~Treflan-5 (EPA Reg. No. 62719-118)~~

Fall Postplant Soil Incorporated Application for Suppression of Common Windgrass in Winter Wheat (For Distribution and Use Only in the State of Michigan)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan.

Directions For Use

Treflan* herbicide may be applied as a postplant soil incorporated treatment for suppression of common windgrass in winter wheat. Apply within 2 days after planting and incorporate one time within 24 hours. Use tillage equipment to destroy existing weeds prior to planting. Soil should be relatively free of plant residue, in good tilth and free from clods.

Broadcast Application Rates:

Soil Texture	Treflan H.F.P. Treflan E.C. or Treflan M.T.F.	Treflan-5
	(pVA)	(pVA)
Coarse	1.0	0.8
Medium	1.0 - 1.5	0.8 - 1.2
Fine	1.5 - 2.0	1.2 - 1.6

Incorporation Directions

Use a rotary hoe or other suitable equipment to incorporate Treflan into the upper one-inch of soil. Best results are obtained when the soil is not crusted. Operate the equipment at a speed necessary to achieve uniform soil mixing (rotary hoe, 8 to 10 mph). Set equipment so as to not disturb planted seed or mix the herbicide into the seed zone as crop injury may result if wheat seed is in direct contact with treated soil.

Planting Directions

Set planting equipment to place seed at least 1.5 inches deep.

Precautions: Under certain conditions, delayed crop emergence and or stand reduction may occur when Treflan is applied to wheat. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield. Carefully follow Special Precautions for Use of Treflan in Small Grains in main product label to minimize potential crop stress.

Note: Risk of crop injury from Treflan applied to winter wheat may be increased if Treflan, Senalant, or Prowl herbicides were applied during the current growing season to the preceding crop.

*Trademark of DowElanco

Amendment(s):

- 1) Added postplant application for windgrass control in winter wheat in the state of Michigan.

(Label Code) Approved: / /
Initial printing.

Supplemental Labeling

Supplemental Labeling for Proposed New Use

91 of 97



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan® H.F.P. (EPA Reg. No. 62719-ELN)

Treflan E.C. (EPA Reg. No. 62719-93)

Treflan M.T.F. (EPA Reg. No. 62719-116)

Treflan S (EPA Reg. No. 62719-118)

Treflan TR-10® (62719-134)

The Coulter Caddy as an Option for the Second Incorporation Pass for Treflan in Soybeans Grown Under Reduced or Conservation Tillage Conditions

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow use precautions and applicable use directions.
- Except as described below, use of Treflan according to this supplemental labeling is subject to all precautions and limitations imposed by the label affixed to the container for Treflan.

Directions For Use

Treflan® herbicide can be applied either in the fall or in the spring as a preplant incorporated treatment for weed control in soybeans grown under reduced or conservation tillage conditions. Make only one application per crop cycle.

Make applications to tilled land, standing or chopped stubble from the previous season's crop. The first incorporation of Treflan must occur within 24 hours. For the first incorporation, a tandem disc or combination tool that can thoroughly mix Treflan into the top 2 to 3 inches of the final seedbed while leaving the desired amount of plant residue on the soil surface is recommended. For fall or spring applications, the second incorporation can occur any time prior to planting or at planting with tillage equipment that provides uniform soil mixing used in conjunction with no-till planters.

The second incorporation can be accomplished with an aggressive Coulter Caddy system used in combination with the grain drill used for broadcast planting of soybeans. This method can be employed on soils that are in good tilth and have moderate soil moisture. For best results, the Coulter Caddy should be equipped and operated as follows:

1. Drills should be spaced no more than 8 inches apart.
2. Use fluted coulters that are 3/4 to 1 1/4 inches wide.
3. Set coulters to penetrate the soil to a depth of 1 1/2 to 2 inches.
4. Operate drill a minimum of 6 mph.
5. The drill should be followed by a First Harrow or Two-bar tine harrow to aid in leveling of the soil and crop residue.

Note: The Coulter Caddy system is not recommended for the second incorporation pass for spring applied Treflan TR-10® herbicide or dry bulk fertilizer impregnated with liquid formulations of Treflan.

*Trademark of DowElanco

(Label Code) Approved / /
Initial printing.

Amendment(s):

- 1) Adds incorporation options for reduced or conservation tillage in soybeans.

Supplemental Labeling



92 of 97

DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® E.C. (EPA Reg. No. 62719-ELN)~~
~~Treflan E.C. (EPA Reg. No. 62719-93)~~
~~Treflan M.T.F. (EPA Reg. No. 62713-116)~~
~~Treflan 5 (EPA Reg. No. 62719-118)~~
~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

Preplant Incorporated Treatment for Weed Control in Direct Seeded Chinese Cabbage or Kohlrabi

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

IMPORTANT: WAIVER OF LIABILITY REQUIRED FOR USE. Do not apply Treflan herbicide according to this supplemental labeling unless the crop owner has signed a Waiver of Liability Certificate to release DowElanco and seller from all claims and liability relating to crop damage or loss of yield resulting from the use of Treflan in Chinese cabbage or kohlrabi. The Waiver of Liability Certificate is attached to this label.

Directions for Use

Direct Seeded Chinese Cabbage or Kohlrabi: Apply Treflan® herbicide as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P. Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan TR-10
	(pints)	(pints)	(pounds)
Coarse	1.0	0.8	5.0
Medium	1.0	0.8	5.0
Fine	1.5	1.2	7.5

- All Soils with 2-5% organic matter - 1.5 pints of ~~Treflan H.F.P.~~, ~~Treflan E.C.~~ or ~~M.T.F.~~; 1.2 pints of ~~Treflan 5~~; or 7.5 pounds of ~~Treflan TR-10~~.

Precaution: Direct seeded cole crops exhibit marginal tolerance to Treflan. Stunting or reduced stands may occur if labeled rates are exceeded.

*Trademark of DowElanco

(Label Code) Approved / /
Initial printing.

Amendment(s):

1. Supplemental labeling to allow use of Treflan herbicides in Chinese cabbage or kohlrabi with signed waiver of liability.

93 of 97

WAIVER OF LIABILITY CERTIFICATE

I, the undersigned, acknowledge and understand that DowElanco (1) does not assume any liability for the use of Treflan® E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10® herbicides as a preplant incorporated treatment in Chinese cabbage or kohlrabi, and (2) makes no representations regarding its weed control or crop safety when applied in Chinese cabbage or kohlrabi.

I hereby release DowElanco and the Seller from whom I purchased Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides, their owners, officers, employees and agents, from any and all losses, damages, claims or causes of action, including reasonable attorney's fees and costs, relating to damage to or loss of any Chinese cabbage or kohlrabi crop resulting from use of Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides whether used by me or by a third party.

I understand that any application of Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides in Chinese cabbage or kohlrabi without a prior signed Waiver of Liability Certificate is an unlawful use of the product.

I warrant and represent that I have sole interest in the crop to be treated or am authorized to act on behalf of and bind all parties with an interest in such crop.

I have read this Waiver of Liability Certificate and have had the opportunity to have it reviewed by an attorney of my choosing.

Signature of Crop Owner: _____

Name of Crop Owner (Please Print): _____

Address of Crop Owner: _____

Seller send copy of signed Waiver of Liability Certificate to:

State Regulatory Affairs Manager
DowElanco
9330 Zionsville Road
Indianapolis, IN 46268-1054

*Trademark of DowElanco

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Treflan* H.F.P. (EPA Reg. No. 62719-ELN)

Treflan E.C. (EPA Reg. No. 62719-93)

Treflan M.T.F. (EPA Reg. No. 62719-116)

Treflan 5 (EPA Reg. No. 62719-118)

Treflan TR-10* (EPA Reg. No. 62719-131)

Weed Control in Eggplant

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

IMPORTANT: WAIVER OF LIABILITY REQUIRED FOR USE. Do not apply Treflan herbicide according to this supplemental labeling unless the crop owner has signed a Waiver of Liability Certificate to release DowElanco and seller from all claims and liability relating to crop damage or loss of yield resulting from the use of Treflan in eggplant. The Waiver of Liability Certificate is attached to this label.

Directions for Use

Transplant Eggplant

Apply and incorporate Treflan* herbicide before transplanting or apply post-transplant and incorporate. When applied post-transplant, direct liquid sprays to the soil between rows and beneath plants or broadcast Treflan TR-10* over the top.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan TR-10
	(pints)	(pints)	(pounds)
Coarse	1.0	0.8	5.0
Medium	1.0	0.8	5.0
Fine	1.5	1.2	7.5

- Coarse and medium soils with 2-5% organic matter - 1.5 pints of ~~Treflan H.F.P., Treflan E.C. or Treflan M.T.F.~~, 1.2 pints of ~~Treflan 5~~, or 7.5 pounds of ~~Treflan TR-10~~

*Trademark of DowElanco

(Label Code) Approved / /
Initial printing.

Amendment(s):

1. Supplemental labeling to allow use of Treflan herbicides in eggplant with signed waiver of liability.

WAIVER OF LIABILITY CERTIFICATE

I, the undersigned, acknowledge and understand that DowElanco (1) does not assume any liability for the use of Treflan H.F.P., Treflan* E.C., Treflan M.T.F., Treflan 5 or Treflan TR-10 herbicides as a preplant incorporated treatment in eggplant, and (2) makes no representations regarding its weed control or crop safety when applied in eggplant.

I hereby release DowElanco and the Seller from whom I purchased Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5 or Treflan TR-10 herbicides, their owners, officers, employees and agents, from any and all losses, damages, claims or causes of action, including reasonable attorney's fees and costs, relating to damage to or loss of any eggplant crop resulting from use of Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5 or Treflan TR-10 herbicides whether used by me or by a third party.

I understand that any application of Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5 or Treflan TR-10 herbicides in eggplant without a prior signed Waiver of Liability Certificate is an unlawful use of the product.

I warrant and represent that I have sole interest in the crop to be treated or am authorized to act on behalf of and bind all parties with an interest in such crop.

I have read this Waiver of Liability Certificate and have had the opportunity to have it reviewed by an attorney of my choosing.

Signature of Crop Owner: _____

Name of Crop Owner (Please Print): _____

Address of Crop Owner: _____

Seller send copy of signed Waiver of Liability Certificate to:

State Regulatory Affairs Manager
DowElanco
9330 Zionsville Road
Indianapolis, IN 46268-1054

*Trademark of DowElanco

Supplemental Labeling



DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

~~Treflan® H.F.P. (EPA Reg. No. 62719-ELN)~~

~~Treflan E.C. (EPA Reg. No. 62719-93)~~

~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~

~~Treflan 5 (EPA Reg. No. 62719-118)~~

~~Treflan TR-10® (EPA Reg. No. 62719-131)~~

Preplant Incorporated Treatment for Weed Control in Lentils (For Distribution and Use Only in the State of North Dakota)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

IMPORTANT: WAIVER OF LIABILITY REQUIRED FOR USE. Do not apply Treflan herbicide according to this supplemental labeling unless the crop owner has signed a Waiver of Liability Certificate to release DowElanco and seller from all claims and liability relating to crop damage or loss of yield resulting from the use of Treflan in lentils. The Waiver of Liability Certificate is attached to this label.

Directions for Use

Lentils

Apply and incorporate Treflan® herbicide in the spring before planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan H.F.P., Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan TR-10
	(pints)	(pints)	(pounds)
Coarse	1.0	0.8	5.0
Medium	1.0	0.8	5.0
Fine	1.5	1.2	7.5

*Trademark of DowElanco

(Label Code) Approved / /
Initial printing.

Amendment(s):

1. Supplemental labeling to allow use of Treflan herbicides in lentils with signed waiver of liability

WAIVER OF LIABILITY CERTIFICATE

I, the undersigned, acknowledge and understand that DowElanco (1) does not assume any liability for the use of Treflan H.F.P., Treflan* E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10* herbicides as a preplant incorporated treatment in lentils, and (2) makes no representations regarding its weed control or crop safety when applied in lentils.

I hereby release DowElanco and the Seller from whom I purchased Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides, their owners, officers, employees and agents, from any and all losses, damages, claims or causes of action, including reasonable attorney's fees and costs, relating to damage to or loss of any lentils crop resulting from use of Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides whether used by me or by a third party.

I understand that any application of Treflan H.F.P., Treflan E.C., Treflan M.T.F., Treflan 5, or Treflan TR-10 herbicides in lentils without a prior signed Waiver of Liability Certificate is an unlawful use of the product.

I warrant and represent that I have sole interest in the crop to be treated or am authorized to act on behalf of and bind all parties with an interest in such crop.

I have read this Waiver of Liability Certificate and have had the opportunity to have it reviewed by an attorney of my choosing.

Signature of Crop Owner: _____

Name of Crop Owner (Please Print): _____

Address of Crop Owner: _____

Seller send copy of signed Waiver of Liability Certificate to:

State Regulatory Affairs Manager
DowElanco
9330 Zionsville Road
Indianapolis, IN 46268-1054

*Trademark of DowElanco