

62719-255

11/06/2000

1 acet copy 1/18

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

November 6, 2000

Dr. John Jachetta  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268-1054

Subject: Dursban 50W Nursery  
EPA File No. 62719-255  
Resubmission dated Nov 1, 2000

Dear Dr. Jachetta:

The revised product labeling referred to above, submitted in connection with the Chlorpyrifos Memorandum of Agreement (MOA) as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended label:

1. On pages 1 and 3 delete the sentence: "To be applied by or under the direct supervision of trained applicators responsible for insect control programs." The product is now a Restricted Use Pesticide and this statement is redundant.

The conditions of paragraph 5 of the June 2000 Memorandum of Agreement apply to this label amendment, specifically this amended label is effective December 1, 2000. Failure to comply with any of the conditions of registration set forth in Paragraph 10 of the MOA shall be grounds for cancellation of the affected registration(s) under FIFRA section 6(e). Submit one copy of the revised final printed label before releasing the product for shipment. If the conditions enumerated above are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e).

A stamped copy of the label is enclosed for your records. If you have questions, please contact Dennis McNeilly at (703) 308-6742 or electronically at [McNeilly.Dennis@EPA.gov](mailto:McNeilly.Dennis@EPA.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Dennis McNeilly", written over a horizontal line.

Dennis McNeilly, Chemist  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

2/118

(Base label):

**RESTRICTED USE PESTICIDE**  
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(Logo) Dow AgroSciences LLC

# Dursban\* 50W Nursery

## In Water Soluble Packets

**To be applied only by or under the direct supervision of trained applicators responsible for insect control programs.**

Active Ingredient:

chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl)phosphorothioate .....	50%
Inert Ingredients .....	50%
Total .....	100%

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
NOV 6 2000

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

62719-255

**Keep Out of Reach of Children**

# DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Corrosive • Causes Irreversible Eye Damage • May Be Fatal If Swallowed • Harmful If Inhaled Or Absorbed Through Skin**

**Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust or spray mist. Keep away from food, feedstuffs, and water supplies.**

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

#### Organophosphate

**If in eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**If swallowed:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**If on skin:** Wash with plenty of soap and water. Get medical attention.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**Note to physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

### Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to label booklet for Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-255

EPA Est. 00000-XX-00

\*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

## Specialty Insecticide

Net Weight \_\_ lb

4/18

(Datapack cover):

**RESTRICTED USE PESTICIDE**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(Logo) Dow AgroSciences LLC

**Dursban\* 50W Nursery**

**In Water Soluble Packets**

**To be applied only by or under the direct supervision of trained applicators responsible for insect control programs.**

Active Ingredient:

chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl)phosphorothioate..... 50%

Inert Ingredients ..... 50%

Total ..... 100%

**Keep Out of Reach of Children**

**DANGER PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-255

EPA Est. 00000-XX-00

\*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

**Specialty Insecticide**

**Net Weight \_\_ lb**

5/18

(Page 1 through end):

<b>Table of Contents</b>	<b>Page</b>
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Protective Personal Equipment (PPE)	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Mixing Directions	-
Approved Uses	-
Ornamentals Grown in Nurseries	-
Ornamentals Grown in Nurseries (Dormant Spray of Tree Pests)	-
Ornamentals (Pre-Plant Incorporation Treatment of Field Grown Nursery Stock)	-
Fruit, Nut, and Citrus Trees Grown in Nurseries	-
Agricultural Sod Farm Turf and Turf Grown for Seed	-
Turf In Golf Courses, Road Medians and Industrial Plant Sites	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## DANGER

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Corrosive • Causes Irreversible Eye Damage • May Be Fatal If Swallowed • Harmful If Inhaled Or Absorbed Through Skin**

**Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust or spray mist. Keep away from food, feedstuffs, and water supplies.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

#### Organophosphate

**If in eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**If swallowed:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**If on skin:** Wash with plenty of soap and water. Get medical attention.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**Note to physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

### Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**Do not apply this product through any type of irrigation system.**  
**Do not formulate this product into other end-use products.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Chemical-resistant headgear for overhead exposure

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Adults, children, and pets should not contact treated surfaces until the spray has dried.

### **Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

**Pesticide Disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** When all packets are used, dispose of empty package in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### **General Information**

Dursban\* 50W Nursery In Water Soluble Packets insecticide is a wettable powder formulation designed for use as a spray to control various pests injurious to trees, turf, and ornamental plants on the following sites: ornamentals and trees in nurseries, sod farm turf, turf grown for seed, and turf on golf courses, road medians and around industrial plant sites. The pests controlled are listed in the accompanying tables. Dursban 50W Nursery is compatible with insecticides, miticides, and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

**Attention:** Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock or use hay for livestock bedding. Do not use in poultry houses.

#### **Handling Precautions for Water Soluble Packets**

Do not allow packets to become wet prior to adding to the spray tank. Wear gloves when handling water-soluble packets. Do not handle with wet gloves. Do not handle water-soluble packets excessively since this may cause breakage. Reseal outer package to protect remaining packets. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

#### **Mixing Directions**

For paddle, sparger, and bypass agitation systems.

1. Fill the spray tank 1/3 full of water.

**Note:** For paddle agitation systems, paddles should be covered with water.

2. Activate agitation system.
3. Add required number of water-soluble packets to obtain desired dilution.
4. Complete tank filling while the packet(s) dissolve and Dursban 50W Nursery disperses.
5. Before beginning to spray, make sure water-soluble packet(s) have dissolved and Dursban 50W Nursery has completely dispersed.

**Note:** Depending on the water temperature and amount of agitation, the packets and Dursban 50W Nursery should be completely dispersed within approximately 5 minutes after adding to water.

### **Approved Uses**

Use of this product is limited to pest control only on the following sites: ornamentals and trees in nurseries, sod farm turf, turf grown for seed, and turf on golf courses, road medians and around industrial plant sites.



9/18

## Ornamentals Grown in Nurseries

Use Dursban 50W Nursery insecticide to treat flowers, shrubs, evergreens, vines, shade and flowering trees, and non-bearing fruit trees found to be infested with the types of pests listed in the following tables including: (1) Sucking Insects and Mites; (2) Defoliators and Leafminers; (3) Borers, Bark Beetles, and Weevils; and (4) Ants, Termites, and Miscellaneous Pests.

### General Use Directions

Dilute Dursban 50W Nursery insecticide with water according to directions given in the table and apply using suitable hand or power-operated spray equipment in a manner to provide complete and uniform coverage. Attempt to penetrate dense foliage, but avoid over spraying to the point of excessive runoff. When using spray equipment delivering less than 200 gallons of finished spray per acre, use the rate recommended in the per acre column. Uniform coverage is critical for effective insect control.

**Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions following each table.

Consult your State Agricultural Experiment Station or Extension Service specialist for application timing and other specific use information

**Note: Phytotoxicity:** Environmental factors and varietal differences have significant effects on phytotoxic expression. Dursban 50W Nursery insecticide has been tested on numerous ornamental plants without causing serious phytotoxicity at recommended use rates. Before treating large numbers of plants, it is recommended that a small block of plants be sprayed and observed for 7 to 10 days to determine phytotoxic potential.

### (1) Sucking Insects and Mites

Pest	Amount of Dursban 50W Nursery per	
	100 gallons	Acre
Adelgids (such as: Cooley and Eastern spruce galls, Pine bark) Aphids (Such as: Apple, Balsam twig, Black pecan, Chrysanthemum, Cottonwood, Grape myrtle, Elm leaf, Melon, Peach, Rose, Spirea, White pine, Woolly apple, Yellow pecan) Boxelder bugs Lace bugs (such as: Hawthorn) Leafhoppers Periodical cicada Plant bugs Psyllids Spittlebugs Thornbug Whiteflies	0.5 - 1 lb (two to four 4 oz packets)	1 - 2 lb (four to eight 4 oz packets)
Leafhoppers Mealybugs (Such as: Citrus, Taxus) Mites (Such as: Clover, Red spider, Southern red) (1) Thrips (exposed)	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Scale insects (2) (Such as: Cottonycushion, Cottony maple, Dearness, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oystershell, Pine needle, San Jose, Tea, White birch)	2 lb (eight 4 oz packets)	2 - 4 lb (eight to sixteen 4 oz packets)

**Pest-Specific Directions**

1. For effective control of **spider mites** when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly hatched nymphs.
2. Time applications for control of **scale insects** when crawlers or first two stages of settled scales are present.

**(2) Defoliators and Leafminers**

Pest	Amount of Dursban 50W Nursery per		
	100 gallons	Acre	
Armyworms (Such as: Fall, Yellowstriped) Bagworms (1) Cankerworms Catalpa sphinx Elm spanworms Fall webworms Grasshoppers Greenstriped mapleworms Green fruitworms Hornworms Jackpine budworms Juniper webworms Katydids Leafrollers (2) Maple leafcutters (3) Oak skeletonizers Oleander caterpillars	Orange tortrix Poplar tentmaker Puss caterpillars Rose chafers Sawflies, exposed (Such as: European pine, Pine, Pin oak, Redheaded) Spring elm caterpillars Springtails Spruce budworms Tent caterpillars (Such as: Eastern, Forest, Western) Walnut caterpillars Western spruce budworms Yellownecked caterpillars	0.5 - 1 lb (two to four 4 oz packets)	1 - 2 lb (four to eight 4 oz packets)
Balsam gall midge Beet armyworms Beetles (Such as: Fuller rose, Native elm bark) (5) Browntail moth Cutworms Cypress tip moth Douglas fir tussock moth European pine shoot moth Gypsy moth (4) Holly bud moth Mahogany webworms	Mimosa webworms Nantucket pine tip moth Oakworms (such as: California, Orangestriped, Redhumped) Pandora moth Pitch pine moth Redhumped caterpillars Subtropical pine tip moth Tussock moth	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Beetles (Such as: Cottonwood leaf, Elm leaf, Flea, Willow leaf) (6)		1 - 2 lb (four to eight 4 oz packets)	2 lb (eight 4 oz packets)
Leafminers Needleminers (Such as: Jeffrey pine, Lodgepole pine, Spruce) Pine needle midge		2 lb (eight 4 oz packets)	2 - 4 lb (eight to sixteen 4 oz packets)

**Pest-Specific Directions**

1. For **bagworms**, treat when larvae are small and actively feeding.
2. For effective control of **leafrollers**, spray should be applied before leaves are tightly rolled.
3. For **maple leafcutter** on maple trees, apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production.
4. To control migrating and invading **gypsy moth larvae**, treat trunks and foliage.

5. To reduce twig and branch feeding by bark beetles, applications should be made in the spring or early summer.
6. For cottonwood leaf beetles, use Dursban 50W Nursery in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present.

### (3) Borers, Bark Beetles, and Weevils

Pest	Amount of Dursban 50W Nursery per	
	100 gallons	Acre
Weevils (such as: Blackvine (1), Cranberry girdler (2), Yellow poplar, Pine reproduction)	1 lb (four 4 oz packets)	2 - 4 lb (eight to sixteen 4 oz packets)
Borers (3): Clearwing moths (Such as: Ash, Dogwood, Lesser peachtree, Lilac, Oak, Peachtree, Rhododendron), Metallic wood (Such as: Bronze birch, Flatheaded appletree, Twolined chestnut), Longhorned beetles (Such as: Cottonwood, Locust, Red oak) Pales weevil adults Zimmerman pine moth	2 lb (eight 4 oz packets)	-
Pales weevil (4) Northern pine weevil (4)	6 lb (twenty-four 4 oz packets)	-
Beetles (Such as: Ambrosia, Anobiidae, Black turpentine, Blister, Cottonwood leaf, Elm leaf, European elm bark, Fuller rose, Japanese, June, Mountain pine, Native elm bark, Southern pine, Spruce, Western pine, Willow leaf) (5)	16 lb (sixty-four 4 oz packets)	-
Weevils (Such as: Northern pine, Pitch eating, Twig) (6)	32 lb (one hundred twenty-eight 4 oz packets)	-

### Pest-Specific Directions

1. **Blackvine weevils** are night feeders. Late afternoon spraying will give control in some areas.
2. For **cranberry girdler larvae** infesting Douglas fir seedlings, direct spray at lower crown and stems following egg laying during summer. Apply a minimum of 50 gallons of dilute spray per acre and irrigate immediately after application to move the insecticide into the top 1 to 2 inches of soil.
3. For **borers**, spray trunks and lower limbs of trees and shrubs when adults begin to emerge. For **peachtree borers**, spray flowering trees and shrubs of the genus *Prunus* as a trunk spray before newly-hatched larvae enter the trees and thoroughly wet all bark areas from ground level to scaffold limbs. Pheromone traps may aid in detection of adult clearwing moths. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat.
4. For control of **pales and northern pine weevil larvae**, apply as a cut stump spray or drench.
5. For **preventive treatment**, spray the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For **remedial treatment**, spray the main trunk of infested trees or logs when damage occurs, but before adult beetles begin to emerge.
6. For pine seedlings, treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff. Do not use more than 6 gallons of spray dilution per acre.

**(4) Ants, Termites, and Miscellaneous Pests**

Pest	Amount of Dursban 50W Nursery per	
	100 gallons	Acre
Ants Cockroaches (Such as: American, Asian wood, Brownbanded, German, Oriental, Smokey brown) Fire ants (foraging workers) Fire ants (mounds) (1) Sowbugs Springtails	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Carpenter ants (2) Termites	16 lb (sixty-four 4 oz packets)	-

**Pest-Specific Directions**

1. For individual **fire ant mounds** apply Dursban 50W Nursery insecticide as a drench. Dilute 0.75 lb of Dursban 50W Nursery (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
2. If possible, locate carpenter ant nests and drench thoroughly.

**Ornamentals Grown in Nurseries (Dormant Spray of Tree Pests)**

Use Dursban 50W Nursery insecticide as a dormant or delayed dormant spray at the rates indicated to control the listed insects.

Pest	Amount of Dursban 50W Nursery per 100 gallons <sup>1</sup>
Aphids (Such as: Mealy plum, Rosy apple, Woolly apple) Borers (such as: Peach twig) Cutworms (Such as: Climbing) Leafrollers (Such as: Oblique banded, Pandemis) Pear psylla adults Scale insects (Such as: San Jose)	0.5 - 1 lb (two to four 4 oz packets)

<sup>1</sup>Based on 200 to 600 gallons per acre as a dilute spray.

**General Use Directions**

While Dursban 50W Nursery insecticide may be used without oil, oil is recommended to control additional pests such as European red mite.

For high volume (dilute) sprays, tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water. Spray the entire tree to runoff using suitable ground spray equipment.

For low volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of Dursban 50W Nursery insecticide as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

**Precautions:** Because cold, dry conditions may cause Dursban 50W Nursery insecticide plus oil to infuse trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated. Do not use more than 4 lb (sixteen 4 oz packets) of Dursban 50W Nursery insecticide per acre.

**Restrictions:** Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated areas.

### **Ornamentals (Pre-Plant Incorporation Treatment of Field Grown Nursery Stock)**

Apply Dursban 50W Nursery insecticide to soil and incorporate before transplanting or planting to control white grubs during transplant or seedling establishment. Apply to the soil surface as a broadcast spray at a rate of 8 pounds per acre (thirty-two 4 oz packets) using sufficient water to obtain adequate coverage. Do not aerially apply. On the same day of the treatment, incorporate the insecticide into the top 2 to 4 inches of the soil using a tandem disc, field cultivator or equivalent incorporation equipment capable of thorough soil mixing.

### **Fruit, Nut, and Citrus Trees Grown in Nurseries**

Use Dursban 50W Nursery insecticide to treat (**non-bearing**) apple, almond, cherry, filbert, grapefruit, lemon, nectarine, pecan, peach, plum, prune, and walnut trees or tree fruit not grown for commercial consumption which are infested with pests listed in the following table.

Pest	Amount of Dursban 50W Nursery per		
	100 gallons	Acre	
Aphids (Such as: Apple, Black cherry, Black pecan, Filbert, Rosey apple, Woolly apple, Yellow pecan) Apple Maggot Borers (Such as: American plum, Dogwood, Lesser peachtree, Pacific flathead, Peach twig, Peachtree, Shothole) Cherry fruit fly Climbing cutworm Codling Moth European apple sawfly Eyespotted bud moth European corn borer Fall webworm Filbert worm Grasshoppers (1) Green fruitworm Hickory shuckworm Katydid Leaf rollers and tiers (Such as: Avacado leafroller, Filbert leafroller, Fruit tree leafroller, Oblique banded leafroller, Omnivorous leaftier, Orange tortrix, Pandemis leafroller, Redbanded leafroller, Variegated leafroller)	Lesser appleworm Mineola Moth Mites (2) (such as: European red, Pecan leaf scorch, Two spotted spider) Navel orangeworm Oriental fruit moth Pecan nut casebearer Pecan weevil Periodical cicada <i>Phylloxera</i> spp. Plum curculio Potato leafhopper Rose chafer Scale insects (such as: San Jose, Walnut, European fruit lecanium) Spittlebug Spotted tentiform leafminer Stinkbug Tarnished plant bug Tufted apple budmoth Walnut husk fly Western tussock moth White apple leafhopper Winter moth	2 lb (eight 4 oz packets)	2 - 4 lb (eight to sixteen 4 oz packets)
Scale insects (3) (Such as: Black scale, Brown soft scale, California red scale, Florida red scale, Snow scale, Chaff scale, Purple scale, Long scale) Mealybugs Thrips		4 lb (sixteen 4 oz packets)	4 - 8 lb (sixteen to thirty- two 4 oz packets)

### General Use Directions

A petroleum spray oil recommended for use on **citrus trees** may be added to spray mixtures only at rates of up to 1.8 gallons per 100 gallons of water to improve control of aphids, mealybugs, scale insects, and thrips. Treat when insects become a problem or in accordance with the local spray schedule recommended by your State Extension Service Specialist. **Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions.

### Precautions:

Do not apply as a foliar spray to **sweet cherries**. Contact of spray solution with leaves may result in premature leaf drop. Observe local use directions for tank mix combinations especially in regard to applications of Dursban 50W Nursery plus spray oil. Do not apply Dursban 50W Nursery insecticide in combination with oil to walnuts. Dry or cold conditions may cause Dursban 50W Nursery insecticide plus oil to infuse trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated. Do not apply when trees are stressed by drought or if temperature exceeds 95°F.

**Pest-Specific Directions**

1. **Lubber grasshoppers** must be controlled when they are small (less than 1 inch in length) by direct contact with spray.
2. For effective control of **spider mites** when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly hatched nymphs.
3. For control of Black scale, Brown soft scale, California red scale, and Chaff scale in Arizona, California and Texas, apply 8 lb (thirty-two 4 oz packets) of Dursban 50W Nursery per acre (8 lb/100 gal).

**Agricultural Sod Farm Turf and Turf Grown for Seed**

Use Dursban 50W Nursery insecticide to control the pests listed in the following table by application at the recommended dosages.

<b>Pest</b>	<b>Amount of Dursban 50W Nursery per Acre</b>
Ants Armyworm (such as: Beet, Fall, Yellowstriped) Chiggers (1) Chinch bugs Crickets Cutworms Deer ticks (2) Earwigs European crane fly larvae Fiery skipper Fire ants (foraging workers) Fire ants (mounds) (3) Fleas Grasshoppers Greenbugs Green June beetle grubs Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Bermudagrass stunt, Winter grain) Mosquitoes (4) Pillbugs Sod webworms (lawn moths) (5) Sowbugs Springtails Ticks (1)	2 lb (eight 4 oz packets)
Billbugs (Such as: Bluegrass, Denver, Hunting) (6)	2 - 4 lb (eight to sixteen 4 oz packets)
Black turfgrass ataenius (7)	4 lb
Annual bluegrass weevil ( <i>Hyperodes</i> ) (8)	(sixteen 4 oz packets)
Mole Crickets (9)	4 - 6 lb (sixteen to twenty four 4 oz packets)
White grubs (Such as: Black turfgrass ataenius, European chafer, Southern masked chafer and Japanese beetle larvae) (10)	4 - 8 lb (sixteen to thirty-two 4 oz packets)
Imported fire ants in commercial sod (11)	16 lb (sixty-four 4 oz packets)

**General Use Directions**

Dilute Dursban 50W Nursery insecticide in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment. **Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions.

**Pest-Specific Directions**

1. Use Dursban 50W Nursery insecticide for area control of **ticks** and **chiggers** infesting sod farm turf and turf grown for seed. Apply Dursban 50W Nursery insecticide in water at the rate of 0.25 lb per acre (one 4 oz packet) using a hydraulic sprayer, mist applicator, backpack sprayer, or other suitable hand or power operated spray equipment.
2. For control of **deer ticks** apply Dursban 50W Nursery insecticide in water at the rate of 2 lb per acre (eight 4 oz packets). Treat turf, grassy areas, weeds, ground surface and debris, using enough spray volume to obtain thorough coverage.
3. **For professional use only.** For individual **fire ant mounds** apply Dursban 50W Nursery insecticide as a drench. Dilute 0.75 lb Dursban 50W Nursery (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
4. Mosquitoes coming to rest on areas treated for control of turf pests will be controlled for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.
5. For **sod webworms**, delay watering or mowing of the treated area for 12 to 24 hours after treatment.
6. For **billbugs**, spray early in the season just prior to, or coinciding with first appearance of adults as recommended by your local Agricultural Extension Service Specialist.
7. For **black turfgrass ataenius** adults, spray early in the season as recommended by your local Agricultural Extension Service Specialist. A repeat application may be needed 1 to 2 weeks later.
8. To control **annual bluegrass weevil**, spray suspected problem areas in mid-April and again in Mid-May, or as recommended by your local Agricultural Extension Service specialist.
9. To control **mole crickets** in turfgrass, apply Dursban 50W Nursery insecticide in a minimum spray volume of 50 gallons of water per acre using broadcast or suitable hand held application equipment. Apply when early stage nymphs are active. Efficacy may be enhanced by spraying late in the afternoon or early evening and irrigating the turf prior to treatment to move mole crickets to the soil surface.
10. For **white grubs**, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. For best results, soil should be moist prior to treatment. **For best results, immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.**
11. **For professional use only.** Imported **fire ants**: Apply Dursban 50W Nursery as a spray to the area of sod to be cut. Immediately after application, irrigate treated area with at least 1/2 acre-inch of water or a sufficient volume to thoroughly soak below the cut line. Do not apply a volume or rate of water to cause runoff. Sod can be cut when **vegetation** has dried and when soil is moist but not wet.

**Precaution:** Personnel handling treated sod should wear elbow-length chemically resistant gloves and apron. Protective clothing and boots should be worn to prevent repeated skin exposure to treated sod.

**Certification Period:** Forty-two days from day of application.



### Turf In Golf Courses, Road Medians and Industrial Plant Sites (Maximum Use Rate of 2 lb (1 lb a.i.)/Acre)

Use Dursban 50W Nursery insecticide to control the pests listed in the following table by application at the recommended dosages.

Pest	Amount of Dursban 50W Nursery per Acre
Ants Armyworm (such as: Beet, Fall, Yellowstriped) Chiggers (1) Chinch bugs Crickets Cutworms Deer ticks (2) Earwigs European crane fly larvae Fiery skipper Fire ants (foraging workers) Fire ants (mounds) (3) Fleas	Grasshoppers Greenbugs Green June beetle grubs Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Bermudagrass stunt, Winter grain) Mosquitoes (4) Pillbugs Sod webworms (lawn moths) (5) Sowbugs Springtails Ticks (1)

#### General Use Directions

Dilute Dursban 50W Nursery insecticide in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment. **Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions.

#### Pest-Specific Directions

1. Use Dursban 50W Nursery insecticide on golf course turf, road medians and around industrial plant sites. Do not allow public use of treated areas during application or until spray has dried. Apply Dursban 50W Nursery insecticide in water at the rate of 0.25 lb per acre (one 4 oz packet) using a hydraulic sprayer, mist applicator, backpack sprayer, or other suitable hand or power operated spray equipment.
2. For control of deer ticks apply Dursban 50W Nursery insecticide in water at the rate of 2 lb per acre (eight 4 oz packets). Treat low underbrush, turf, grassy areas, weeds, ground surface and debris, using enough spray volume to obtain thorough coverage.
3. **For professional use only.** For individual fire ant mounds apply Dursban 50W Nursery insecticide as a drench. Dilute 0.75 lb Dursban 50W Nursery (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
4. Mosquitoes coming to rest on areas treated for control of turf pests will be controlled for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.
5. For sod webworms, delay watering or mowing of the treated area for 12 to 24 hours after treatment.

18/18

---

### **Warranty Disclaimer**

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

\*Trademark of Dow AgroSciences LLC  
EPA-Accepted \_\_\_/\_\_\_/\_\_\_