62719-259

12-25-2011



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

> OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Niamh McMahon Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268

DEC 23 2011

Subject: Supplemental Label for post emergence broadleaf weed control in cottonwood, poplar and eucalyptus tree plantations.

Transline EPA Reg. No: 62719-259 Application Dated: September 22, 2011

Dear Ms. McMahon

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

The supplemental labeling must be incorporated into the main product labeling within eighteen (18) months of the date of this letter.

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Grant Rowland at rowland.grant@epa.gov or 703-347-0254.

Sincerely,

Kathryn Montague Product Manager 23 Herbicide Branch Registration Division (7505P)

Supplemental Labeling

Under the Federal Insecticide, Forstolde, and Recenticide Act, as aduanded, for the pessicide stered ander Dow AgroSciences 62719-29

ACCEPTED

NFC 23 2011

Dow AgroSciences LLC

9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Transline[®]

EPA Reg. No. 62719-259

Postemergence Broadleaf Weed Control in Cottonwood/Poplar and Eucalyptus **Tree Plantations**

(Not for Use or Distribution Within the State of Florida)

This supplemental label expires on December 31, 2014, and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application. Read the label affixed to the container for Transline[®] herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Transline according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Transline.

Directions for Use

Refer to product label for Precautions and Restrictions, Mixing and Application directions.

Use Transline for postemergence control of labeled broadleaf weeds in new and established plantings of cottonwood/poplar and eucalyptus tree plantations. Apply as a broadcast foliar spray over trees, as a banded or directed spray, or as a spot application. Broadcast apply in 10 gallons or more per acre total spray volume using ground or aerial equipment. Apply either during or after site preparation or after trees are planted. Use sufficient spray volume to ensure thorough coverage of weeds. Apply to new plantings only after they are well established as indicated by several inches of new healthy growth. Sequential applications may be made as long as the total rate per annual use season does not exceed 1 1/3 pints per acre.

If applying as a spot treatment, apply to weeds on a spray to wet basis (not to runoff) with uniform and complete spray coverage. Avoid contact with foliage of cottonwood/poplar trees as much as possible.

Hand-Held Sprayers: Mix the amount of Transline corresponding to the desired broadcast rate in 1 gallonor more of water and apply to an area of 1000 sq ft. For larger areas, multiply the values in the table below by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3500 sq ft, multiply the table value by 3.5 (3500 ÷ 1000 = 3.5). Prepare a spray solution by adding 1/4 fl oz of Transline per gallon of water. When applied at 1 gallon of spray per 1000 sg ft, this spray concentration is equivalent to a broadcast rate of 2/3 pint per acre. Using the example of 3500 sq ft, the applicator would use $3.5 \times 1/4$ fl oz = 7/8 fl oz of Transline in 3.5 gallons of water.

Amount of Transline to Treat an Area of 1000 sq ft (pint/acre) (Mix in 1 Gallon or More of Spray)				
1/3	2/3	1	1 1/3	
1/8 fl oz	1/4 fl oz	3/8 fl oz	1/2 fl oz	
(3.6 mL)	(7.3 mL)	(11 mL)	(15 mL)	

Tank Mix: This product may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product. Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.

Use Site	Weed Species	Application Rate (pint/acre)	Application Timing
cottonwood/poplar plantations	ageratum (Ageratum conyzoides) fireweed (Chamerion angustifolium) thickhead (Crassocephalum crepidioides)	1/3 – 2/3	Smaller plants may be controlled using the lower rate. Plants may be slow to show control at lower rates.
eucalyptus tree plantations	beggarweed (<i>Desmodium</i> <i>intortum</i>) horseweed (<i>Conyza canadensis</i>)	1/2 – 1 1/3	Smaller plants may be controlled using the lower rate. Use 1 to 1 1/3 pints per acre for <i>Desmodium intortum</i> in flower stage.

Specific Use Precautions and Restrictions:

- Do not tank mix Transline with other herbicides labeled for use in cottonwood/poplar tree plantations unless spray avoids all contact with tree foliage.
- A tank mix of Transline with another herbicide may be foliarly applied over eucalyptus trees provided that foliar application is specifically allowed by the label of the tank mix partner.
- Certain broadleaf weeds, such as mustards, henbit, chickweed, kochia, lambsquarters, pigweed, Russian thistle and bindweed, will not be controlled or suppressed.

[®]Trademark of Dow AgroSciences LLC R113-010 EPA accepted: __/_/__ Initial printing