

62719-259

6/26/2012

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington D C 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Niamh McMahon
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268

JUN 26 2012

Subject Supplemental Label For Postemergence Broadleaf Weed
Control in Cottonwood/Poplar and Eucalyptus Tree Plantations
Transline®
EPA Reg No 62719 259
Application Dated March 27 2012

Dear Ms McMahon

The supplemental labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act as amended is acceptable

The supplemental labeling must be incorporated into the main product labeling within eighteen (18) months of the date of this letter

A stamped copy of your label is enclosed for your records You must submit one (1) copy of the final printed label before you release the product for shipment Products shipped after eighteen (18) months from the date of this letter must bear the new revised label If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA §6(e) Your release for shipment of the product constitutes acceptance of these conditions

If you have any questions please contact Grant Rowland at rowland_grant@epa.gov or 703 347 0254

Sincerely

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2/3

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Transline® EPA Reg No 62719 259

Postemergence Broadleaf Weed Control in Cottonwood/Poplar and Eucalyptus Tree Plantations (Not for Use or Distribution Within the State of Florida)

This supplemental label expires on December 26 2013 and must not be used or distributed after this date

ATTENTION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
This labeling must be in the possession of the user at the time of application
Read the label affixed to the container for Transline® herbicide before applying Carefully follow all precautionary statements and applicable use directions
Use of Transline according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Transline

Directions for Use

Refer to product label for Precautions and Restrictions Mixing and Application directions

Use Transline for postemergence control of labeled broadleaf weeds in new and established plantings of cottonwood/poplar and eucalyptus tree plantations Apply as a broadcast foliar spray over trees as a banded or directed spray or as a spot application Broadcast apply in 10 gallons or more per acre total spray volume using ground or aerial equipment Apply either during or after site preparation or after trees are planted Use sufficient spray volume to ensure thorough coverage of weeds Apply to new plantings only after they are established as indicated by several inches of new healthy growth

Application Rates Apply at a rate of 1/3 to 1 1/3 pints/acre in eucalyptus tree plantations and 1/3 to 2/3 pints/ acre in cottonwood/poplar plantations Sequential applications may be made as long as the total rate per annual use season does not exceed 1 1/3 pints per acre

In Hawaii For control of additional weeds in eucalyptus tree plantations (listed below) apply as a broadcast spray over the top as a directed or branded spray or as a spot treatment

Additional Weeds Controlled in Eucalyptus Tree Plantations in Hawaii

Weed Species	Application Rate (pint/acre)	Application Timing
ageratum (<i>Ageratum conyzoides</i>) fireweed (<i>Erechtites hieracifolia</i>) thickhead (<i>Crassocephalum crepidioides</i>)	1/3 – 2/3	Smaller plants may be controlled using the lower rate Plants may be slow to show control at lower rates
beggarweed (<i>Desmodium intortum</i>) horseweed (<i>Conyza bonariensis</i>)	2/3 – 1 1/3	Smaller plants may be controlled using the lower rate Use 1 to 1 1/3 pints per acre for <i>Desmodium intortum</i> in flower stage

JUN 26 2012
62719-259

Spot Application If applying as a spot treatment apply to weeds on a spray to wet basis (not to runoff) with uniform and complete spray coverage Avoid contact with foliage of cottonwood/poplar trees as much as possible See use instructions for Hand Held Sprayers

Hand Held Sprayers Mix the amount of Transline corresponding to the desired broadcast rate in 1 gallon or more of water and apply to an area of 1000 sq ft For larger areas multiply the values in the table below by the area to be treated in thousands of square feet e g if the area to be treated is 3500 sq ft multiply the table value by 3.5 (3500 - 1000 = 3.5) Prepare a spray solution by adding 1/4 fl oz of Transline per gallon of water When applied at 1 gallon of spray per 1000 sq ft this spray concentration is equivalent to a broadcast rate of 2/3 pint per acre Using the example of 3500 sq ft the applicator would use 3.5 x 1/4 fl oz = 7/8 fl oz of Transline in 3.5 gallons of water

Amount of Transline to Treat an Area of 1000 sq ft (pint/acre) (Mix in 1 Gallon or More of Spray)			
1/3	2/3	1	1 1/3
1/8 fl oz (3.6 mL)	1/4 fl oz (7.3 mL)	3/8 fl oz (11 mL)	1/2 fl oz (15 mL)

Tank Mix This product may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) tank mixing is not prohibited by the label of the tank mix product Read carefully and follow all applicable use directions precautions and limitations on the respective product labels

Specific Use Precautions and Restrictions

- In California do not exceed 2/3 pint per acre per annual use season
- Do not tank mix Transline with other herbicides labeled for use in cottonwood/poplar tree plantations unless allowed by all herbicide labels As a precaution it may be advisable to avoid contact with tree foliage
- A tank mix of Transline with another herbicide may be made as a foliar application over eucalyptus trees provided that it is specifically allowed by the label of the tank mix partner
- Certain broadleaf weeds such as mustards henbit chickweed kochia common lambsquarters pigweed Russian thistle and bindweed will not be controlled or suppressed

®Trademark of Dow AgroSciences LLC

R113 011
 EPA accepted ___/___/___
 Replaces R113 010