


printed April 6, 1994



page 2

(Front Panel of Label)

(logo) DowElanco

Turf Fertilizer Contains Confront\*

- Selectively controls annual and perennial broadleaf weeds
- Provides a feeding of fertilizer

Net Weight 34.21 lb (15.52 kg)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:** CAUTION, causes eye irritation. Harmful if inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not contaminate feed or foodstuffs. Do not contaminate water used for irrigation or domestic purposes. Do not graze treated areas. Do not feed clippings to livestock.

**Statement of Practical Treatment**

**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.  
**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**Environmental Hazards:** Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**Notice:** Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies".

2 of 7

(Front Panel of Label - Continued)

**Turf Fertilizer Contains Confront\***

**ACTIVE INGREDIENTS:**

triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid,  
triethylamine salt ..... 0.50%

clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid,  
triethylamine salt ..... 0.18%

**INERT INGREDIENTS:** ..... 99.32%

**Total** ..... 100.00%

**Acid Equivalent:**

triclopyr - 0.36%

clopyralid - 0.12%

**Guaranteed Analysis**

Total nitrogen (N) ..... XX%

X.X% ammoniacal nitrogen

XX.X% urea, methylene ureas nitrogen

X.X% water insoluble nitrogen

Available phosphoric acid (P<sub>2</sub>O<sub>5</sub>) ..... X%

Soluble potash (K<sub>2</sub>O) ..... X%

Derived from:

EPA Reg. No. 62719-XXX

EPA Est. XXXX-XX-X

Product of USA

\*Trademark of DowElanco

**[Editor's note: Registrant: DowElanco • Indianapolis, IN 46268-1054]**

Distributed by:

(Name and address of subregistrant)

---

**Recommended for Use by Professional Turfgrass Managers**

---

(Back Panel of Label)

**Directions for use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

**General Information**

This turf fertilizer product contains Confront herbicide which is a broad spectrum non-phenoxy weed killer that will control broadleaf weeds in established cool season and warm season turfgrass.

**Weeds Controlled by Turf Fertilizer Contains Confront**

Apply Turf Fertilizer Contains Confront to moist or dry foliage at the NORMAL RATE (0.65 lb ai/A) to control the following weeds:

- |                           |                                 |
|---------------------------|---------------------------------|
| black medic               | hop clover                      |
| bristly hawksbeard        | narrow leaf plantain (buckhorn) |
| broadleaf plantain        | red clover                      |
| catsear (coast dandelion) | white clover                    |
| common dandelion          |                                 |
| false dandelion           |                                 |
| hawkweed                  |                                 |

**Note:** Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

Apply Turf Fertilizer Contains Confront to moist foliage at the HEAVY RATE (0.75 lb ai/A) to control the following weeds.

- |                     |                          |
|---------------------|--------------------------|
| creeping beggarweed | roundleaf mallow         |
| dwarf beggarweed    | sheep sorrel             |
| burdock             | shepherdspurse           |
| coffee weed         | slender speedwell        |
| common chickweed    | smartweed                |
| mouseear chickweed  | spurweed                 |
| common cocklebur    | Canada thistle           |
| curly dock          | musk thistle             |
| dollarweed          | old world diamond flower |
| English lawn daisy  | common vetch             |
| goldenrod           | Virginia pepperweed      |
| ground ivy          | wild buckwheat           |
| henbit              | common yellow woodsorrel |
| lambquarters        | creeping woodsorrel      |
| lespedeza           | prostrate spurge         |
| matchweed           | wild violet              |
| poison ivy          | Virginia buttonweed      |
| common ragweed      |                          |

A repeat application may be needed for harder to control weeds in 4 to 6 weeks.

(Back Panel of Label - Continued)

Turf Fertilizer Contains Confront is recommended for use on the following turf species:

**Established Cool Season Turfgrasses**

Common Name

Scientific Name

bluegrass, Kentucky  
fescue, chewings  
fescue, creeping red  
fescues, sheeps  
fescue, tall  
ryegrass, perennial

*Poa pratensis*  
*Festuca rubra var. commutata*  
*Festuca rubra*  
*Festuca ovina*  
*Festuca arundinaceae*  
*Lolium perenne*

**Established Warm Season Turfgrasses†**

Common Name

Scientific Name

bahiagrass (Argentine or Pensacola)  
common Bermudagrass  
  
buffalograss  
centipedegrass  
zoysiagrass  
zoysiagrass  
fescue, tall (growing in warm season area)

*Paspalum spp.*  
*Cynodon dactylon*  
*(common bermuda only)*  
*Buchloe dactyloides*  
*Eremochloa ophiuroides*  
*Zoysia japonica*  
*Zoysia tenuifolia*  
*Festuca arundinaceae*

†Do not treat warm season turfgrasses with Turf Fertilizer Contains Confront when mowing height is less than 1/2 inch.

Do not use Turf Fertilizer Contains Confront on golf course putting greens and tees.

Turf Fertilizer Contains Confront may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply Turf Fertilizer Contains Confront only to healthy, well-established turf that has a well-anchored root system.

Common bermudagrass, centipedegrass, zoysiagrass and bahiagrass may be temporarily discolored by Turf Fertilizer Contains Confront.

Do not apply Turf Fertilizer Contains Confront to bentgrass; hybrid bermudagrass varieties such as Tifway, Tifdwarf, and Tifgreen; St. Augustinegrass; and ground covers such as lippia.

On bahiagrass and centipedegrass, do not apply before mid-March or until after the second mowing of the year. In the Fall, do not treat centipedegrass after the first of October or bahiagrass after the first November.

5 4 7

(Back Panel of Label - Continued)

### **Application Directions**

Make application using equipment that will ensure uniform coverage. Application should be made when weeds are actively growing. Good soil moisture conditions should be present before application to achieve optimal results. Application under drought conditions may provide less than desirable results. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled.

Additional applications should not be made less than four weeks apart to minimize grass injury. Newly seeded turf should be mowed four times before being treated.

### **General Use Precautions**

**Apply this product only as specified on this label.**

**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

Do not apply Turf Fertilizer Contains Confront directly to, or otherwise permit it to come into direct contact with vegetables, flowers, fruit or ornamental trees, cotton, grapes, tobacco, or other non-target broadleaf plants.

Do not apply when weather conditions favor drift from target areas. Drift or runoff may adversely affect non-target plants.

Do not apply to exposed roots of shallow rooted trees and shrubs and allow to contact exposed suckers or injury may occur.

Do not reseed until four weeks after application.

Do not apply at greater than recommended rates, especially under the branch spread (dripline) of trees.

Do not use grass clippings from turf treated with Turf Fertilizer Contains Confront for mulch.

Do not use compost containing grass clippings from turf treated with Turf Fertilizer Contains Confront in the growing season of application.

Do not apply on ditches used to transport irrigation water.

Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

5

6 4 7

Do not seed, sod or sprig after application. Delay until no sooner than four weeks after application.

(Back Panel of Label - Continued)

**Treatment of Turf Species Not Listed on the Confront Fertilizer Label**

Users who wish to use Confront on turf species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the treatment is safe to the target species. The user assumes the responsibility for any plant damage or other liability resulting from the use of Confront on species not recommended on this label.

**Storage and disposal**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

**Pesticide Disposal:** Pesticide wastes are toxic. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office of guidance.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Suggested Spreader Settings**

To provide proper distribution, calibrate spreader before application.

34.21 lb treats 11,000 ft at the NORMAL RATE (0.9 lb N/M) 0.65 lb ai/A  
 34.21 lb treats 9,529 ft at the HEAVY RATE (1.04 lb N/M) 0.75 lb ai/A

<u>SPREADER</u>	<u>GROUND SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>	
			<u>(lb product/acre)</u>	
			<u>NORMAL RATE</u>	<u>HEAVY RATE</u>
			<u>135.5</u>	<u>156.4</u>
Scotts Drop Type	3 mph	overlap wheels	X	X
Gandy	3 mph	overlap wheels	X	X
Scotts Rotaries				
R-7†, R-7X†	3 mph	X feet	X	X
R-8 (cone X)	3 mph	X feet	X	X
Lely models				
(see below)	4.5 mph	XX feet	X	X
Vicon (Model 02)	4.5 mph	XX feet	X	X
Vicon (Model 03)	4.5 mph	XX feet	X	X

†For the R-7X model, use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

**IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY**

**[Editor's note: Subregistrant may insert appropriate warranty and disclaimer information.]**