



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 6, 2017

Claire Terry
Regulatory Manager
Dow AgroSciences LLC
9330 Zionsville Road, 308/2E
Indianapolis, IN 46268-1054

Subject: Label Amendment – MOA and resistance-management info, fertilizer and state directions and restriction, claims, other edits
Product Name: Turf Fertilizer Contains Confront
EPA Registration Number: 62719-262
Application Date: May 9, 2017
Decision Number: 530005

Dear Ms. Terry:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 62719-262
Decision No. 530005

with FIFRA section 6. If you have any questions, please contact Mindy Ondish by phone at 703-605-0723, or via email at ondish.mindy@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending to the right.

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

(Base label):

Turf Fertilizer Contains Confront®

HERBICIDE

To be applied only under the direct supervision of licensed pesticide applicators responsible for turf weed control programs.

- Selectively controls annual and perennial broadleaf weeds
- Provides a feeding of fertilizer

[XX lbs] Covers up to XX,000 sq ft (XXX' x XX')
and/or

This bag covers XX,000 sq ft

Group	4	HERBICIDES
--------------	----------	-------------------

Active Ingredients:


triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid, triethylamine salt	0.50%
clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid, triethylamine salt.....	0.18%
Other Ingredients.....	99.32%
Total	100.00%

Acid Equivalent:

triclopyr	-	0.36%
clopyralid	-	0.12%

Guaranteed Fertilizer Analysis: XX-XX-XX †

Total Nitrogen (N).....	XX%
XX% Ammoniacal Nitrogen	
XX% Nitrate Nitrogen	
XX% Water-Insoluble Nitrogen*	
XX% Urea Nitrogen**	
XX% Slowly available water soluble Nitrogen.	
XX% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)***	XX%
Soluble Potash (K ₂ O).....	XX%
Calcium (Ca)	XX%
Magnesium (Mg)	XX%
XX% Water Soluble Magnesium (Mg)	
Sulfur****	
XX% Combined Sulfur (S)	
Boron (B)	XX%
Chlorine (Cl) [Max, Not more than]	XX%
Cobalt (Co).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
XX% Chelated Iron (Fe)	
XX% Water Soluble Iron (Fe) Manganese (Mn)XX%	
Water Soluble Manganese (Mn)	XX%



Molybdenum (Mo) XX%
 Nickel (Ni)..... XX%
 Sodium (Na)..... XX%
 Zinc (Zn)..... XX%
 XX% Chelated Zinc (Zn)

*[XX]% Water insoluble nitrogen from biosolids.
 **[XX]% coated slow release Nitrogen (N)
 **[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea
 **[XX]% Slowly available water soluble nitrogen
 **[XX]% Slowly available nitrogen from methylene ureas
 **[XX]% Slowly available nitrogen from ureaform
 **[XX]% Slowly available nitrogen from BioSolids
 **[XX]% Stabilized urea nitrogen [or [urea] nitrogen stabilized] with dicyandiamide
 ****[XX]% Free Sulfur (S)

Derived From: [XX][†]
 Primary plant foods derived from: [XX][†]
 Secondary plant foods derived from [XX][†]
 Iron derived from: [XX][†]

Density – XX lb./cu.ft.

[Editor’s Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]

[†] Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous sucate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdcic oxide, manganese sulfate, manganese sucate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molybdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

Keep Out Of Reach Of Children

WARNING

[Optional Label Claims]

This Bag Treats Up To _____ Square Feet

Controlled [Slow] release nitrogen for extended feeding

Contains Iron for deep greening

Iron enriched [formula]

Deep greening [formula]

Slow release nitrogen feeds your turfgrass over [x] months. [Note: The “x” represents the amount of months the fertilizer feeds depending on what we use (i.e. methylene urea, PCSCU)]

Phosphorus [Phosphate] free [formula] [Approved for use only on Phosphorus-Free Fertilizer/Carrier blends]

Zero Phosphorus [Approved for use only on Phosphorus-Free Fertilizer/Carrier blends]

Rich Green Turfgrass

Long-lasting Feeding

100% Satisfaction Guaranteed or Your Money Back!

Fortified with Milorganite [Note: Milorganite (logo/graphic) Need to move logo outside of label text (above product name, on side of label or below text)]

Contains Non-Burning, Slow-Release Nitrogen and Iron for Deep Greening

Contains Non-Burning Nitrogen

Contains Slow-Release Nitrogen

Controlled Release [Fertilizer] Technology feeds turfgrass] for 2+ months

UFLEXX™

UFLEXX™ is a trademark of AGROTAIN International, LLC.

UFLEXX™ is a trademark of Koch Fertilizer, LLC.

UMAXX®

UMAXX® is a registered trademark of AGROTAIN International, LLC.

Contains SCU [Sulfur Coated Urea]

N-Sure®

N-Sure is a registered trademark of Tessenderlo Kerley

with XCU®

XCU is registered trademark of Agrium

XCU is registered trademark of Koch Fertilizer, LLC

Apex-10™ Specialty Nutritional

Crystal Green® slow-release nutrient technology

Crystal Green is a registered trademark of Ostara Nutrient Recovery Technologies

Agrotain® Ultra

Agrotain is a registered trademark of Koch Agronomic Services

RXN XXX – [The % RxN (Agrotain Ultra) is indicated by the number built into the logo and can range from 20 to 100.]

STN XXX – [The % STN is indicated by the number built into the logo and can range from 25 to 100]

STU [Agrotain Ultra]

Growing a Better World [Harrell's Private Labels Only]

[Contains] Harrell's PROFertilizer

Harrell's Logo

Regain [Spring Valley Labels Only]

MESA® Patented [Lebanon Seaboard Private Labels Only]

Meth-Ex® [Lebanon Seaboard Private Labels Only]

Expo potassium sulfate [Lebanon Seaboard Private Labels Only]

With LSN – LebanonTurf Stabilized Nitrogen [Lebanon Seaboard Private Labels Only]

Green-Smart Technology [Lebanon Seaboard Private Labels Only]

Enhanced efficiency fertilizer GreenSmart for a Green Turfgrass [Lebanon Seaboard Private Labels Only]

Contains LESCO® Poly Plus® Polymer Coated Urea to provide uniform growth with extended nitrogen feeding. Poly Plus is comprised of Polymer Coated Urea. [John Deere Landscapes/LESCO Private Labels Only]

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. [John Deere Landscapes/LESCO Private Labels Only]

Wolf Trax DDP (Dispersable Dry Powder) Fe (Iron) Mn (Manganese), Zn (Zinc) Micronutrients

Professional Turf Grade/Premium Turf Grade/Mini Fairway Grade: Size Guide Number SGN XXX [The text box may have any combination of SGNs listed within the box. 80, 100, 145, 150, 195, 200, 210, 220, 230]

Duration XXX – Average nutrient release XXX days [The text box below may have any combination of the % Durations listed within the box. Duration 180 - Duration 120 - Duration 90 - Duration 45]

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.html>

Information regarding the contents and levels of metals in this product is available on the Internet at:

- **California Regulatory Site: <https://apps1.cdfa.ca.gov/FertilizerProducts/>**
- **Oregon Regulatory Site: http://oda.state.or.us/dbs/heavy_metal/search.lasso**
- **Washington Regulatory Site: <http://agr.wa.gov/> or <http://agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx>**
- **<http://www.aapfco.org/metals.html>**

- o <http://www.regulatory-info-lpi.com>

It is a violation of law in Maryland, New Hampshire, and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. [Maryland, New Hampshire, or New Jersey Applications]

No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period. [Virginia Applications]

Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to turf and sweep any product that lands on driveways, sidewalks, or streets back onto the turfgrass. [Maryland Applications]

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area. [Maryland Applications which contain Phosphorus] Not for use on established lawns or grass in MD. [If phos >5% and is not classified as a "starter" fertilizer]

In Virginia and New Jersey this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf. [Virginia and New Jersey Applications which contain Phosphorus]

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use. [FL Language for Specialty Products]

The maximum rate of application of this product must not exceed the lesser of the applicable rate determined from the [Table [XX]: Use Rates or Broadcast Application Rates], and the applicable rate determined in the State Nutrient Restrictions table.

STATE NUTRIENT RESTRICTIONS

State	Readily Available Nitrogen* lbs/1,000 sq ft	Total Nitrogen lbs/1,000 sq ft
MD, NH, VA	0.7	0.9
NJ	0.7	1.0
XXX	XXX	XXX

* Nitrogen present beyond the amount shown as Readily Available must be in the form of Slow-Release nitrogen

Note: Supplemental applications of fertilizer are needed to achieve a minimum 1lb Nitrogen per 1,000 sq ft.

Maximize your turfgrass

Stein Policy - 100% Satisfaction Guaranteed - If for any reason you are not satisfied with the results after using this premium product, you are entitled to replacement bags of fertilizer or your money back. Simply bring proof of purchase to any Stein Garden Center. [EC Grow Private Labels Only]

If, after using this product, you are dissatisfied with the results, simply mail the sales receipt, a brief description of the problem and your contact information to the address listed below for a refund of the purchase price of the product. EC Grow • PO Box 837 • Eau Claire, WI 54702 [EC Grow Private Labels Only]

If you have questions or comments regarding the quality of this product, please write to Turf Care Supply Corp. at the address below, or call Customer Service at 877-220-1014 between the hours of 9 A.M. and 5 P.M., Eastern Time, Monday through Friday. Please refer to the item number found on this bag when writing or calling. [Turf Care Supply Private Labels Only]

This product contains polymer-coated Enhanced Efficiency Fertilizer technology. As your lawn/turfgrass demands nutrients, nitrogen is released slowly into the soil for plant root uptake. This controlled-release approach eliminates surge growth and provides predictable feeding for over two months. All Menards Premium fertilizer products contain Enhanced Efficiency Fertilizer which ensures consistent feeding, deep greening, and promotes overall lawn/turfgrass health.

SPREAD IT & FORGET IT and the **SPREAD IT & FORGET IT** logo are registered trademarks of Koch Agronomic Services, LLC.

DURATION CR and the **DURATION CR** logo are registered trademarks of Koch Agronomic Services, LLC.

The **KOCH** logo is a trademark of Koch Industries, Inc. © 2014 Koch Agronomic Services, LLC.

STA-GREEN® is used under license to Central Garden and Pet Company

UFLEXX is a trademark of Koch Fertilizer LLC

CONTROLLED-RELEASE FERTILIZER

ENGINEERED FOR PERFORMANCE

DRIVEN BY DURATION CR

Contains **XCU Polymer Coated Sulfur Coated Urea**, a trademark of Koch Agronomic Services, LLC

Made in USA

Your Prescription for a Picture Perfect Lawn

Turf Care Supply Corp – Manufacturing

Turf Care Supply Corp - TCS GrowStar Premier Turf Products**Premier Turf Products**

Manufactured in the USA [Note: All inert materials must be produced in the US, and formulation must be manufactured in US to use this claim]

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING**

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin

Do not get in eyes or on clothing. Avoid contact with skin. Wear long-sleeved shirt and long pants, socks and shoes, protective eyewear and chemical resistant gloves made of rubber or other waterproof material. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 [or 1-800-XXX-XXXX] for emergency medical treatment information.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrigid containers any size)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers greater than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the

rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 [or 1-800-XXX-XXXX].

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-262

EPA Est. _____

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced by/for:
Manufactured by/for:
Distributed by/for:
Guaranteed by:

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

NET WEIGHT _____

(Label booklet cover):

Turf Fertilizer Contains Confront®

HERBICIDE

To be applied only under the direct supervision of licensed pesticide applicators responsible for turf weed control programs.

- Selectively controls annual and perennial broadleaf weeds
- Provides a feeding of fertilizer

[XX lbs] Covers up to XX,000 sq ft (XXX' x XX')
and/or

This bag covers XX,000 sq ft

Group	4	HERBICIDES
--------------	----------	-------------------

Active Ingredients:

triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid, triethylamine salt	0.50%
clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid, triethylamine salt.....	0.18%
Other Ingredients.....	99.32%
Total	100.00%

Acid Equivalent:

triclopyr	-	0.36%
clopyralid	-	0.12%

Guaranteed Fertilizer Analysis: XX-XX-XX †

Total Nitrogen (N).....	XX%
XX% Ammoniacal Nitrogen	
XX% Nitrate Nitrogen	
XX% Water-Insoluble Nitrogen*	
XX% Urea Nitrogen**	
XX% Slowly available water soluble Nitrogen.	
XX% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)***	XX%
Soluble Potash (K ₂ O).....	XX%
Calcium (Ca)	XX%
Magnesium (Mg)	XX%
XX% Water Soluble Magnesium (Mg)	
Sulfur****	
XX% Combined Sulfur (S)	
Boron (B)	XX%
Chlorine (Cl) [Max, Not more than]	XX%
Cobalt (Co).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
XX% Chelated Iron (Fe)	
XX% Water Soluble Iron (Fe) Manganese (Mn)XX%	
Water Soluble Manganese (Mn)	XX%

Molybdenum (Mo) XX%
 Nickel (Ni)..... XX%
 Sodium (Na)..... XX%
 Zinc (Zn)..... XX%
 XX% Chelated Zinc (Zn)

*[XX]% Water insoluble nitrogen from biosolids.
 **[XX]% coated slow release Nitrogen (N)
 **[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea
 **[XX]% Slowly available water soluble nitrogen
 **[XX]% Slowly available nitrogen from methylene ureas
 **[XX]% Slowly available nitrogen from ureaform
 **[XX]% Slowly available nitrogen from BioSolids
 **[XX]% Stabilized urea nitrogen [or [urea] nitrogen stabilized] with dicyandiamide
 ****[XX]% Free Sulfur (S)

Derived From: [XX][†]
 Primary plant foods derived from: [XX][†]
 Secondary plant foods derived from [XX][†]
 Iron derived from: [XX][†]

Density – XX lb./cu.ft.

[Editor’s Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]

[†] Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous sucate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdcic oxide, manganese sulfate, manganese sucate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molybdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

Keep Out Of Reach Of Children

WARNING

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 [or 1-800-XXX-XXXX].

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-262

EPA Est. _____

®Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

Produced by/for:

Manufactured by/for:
Distributed by/for:
Guaranteed by:

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

NET WEIGHT _____

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin

Do not get in eyes or on clothing. Avoid contact with skin. Wear long-sleeved shirt and long pants, socks and shoes, protective eyewear and chemical resistant gloves made of rubber or other waterproof material. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 [or 1-800-XXX-XXXX] for emergency medical treatment information.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Product Information

This turf fertilizer product contains Confront® herbicide which is a broad spectrum weed killer that will control broadleaf weeds in established cool season and warm season turfgrass.

Turf Fertilizer Contains Confront may be used on the following turf species:

Common Name

bluegrass, Kentucky
fescue, chewings
fescue, creeping red
fescues, sheeps
fescue, tall
ryegrass, perennial

Scientific Name

Established Cool Season Turfgrass

Poa pratensis
Festuca rubra var. commutata
Festuca rubra
Festuca ovina
Festuca arundinaceae
Lolium perenne

Established Warm Season Turfgrass¹

bahiagrass (Argentine or Pensacola)	<i>Paspalum</i> spp.
buffalograss	<i>Buchloe dactyloides</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
bermudagrass ²	<i>Cynodon dactylon</i>
fescue, tall (growing in warm season area)	<i>Festuca arundinaceae</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

¹Do not treat warm season turfgrasses with Turf Fertilizer Contains Confront when mowing height is less than 1/2 inch.

²Do not apply Confront to bermudagrass on sod farms.

Use Precautions:

Turf Fertilizer Contains Confront may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply Turf Fertilizer Contains Confront only to healthy, well-established turf that has a well-anchored root system.

Bermudagrass, centipedegrass, zoysiagrass, and bahiagrass may be temporarily discolored by Turf Fertilizer Contains Confront.

Use Restrictions

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

In **California, Oregon** and **Washington**, turfgrass and lawn uses are restricted to golf courses only.

Do not use on residential turf. Turfgrass and lawn uses are restricted to non-residential sites.

Apply this product only as specified on this label.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled.

Do not apply to exposed roots of shallow rooted trees and shrubs and allow to contact exposed suckers or injury may occur.

Do not apply Turf Fertilizer Contains Confront directly to, or otherwise permit it to come into direct contact with, cotton, grapes, tobacco, vegetables, flowers, fruit or ornamental trees, or other nontarget broadleaf plants.

Do not reseed until 4 weeks after application.

Do not use Turf Fertilizer Contains Confront on golf course putting greens and tees.

Do not send grass clippings to a compost facility.

Do not collect grass clippings for mulch or compost.

Applicator must give notice to landowners/property managers to not use grass clippings for composting.

Do not apply on ditches used to transport irrigation water.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

Do not contaminate feed or foodstuffs.

Do not graze treated areas. Do not feed clippings to livestock.

Do not apply when weather conditions favor drift from target areas. Drift or runoff may adversely affect nontarget plants.

Do not apply at greater than specified rates, especially under the branch spread (dripline) of trees.

Do not seed, sod, or sprig after application. Delay until no sooner than 4 weeks after application.

Do not apply Turf Fertilizer Contains Confront to bentgrass; hybrid bermudagrass varieties such as tifway, tifdwarf, and tifgreen; St. Augustinegrass; and ground covers such as lippia.

On bahiagrass and centipedegrass, do not apply before mid-March or until after the second mowing of the year.

In the fall, do not treat centipedegrass after the first of October or bahiagrass after the first November

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [1] [2] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [1] [2] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [1] [2] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [1] [2] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [1] [2] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [1] [2] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [1] [2] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [1] [2] applications per year on zoysiagrass in the south.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.
- Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Treatment of Turfgrass Species Not Listed on the Confront Fertilizer Label

Users who wish to use Turf Fertilizer Contains Confront on turfgrass species not listed on this label may determine the suitability for such uses by treating a small area at the specified rate. Prior to treatment of larger areas, observe the treated area for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the treatment is safe to the target species. The user assumes the responsibility for any plant damage or other liability resulting from the use of Turf Fertilizer Contains Confront on species not listed on this label.

Application Directions

Use equipment that will ensure uniform coverage and apply 3.58 lbs per 1000 sq feet (156.46 lb per acre, equivalent to 0.187 lb ae/acre clopyralid and 0.56 lb ae/acre of triclopyr) and [XX] lbs of Nitrogen per 1000 sq ft when weeds are actively growing. Good soil moisture conditions should be present before application to achieve optimal results. Application under drought conditions may provide less than

desirable results. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled.

To minimize the potential for turfgrass injury, do not make additional applications within 4 weeks of a previous application. Apply to newly seeded or sprigged turfgrass only after the fourth mowing or after the turfgrass seedlings have tillered and developed an adequate secondary root system.

- Do not apply more than 2 applications per year and do not use more than 0.375 lb ae clopyralid/1.125 lb ae triclopyr (312.5 lbs of Turf Fertilizer Contains Confront) per acre per year.
- In Florida and New York, the maximum use rate is 0.25 lb ae clopyralid/0.75 lb ae triclopyr per acre (205.6 lbs of Turf Fertilizer Contains Confront) per acre per year.

Suggested Spreader Settings

[Editor's note: With available data Supplemental Distributor may add the following statement.] These settings were calibrated and field tested. However, age, condition of spreader speed of operations and evenness of terrain may require slightly different settings for desired coverage.

[Editor's note: Supplemental Distributor may add a table listing recommended application equipment and calibration settings.]

To provide proper distribution, calibrate spreader before application. [Check equipment frequently to verify calibration and proper functioning.]

<u>Spreader</u>	<u>Ground Speed</u>	<u>Width of Coverage</u>	<u>Spreader Settings</u> (156.46 lb product/acre)
Scotts Drop Type	3 mph	overlap wheels	X
Gandy	3 mph	overlap wheels	X
Scotts Rotaries			
R-7 ¹ , R-7X ¹	3 mph	X feet	X
R-8 (cone X)	3 mph	X feet	X
Lely models			
(see below)	4.5 mph	XX feet	X
Vicon (Model 02)	4.5 mph	XX feet	X
Vicon (Model 03)	4.5 mph	XX feet	X

¹For the R-7X model, use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR, and 1250. PTO at 450 rpm.
[AND/OR]

Calibration Settings	
Brand and Model of Spreader	Calibration Setting for 3.58 lb of Product per 1000 sq ft
XXXXXXX	XX
XXXXXXX	
XXXXXXX	

[*These settings are approved for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product]

Weeds Controlled

Apply Turf Fertilizer Contains Confront to moist or dry foliage at 3.58 lbs per 1000 sq feet (156.46 lb per acre, equivalent to 0.187 lb ae/acre clopyralid and 0.56 lb ae/acre of triclopyr) to control the following weeds:

black medic	hawkweed
bristly hawksbeard	hop clover
broadleaf plantain	narrow leaf plantain (buckhorn)
catsear (coast dandelion)	red clover
common dandelion	white clover
false dandelion	
burdock	lespedeza
Canada thistle	matchweed
coffee weed	mouseear chickweed
common chickweed	musk thistle
common cocklebur	old world diamond flower
common ragweed	poison ivy
common vetch	prostrate spurge
common yellow woodsorrel	roundleaf mallow
creeping beggarweed	sheep sorrel
creeping woodsorrel	shepherdspurge
curly dock	slender speedwell
dollarweed	smartweed
dwarf beggarweed	spurweed
English lawn daisy	Virginia buttonweed
goldenrod	Virginia pepperweed
ground ivy	wild buckwheat
henbit	wild violet
lambsquarters	

Note: Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

For harder to control weeds, a repeat application in 4 to 6 weeks may be necessary.

Weed Resistance Management:

Triclopyr and clopyralid, the active ingredients in this product, are Group 4 herbicides based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 4 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 4 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds or to report herbicide failures.

Best Management Practices:

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. It is

recommended to scout for weeds before Turf Fertilizer Contains Confront application for identification and growth stage and after application to facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
EPA accepted __/__/__