P1125

£

ł

)

)

page 1

1778

Amendment 08/09/95

Front Panel of Label

[photo: house with beautiful lawn] (inset photo: dandelion]

Kills dandelions

(Subregistrant) (Logo) Lawn Fertilizer Plus Confront^{*} Weed Control

For use on bahiagrass, common bermudagrass, bluegrass, centipedegrass, fescue, ryegrass, or zoysiagrass lawns

-- The (Subregistrant) Difference --

- Thick, green, weed-free turf
- Timed-release nitrogen won't burn your lawn or cause surge growth
- Contains the most effective weed control you can buy
- (Subregistrant) No-Quibble Guarantee%
- Toll-free Consumer Hotline: 1-800-000-0000

Patented Particle Technology

[art]

Unlike other weed-and-feeds, every particle of (Subregistrant) Lawn Fertilizer Plus Confront contains

- (Subregistrant)'s patented timed-release nitrogen
- Phosphorus
- Potassium
- Exclusive, highly effective weed control

[Editor's note: The above section may be included, if applicable.]

Do not use on St. Augustinegrass, hybrid bermudagrass, carpetgrass, bentgrass, dichondra, or lippia lawns

Treats and feeds 5,000 sq. ft.

NET WEIGHT 15.55 lb. (7.05 kg)

CAUTION WARNING: KEEP OUT OF REACH OF CHILDREN See back for additional precautionary statements.

Trademark of DowElanco

ACCEPTED SEP - 7 1995 Under the Foderal Insecticide, Fungicide, and Redenticide Act. us amonded, fir the presticide registared unday EPA Rog. No.

R2A FERTLZ LABELS US

27

....

1

`}

)

248

Lawn Fertilizer Plus Confront Weed Control

. •

ACTIVE INGREDIENTS: triclopyr: 3,5,6 trichloro-2- pyridinyloxyacetic
acid, triethylamine salt0.50%
clopyralid: 3,6-dichloro-2-
pyridinyloxyacetic acid,
triethylamine salt0.18%
INERT INGREDIENTS
Total
Acid Equivalent: triclopyr - 0.38% clopyralid - 0.12%
Guaranteed Analysis
Total nitrogen (N)XX%
X.X% ammoniacal nitrogen
XX.X% urea, methylene ureas nitrögen
X.X% water insoluble nitrogen
Available phosphoric acid (P2O5)X%
Soluble potash (K ₂ O)X%

Derived from:

EPA Reg. No. 62719-263

Product of USA

Distributed by: (Name and address of subregistrant)

> PULL TAB and remove for pour spout Easy [Illus.] Open PULL HERE

Treats and feeds 5,000 sq. ft.

EPA Est. XXXX-XX-X

R2A FERTLZ LABELS US

ŧ

)

)

KEEP OUT OF REACH OF CHILDREN CAUTION WARNING PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION-WARNING, Gauses eye irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear waterpreef gloves, long-sleeved shirt, long pants; and shoes plus seeks when handling or applying this product. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or face shield. Harmful if swallowed or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If in eyes: Flush with plenty of water. Get medical attention if imitation persists. Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention if imitation persists. If on skin: Wash with plenty of soap and water. Get medical attention if imitation persists. If swallowed: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loarny sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your locat agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

348

29

Amendment 08/09/95

page 4

Lawn Fertilizer Plus Confront Weed Control

- The (Subregistrant) Difference -

(Subregistrant's) Patented Particle Technology is available only in (Subregistrant @) products, so don't be fooled by weed-and-feeds claiming to be "just like (Subregistrant's)."

Remember, even weed-and-feeds claiming to contain slow- or controlled-release nitrogen are typically just simple mixes, so they can't deliver the consistent great results you'll get from (Subregistrant's) Patented Particle Technology. With (Subregistrant) Lawn Fertilizer Plus Confront Weed Control, every particle contains time-release nitrogen, plus the other primary nutrients and the most effective weed control you can buy.

[photos: simple mix and LF+CWC particles]

ORDINARY SIMPLE MIXES

)

}

- Nutrient and weed control granules of many different sizes and shapes
- Result: granules of unequal size can separate during application

[photos: simple mix and LF+CWC patterns]

ORDINARY SIMPLE MIXES

- The amount of individual nutrients and weed control in each square inch of lawn may vary greatly
- Result: uneven, inconsistent weeding and feeding

(Subregistrant's) TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control granules are consistent in size and shape
- Result: consistent distribution of nutrients and weed control

(Subregistrant's) TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control delivers the same ratio of primary nutrients plus weed control to every square inch of turf
 Result: consistent greening
- Result: consistent greening plus better weed control

What to expect

Dandelions and other weeds will be killed in 4 to 5 weeks and your lawn will be greener in 4 to 6 days.

Small, lightweight granules mean better weed control. Most lawn weed controls must be absorbert through the weeds' leaves to be effective. Lawn Fertilizer Plus Confront Weed Control granules are small and lightweight, so they will stay on even the smallest weed leaves. Weed control in Lawn Fertilizer Plus Confront Weed Control is absorbed by the weeds' leaves and roots, killing the plant completely.

Apply to moist or dry foliage at the REGULAR RATE to control the following weeds:

black medic bristly hawksbeard broadleaf plantain catsear (coast dandelion) hop clover narrow leaf plantain (buckhorn) red clover white clover Amendment 08/09/95

548

common dandelion false dandelion hawkweed

Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

Apply to moist foliage at the HEAVY RATE to control the following weeds.

creeping beggarweed dwarf beggarweed burdock coffee weed common chickweed mouseear chickweed common cocklebur curly dock dollarweed English lawn daisy goldenrod ground ivy henbit lambsquarters lespedeza matchweed poison ivy common ragweed

)

)

roundleaf mailow sheep sorrel shepherdspurse slender speedwell smartweed spurweed Canada thistle musk thistle old world diamond flower common vetch Virginia pepperweed wild buckwheat common yellow woodsorrel creeping woodsorrel prostrate spurge wild violet Virginia buttonweed

A repeat application may be necessary.

How and when to apply

Apply in late spring, summer, or fall

Directions for usa

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

- Apply anytime weeds are actively growing (when daytime temperatures are consistently above 60_F). Avoid applying when temperatures are consistently above 90_F or during drought conditions. Applications during these conditions may provide less control than desirable.
- Apply on a calm day.
- Lawn Fertilizer Plus Confront Weed Control is absorbed by both foliage and roots. Weeds listed in the REGULAR RATE section above can be controlled by application to dry foliage. For best results, apply when good moisture conditions are present.
- To control weeds listed in the HEAVY RATE section, apply to moist grass. To help particles stick to weed leaves, the grass should be thoroughly watered or wet from dew before applying Lawn Fertilizer Plus Confront Weed Control.
- In the Fall, apply up to the last mowing or as long as grasses are actively growing.

<u>R2A FERTLZ LABELS US</u>

Ś

64?

- On bahia and centipedegrass lawns, wait until after the second mowing of the year, but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not use this product on flowers, shrubs, fruit, or vegetable plants or on creeping bentgrass, St. Augustinegrass, dichondra, or lippia lawns. Do not apply this product to hybrid bermudagrass varieties such as Tifdwarf, Tifgreen, and Tifway. On common bermudagrass, bahia, centipedegrass, or zoysia lawns, under certain conditions some temporary discoloration may result.

Watering or rain

Do not water your lawn for 24 hours after application. Do not apply when rain is expected within 24 hours. Water will wash the weed control product off the weed leaves and reduce it's effectiveness.

IMPORTANT What to Expect

Weed stems and leaves will begin to twist and curl within a week, but a month or more may be required for complete control. If weed control is incomplete after 4 to 5 weeks, a repeat application can be made. To minimize grass injury, additional applications should be at least 4 weeks apart.

Seeding/Sodding/Sprigging

Wait until the new grass has been mowed at least four times before applying Lawn Fertilizer Plus Confront Weed Control.

If you are planning to seed, sod, or sprig, you should wait at least one month after you have applied Lawn Fertilizer Plus Confront Weed Control.

This product will control only actively growing weeds; it will not prevent weed seeds from sprouting new plants.

Spreading Tips

)

For best results, use a (_____) spreader.

Be careful when applying this product near shrubs, flower beds, or vegetable gardens, as contact with this weed control product may damage these desirable plants. When using any rotary spreader, take special care near desirable plants.

Do not contaminate feed or foodstuffs.

Do not contaminate water used for imigation or domestic purposes.

Do not graze treated areas. Do not feed clippings to livestock.

Do not apply in a brisk wind, as drift may contact desirable plants and cause injury.

Do not apply to exposed roots of shallow-rooted trees and shrubs or allow contact with exposed suckers, as injury may occur.

Do not reseed until 4 weeks after application.

Do not use Lawn Fertilizer Plus Confront Weed Control on golf course putting greens or tees.

Do not apply at greater than recommended rates, especially under the branch spread (cripline) of trees.

Do not use grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control for mulch.

Do not use compost containing grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control in the growing season of application.

R2A FERTLZ LABELS US

ļ

)

)

page 7

1.10

	REGULAR RATE covers 5,000 sq. ft.	HEAVY RATE covers 4,300 sq. ft.	DIRE	CTIONS	
Scotts Drop Spreaders	xx	XX	2 hea ionge lawn,	lawn first; apply ider strips; apply ist direction of overlapping I marks.	
Scotts EasyGreen® Spreaders	xx	XX	30 in previ	e each pass about ches from ous pass to ap spread width.	
Scotts SpeedyGreen® Rotary	xx	xx	Make each pass about 5 feet from previous pass to overlap spread width.		
Additional Spreader Settings			GULAR	HEAVY RATE_	
EZ Drop, Republic (#530) EZ Broadcast, Republic (#540) EZ Grow, Republic (#545) Red Devil Drop (FH-22) Red Devil Rotary (SB-50)			XX XX XX XX XX	XX XX XX XX XX	

Spreader Settings

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage).

[illustrations; spreading patters]

Quaker Drop (Model #10850)

Ortho Drop

Ortho Rotary

Quaker Rotary (Model #10301)

STORAGE: Store in original container in a clean, dry area away from extremes of heat and cold ar.d out of direct sunlight.

XX

XX

XX

XX

XX

XX

XX

XX

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. When completely empty, discard bag in trash.

Questions? Call (Subregistrant)'s toll-free Consumer Hotline: 1-800-000-0000-

(Subregistrant's) No-Quibble Guarantee **

Amendment 08/09/95

ί,

)

)

page 8

8.78

34

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

(Subregistrant). . . the lawn people (Subregistrant Address)

[Editor's note: Registrant is DowElanco • Indianapolis, IN 46268-1054]