

PM 25 62719-263

10/9

R2A/Lawn Fertilizer Plus Confront/Amend/02-10-97
file: I263feb97d.doc

Registration Notes for Previous Labeling Actions:

Lawn Fertilizer Plus

Confront* Weed Control

EPA Reg. No. 62719-263

Final Printed Labeling: Label text based on EPA-accepted copy dated September 7, 1995.

Amendments EPA-accepted 09/07/95:

Changes to Precautionary Statements based on EPA acute toxicity review dated 04/10/95:

1. Signal word changed from CAUTION to WARNING based on primary eye irritation study.
2. Statement of Hazard statement and Statements for Practical Treatment modified in accordance with EPA instructions except that "get medical attention if irritation persists" was retained for eye and skin exposure routes.

Final Printed Labeling: Label text based on EPA-accepted copy dated April 4, 1995.

Changes to Final Printed Labeling:

1. Revised precautionary statement as per EPA-condition of acceptance.
2. Moved information not included by EPA as part of specified precautionary statements to General Use Precautions section.

*Trademark of DowElanco

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated
 MAR 20 1997
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.
62719-263

2019

(Front Panel of Label)

[photo: house with beautiful lawn]
(inset photo: dandelion)

Kills dandelions

(Subregistrant) (Logo)
**Lawn Fertilizer Plus
Confront* Weed Control**

For use on bahiagrass, common bermudagrass, bluegrass, centipedegrass, fescue, ryegrass, or zoysiagrass lawns.

— The (Subregistrant) Difference —

- Thick, green, weed-free turf
- Timed-release nitrogen won't burn your lawn or cause surge growth
- Contains the most effective weed control you can buy
- (Subregistrant) No-Quibble Guarantee™
- Toll-free Consumer Hotline: 1-800-000-0000

Patented Particle Technology

[art]

Unlike other weed-and-feeds, every particle of (Subregistrant) Lawn Fertilizer Plus Confront contains

- (Subregistrant)'s patented timed-release nitrogen
- Phosphorus
- Potassium
- Exclusive, highly effective weed control

[Editor's note: The above section may be included, if applicable.]

Do not use on St. Augustinegrass, hybrid bermudagrass, carpetgrass, bentgrass, dichondra, or lippia lawns.

Treats and feeds 5,000 sq ft


NET WEIGHT 15.55 lb (7.05 kg)

WARNING: KEEP OUT OF REACH OF CHILDREN

See back for additional precautionary statements.

*Trademark of DowElanco

Lawn Fertilizer Plus Confront Weed Control



3 of 9

ACTIVE INGREDIENTS:

triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid, triethylamine salt 0.50%

clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid, triethylamine salt 0.18%

INERT INGREDIENTS 99.32%

Total 100.00%

Acid Equivalent:

triclopyr - 0.36%

clopyralid - 0.12%

Guaranteed Analysis

Total nitrogen (N) XX%

 X.X% ammoniacal nitrogen

 XX.X% urea, methylene ureas nitrogen

 X.X% water insoluble nitrogen

Available phosphoric acid (P₂O₅) X%

Soluble potash (K₂O) X%

Derived from:

EPA Reg. No. 62719-263

EPA Est. XXXX-XX-X

Product of USA

Distributed by:
(Name and address of subregistrant)

PULL TAB
and remove for
pour spout
Easy [illus.]
Open PULL HERE

Treats and feeds 5,000 sq ft

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: **WARNING,** Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or face shield. Harmful if swallowed or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

4079

Statement of Practical Treatment

If in eyes: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

5 of 9

Lawn Fertilizer Plus Confront Weed Control

— The (Subregistrant) Difference —

(Subregistrant's) Patented Particle Technology is available only in (Subregistrant ®) products, so don't be fooled by weed-and-feeds claiming to be "just like (Subregistrant's)."

Remember, even weed-and-feeds claiming to contain slow- or controlled-release nitrogen are typically just simple mixes, so they can't deliver the consistent great results you'll get from (Subregistrant's) Patented Particle Technology. With (Subregistrant) Lawn Fertilizer Plus Confront Weed Control, every particle contains time-release nitrogen, plus the other primary nutrients and the most effective weed control you can buy.

[photos: simple mix and LF+CWC particles]

ORDINARY SIMPLE MIXES

- Nutrient and weed control granules of many different sizes and shapes
- Result: granules of unequal size can separate during application

(Subregistrant's) TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control granules are consistent in size and shape
- Result: consistent distribution of nutrients and weed control

[photos: simple mix and LF+CWC patterns]

ORDINARY SIMPLE MIXES

- The amount of individual nutrients and weed control in each square inch of lawn may vary greatly
- Result: uneven, inconsistent weeding and feeding

(Subregistrant's) TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control delivers the same ratio of primary nutrients plus weed control to every square inch of turf
- Result: consistent greening plus better weed control

What to expect

Dandelions and other weeds will be killed in 4 to 5 weeks and your lawn will be greener in 4 to 6 days.

Small, lightweight granules mean better weed control. Most lawn weed controls must be absorbed through the weed's leaves to be effective. Lawn Fertilizer Plus Confront Weed Control granules are small and lightweight, so they will stay on even the smallest weed leaves. Weed control in Lawn Fertilizer Plus Confront Weed Control is absorbed by the weed's leaves and roots, killing the plant completely.

Apply to moist or dry foliage at the REGULAR RATE to control the following weeds:

black medic	hop clover
bristly hawksbeard	narrow leaf plantain (buckhorn)
broadleaf plantain	red clover
catsear (coast dandelion)	white clover
common dandelion	
false dandelion	
hawkweed	

6049

Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

Apply to moist foliage at the **HEAVY RATE** to control the following weeds.

creeping beggarweed	roundleaf mallow
dwarf beggarweed	sheep sorrel
burdock	shepherdspurse
coffee weed	slender speedwell
common chickweed	smartweed
mouseear chickweed	spurweed
common cocklebur	Canada thistle
curly dock	musk thistle
dollarweed	old world diamond flower
English lawn daisy	common vetch
goldenrod	Virginia pepperweed
ground ivy	wild buckwheat
henbit	common yellow woodsorrel
lambquarters	creeping woodsorrel
lespedeza	prostrate spurge
matchweed	wild violet
poison ivy	Virginia buttonweed
common ragweed	

A repeat application may be necessary.

How and when to apply

Apply in late spring, summer, or fall.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

- Apply anytime weeds are actively growing (when daytime temperatures are consistently above 60°F). Avoid applying when temperatures are consistently above 90°F or during drought conditions. Applications during these conditions may provide less control than desirable.
- Apply on a calm day.
- Lawn Fertilizer Plus Confront Weed Control is absorbed by both foliage and roots. Weeds listed in the **REGULAR RATE** section above can be controlled by application to dry foliage. For best results, apply when good moisture conditions are present.
- To control weeds listed in the **HEAVY RATE** section, apply to moist grass. To help particles stick to weed leaves, the grass should be thoroughly watered or wet from dew before applying Lawn Fertilizer Plus Confront Weed Control.
- In the Fall, apply up to the last mowing or as long as grasses are actively growing.
- On bahia and centipedegrass lawns, wait until after the second mowing of the year, but **NOT** before mid-March. In the fall, **DO NOT** treat centipede after the first of October and **DO NOT** treat bahia after the first of November.
- Do not use this product on flowers, shrubs, fruit, or vegetable plants, or on creeping bentgrass, St. Augustinegrass, dichondra, or lippia lawns. Do not apply this product to hybrid bermudagrass varieties such as Tifdwarf, Tifgreen, and Tifway. On common bermudagrass, bahia, centipedegrass, or zoysia lawns, under certain conditions some temporary discoloration may result.

7019

Watering or rain

Do not water your lawn for 24 hours after application. Do not apply when rain is expected within 24 hours. Water will wash the weed control product off the weed leaves and reduce its effectiveness.

IMPORTANT**What to Expect**

Weed stems and leaves will begin to twist and curl within a week, but a month or more may be required for complete control. If weed control is incomplete after 4 to 5 weeks, a repeat application can be made. To minimize grass injury, additional applications should be at least 4 weeks apart.

Seeding/Sodding/Sprigging

Wait until the new grass has been mowed at least 4 times before applying Lawn Fertilizer Plus Confront Weed Control.

If you are planning to seed, sod, or sprig, you should wait at least 1 month after you have applied Lawn Fertilizer Plus Confront Weed Control.

This product will control only actively growing weeds; it will not prevent weed seeds from sprouting new plants.

Spreading Tips

For best results, use a (_____) spreader.

Be careful when applying this product near shrubs, flower beds, or vegetable gardens, as contact with this weed control product may damage these desirable plants. When using any rotary spreader, take special care near desirable plants.

Do not contaminate feed or foodstuffs.

Do not contaminate water used for irrigation or domestic purposes.

Do not graze treated areas. Do not feed clippings to livestock.

Do not apply in a brisk wind, as drift may contact desirable plants and cause injury.

Do not apply to exposed roots of shallow-rooted trees and shrubs or allow contact with exposed suckers, as injury may occur.

Do not reseed until 4 weeks after application.

Do not use Lawn Fertilizer Plus Confront Weed Control on golf course putting greens or tees.

Do not apply at greater than recommended rates, especially under the branch spread (dripline) of trees.

Do not use grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control for mulch.

Do not use compost containing grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control in the growing season of application.

8 of 9

Spreader Settings

	REGULAR RATE covers 5,000 sq ft	HEAVY RATE covers 4,300 sq. ft.	DIRECTIONS
Scotts Drop Spreaders	XX	XX	Mow lawn first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts EasyGreen® Spreaders	XX	XX	Make each pass about 30 inches from previous pass to overlap spread width.
Scotts SpeedyGreen® Rotary	XX	XX	Make each pass about 5 feet from previous pass to overlap spread width.

Additional Spreader Settings

	REGULAR RATE	HEAVY RATE
EZ Drop, Republic (#530)	XX	XX
EZ Broadcast, Republic (#540)	XX	XX
EZ Grow, Republic (#545)	XX	XX
Red Devil Drop (FH-22)	XX	XX
Red Devil Rotary (SB-50)	XX	XX
Quaker Drop (Model #10850)	XX	XX
Quaker Rotary (Model #10301)	XX	XX
Ortho Drop	XX	XX
Ortho Rotary	XX	XX

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

[illustrations; spreading patterns]

Storage and Disposal

Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. When completely empty, discard bag in trash.

Questions? Call (Subregistrant)'s toll-free Consumer Hotline: 1-800-000-0000

2 of 9

(Subregistrant's) No-Quibble Guarantee™

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

(Subregistrant) . . . the lawn people
(Subregistrant Address)

[Editor's note: Registrant is DowElanco • Indianapolis, IN 46268-1054]

*Trademark of DowElanco
EPA-Accepted: 09/07/95