627/9-263

9/29/2000

Page / 3 8

R2A / Lawn Fertilizer Contains Confront . Amend With Edits / 09-12-00

page 1

(Front Panel of Label)

[photo: house with beautiful lawn] [inset photo: dandelion]

Kills dandelions

[Subregistrant] (Logo)
Lawn Fertilizer Plus
Confront\* Weed Control

For use on bahiagrass, common bermudagrass, bluegrass, centipedegrass, fescue, ryegrass, or zoysiagrass lawns.

-- The [Subregistrant] Difference --

· Thick, green, weed-free turf

- Timed-release nitrogen won't burn your lawn or cause surge growth
- Contains the most effective weed control you can buy
- [Subregistrant] No-Quibble Guarantee ]
- Toll-free Consumer Hotline: 1-800-000-0000

ACCEPTED
with COMMENTS
In EPA Letter Dated:

SEP 2 9 2000
Under the Federal Insecticide,
Fungicide, and Redouticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

### Patented Particle Technology [art]

Unlike other weed-and-feeds, every particle of (Subregistrant) Lawn Fertilizer Plus Confront contains

- (Subregistrant)'s patented timed-release nitrogen
- Phosphorus
- Potassium
- · Exclusive, highly effective weed contro

[Editor's note: The above section may be included, if applicable.]

Do not use on St. Augustinegrass, hybrid bermudagrass, carpetgrass, bentgrass, dichondra, or lippia lawns.

Treats and feeds 5,000 sq ft

NET WEIGHT 15.55 lb (7.05 kg)

# KEEP OUT OF REACH OF CHILDREN WARNING

See back for additional precautionary statements.

\*Trademark of Dow AgroSciences LLC

Lawn Fertilizer Plus Confront Weed Control

page 2

**ACTIVE INGREDIENTS:** 

triclopyr: 3,5,6 trichloro-2pyridinyloxyacetic acid,

triethylamine salt ......0.50%

clopyralid: 3,6-dichloro-2pyridinyloxyacetic acid,

Acid Equivalent:

triclopyr - 0.36% clopyralid - 0.12%

Guaranteed Analysis

Total nitrogen (N) ......XX.X%

X.X% ammoniacal nitrogen

XX.X% urea, methylene ureas nitrogen

X.X% water insoluble nitrogen

Derived from:

EPA Reg. No. 62719-263

EPA Est. XXXX-XX-X

Product of USA

Distributed by:

[Name and address of subregistrant]

PULL TAB
and remove for
pour spout
Easy [Illus.]
Open PULL HERE

Treats and feeds 5,000 sq ft

# KEEP OUT OF REACH OF CHILDREN WARNING

#### PRECAUTIONARY STATEMENTS

#### **Hazards to Humans and Domestic Animals**

WARNING, Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or face shield. Harmful if swallowed or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin

Do not get in eyes or on clothing. Avoid contact with skin. Wear long-sleeved shirt and long pants, socks and shoes, gogles or face shield and chemical resistant gloves made of rubber or other waterproof material. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

#### **User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### Statement of Practical Treatment First Aid

If in eyes: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Wash with plenty of soap and water. Get medical attention if irritation persists Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Gall a doctor or get medical attention. Do not induce vermiting. Drink premptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

#### **Environmental Hazards**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

# Lawn Fertilizer Plus Confront Weed Control

-- The [Subregistrant's Name] Difference --

[Subregistrant's] Patented Particle Technology is available only in [Subregistrant ®] products, so don't be fooled by weed-and-feeds claiming to be "just like [Subregistrant's]"

Remember, even weed-and-feeds claiming to contain slow- or controlled-release nitrogen are typically just simple mixes, so they can't deliver the consistent great results you'll get from (Subregistrant's) Patented Particle Technology. With (Subregistrant) Lawn Fertilizer Plus Confront Weed Control, every particle contains time-release nitrogen, plus the other primary nutrients and the most effective weed control you can buy.

#### [photos: simple mix and LF+CWC particles]

#### **ORDINARY SIMPLE MIXES**

- Nutrient and weed control granules of many different sizes and shapes
- Result: granules of unequal size can separate during application

#### [Subregistrant's] TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control granules are consistent in size and shape
- Result: consistent distribution of nutrients and weed control

[photos: simple mix and LF+CWC patterns]

#### **ORDINARY SIMPLE MIXES**

- The amount of individual nutrients and weed control in each square inch of lawn may vary greatly
- · Result: uneven, inconsistent weeding and feeding

#### [Subregistrant's] TECHNOLOGY

- Lawn Fertilizer Plus Confront Weed Control delivers the same ratio of primary nutrients plus weed control to every square inch of turf
- · Result: consistent greening plus better weed control

#### What to expect

Dandelions and other weeds will be killed in 4 to 5 weeks and your lawn will be greener in 4 to 6 days.

Small, lightweight granules mean better weed control. Most lawn weed controls must be absorbed through the weed's leaves to be effective. Lawn Fertilizer Plus Confront Weed Control granules are small and lightweight, so they will stay on even the smallest weed leaves. Weed control in Lawn Fertilizer Plus Confront Weed Control is absorbed by the weed's leaves and roots, killing the plant completely.

#### **Directions for Use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

Apply to moist or dry foliage at the REGULAR RATE to control the following weeds:

black medic

hop clover

bristly hawksbeard

narrow leaf plantain (buckhorn)

broadleaf plantain

red clover

catsear (coast dandelion)

white clover

common dandelion

false dandelion

hawkweed

Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

Apply to moist foliage at the HEAVY RATE to control the following weeds.

creeping beggarweed dwarf beggarweed

roundleaf mallow sheep sorrel shepherdspurse

burdock coffee weed

siender speedweil

common chickweed mouseear chickweed common cocklebur

smartweed spurweed Canada thistle

curly dock

musk thistle

dollarweed

old world diamond flower

English lawn daisy

common vetch Virginia pepperweed

goldenrod ground ivy

wild buckwheat

henbit lambsquarters common yellow woodsorrel creeping woodsorrel

lespedeza

prostrate spurge

matchweed

wild violet

poison ivy

Virginia buttonweed

common ragweed

A repeat application may be necessary.

#### How and when to apply

Apply in late spring, summer, or fall.

Application Restrictions: Avoid contact with eyes, skin, or clothing during and after application. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not enter or allow people or pets to enter the treated area until dusts have settled.

- Apply anytime weeds are actively growing (when daytime temperatures are consistently above 60 = F).
   Avoid applying when temperatures are consistently above 90 = F or during drought conditions.
   Applications during these conditions may provide less control than desirable.
- · Apply on a calm day.
- Lawn Fertilizer Plus Confront Weed Control is absorbed by both foliage and roots. Weeds listed in the REGULAR RATE section above can be controlled by application to dry foliage. For best results, apply when good moisture conditions are present.
- To control weeds listed in the HEAVY RATE section, apply to moist grass. To help particles stick to weed leaves, the grass should be thoroughly watered or wet from dew before applying Lawn Fertilizer Plus Confront Weed Control.
- In the Fall, apply up to the last mowing or as long as grasses are actively growing.
- On bahia and centipedegrass lawns, wait until after the second mowing of the year, but NOT before
  mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after
  the first of November.
- Do not use this product on flowers, shrubs, fruit, or vegetable plants, or on creeping bentgrass, St.
  Augustinegrass, dichondra, or lippia lawns. Do not apply this product to hybrid bermudagrass varieties such as Tifdwarf, Tifgreen, and Tifway. On common bermudagrass, bahia, centipedegrass, or zoysia lawns, under certain conditions some temporary discoloration may result.

#### Watering or rain

Do not water your lawn for 24 hours after application. Do not apply when rain is expected within 24 hours. Water will wash the weed control product off the weed leaves and reduce it's effectiveness.

## IMPORTANT What to Expect

Weed stems and leaves will begin to twist and curl within a week, but a month or more may be required for complete control. If weed control is incomplete after 4 to 5 weeks, a repeat application can be made. To minimize grass injury, additional applications should be at least 4 weeks apart.

#### Seeding/Sodding/Sprigging

Wait until the new grass has been mowed at least 4 times before applying Lawn Fertilizer Plus Confront Weed Control.

If you are planning to seed, sod, or sprig, you should wait at least 1 month after you have applied Lawn Fertilizer Plus Confront Weed Control.

This product will control only actively growing weeds; it will not prevent weed seeds from sprouting new plants.

#### **Spreading Tips**

For best results, use a [\_\_\_\_\_] spreader.

Be careful when applying this product near shrubs, flower beds, or vegetable gardens, as contact with this weed control product may damage these desirable plants. When using any rotary spreader, take special care near desirable plants.

Do not contaminate feed or foodstuffs.

Do not contaminate water used for irrigation or domestic purposes.

Do not graze treated areas. Do not feed clippings to livestock.

Do not apply in a brisk wind, as drift may contact desirable plants and cause injury.

Do not apply to exposed roots of shallow-rooted trees and shrubs or allow contact with exposed suckers, as injury may occur.

Do not reseed until 4 weeks after application.

Do not use Lawn Fertilizer Plus Confront Weed Control on golf course putting greens or tees.

Do not apply at greater than recommended rates, especially under the branch spread (dripline) of trees.

Do not use grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control for mulch.

Do not use compost containing grass clippings from turf treated with Lawn Fertilizer Plus Confront Weed Control in the growing season of application.

Spreader Settings					
Spreader	REGULAR RATE covers 5,000 sq ft	HEAVY RATE covers 4,300 sq ft	DIRECTIONS		
Scotts Drop Spreaders	XX	XX	Mow lawn first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.		
Scotts EasyGreen Spreaders	XX	XX	Make each pass about 30 inches from previous pass to overlap spread width.		
Scotts SpeedyGreen Rotary	XX	XX	Make each pass about 5 feet from previous pass to overlap spread width.		

	REGULAR	HEAVY
Additional Spreader Settings	RATE	RATE
EZ Drop, Republic (#530)	XX	XX
EZ Broadcast, Republic (#540)	XX	XX
EZ Grow, Republic (#545)	XX	XX
Red Devil Drop (FH-22)	XX	XX
Red Devil Rotary (SB-50)	XX	XX
Quaker Drop (Model #10850)	XX	XX
Quaker Rotary (Model #10301)	XX	XX
Ortho Drop	XX	XX
Ortho Rotary	XX	XX

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

[Editor's note: Illustrations; spreading patterns.]

#### Storage and Disposal

Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse empty bag. When completely empty, discard bag in trash.

Questions? Call [Subregistrant's] toll-free Consumer Hotline: 1-800-000-0000

#### [Subregistrant's] No-Quibble Guarantee™

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

[Subregistrant] . . . the lawn people [Subregistrant Address]

[Editor's note: Registrant is Dow AgroSciences LLC • Indianapolis, IN 46268-1054]



#### **Warranty Disclaimer**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

#### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

#### **Limitation of Remedies**

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

*Trademark of I	Dow	AgroSciences	LLC
EPA-Accepted:	_/_	_/	