

62719-267

6/20/2001

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

JUN 20 2001

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

John J. Jachetta, PhD  
Regulatory Manager  
Regulatory Success-Americas  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: Amendment - Change in PHI for Corn Grown for Seed  
Tracer (A.I. SPINOSAD)  
EPA Registration No. 62719-267  
Your Submission Dated March 21, 2001

Dear Dr. Jachetta:

The amendment referred to above, submitted in connection with registration under section 3 (c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable, subject to the comments listed below.

- 1 Under "Restrictions", add the following statements: "Do not apply within 28 days of grain or fodder harvest or within 7 days of forage harvest."

A stamped copy of the label is enclosed for your records. If you have any questions, please call Dr. William Sproat of my team at 703-308-8587.

Sincerely,

George T. LaRocca  
Product Manager 13  
Insecticide Branch  
Registration Division (7505C)

Enclosure

7505C  
SPROAT  
06/20/01

# Supplemental Labeling



2/2

Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA

## Tracer\*

EPA Reg. No. 62719-267

### For Insect Control in Corn Grown for Seed

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Tracer\* Naturalyte\* insect control before applying. Carefully follow precautionary statements and applicable use directions.
- Use of Tracer according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Tracer.

#### Directions for Use

##### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
European corn borer	1 - 3
armyworms (such as fall armyworm) corn earworm southwestern corn borer western bean cutworm	2 - 3

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
JUN 20 2001

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
62719-267

##### Specific Use Directions:

**Application Timing:** Scout for European corn borer and armyworms with enough regularity to monitor egg laying and egg hatch. Applications of Tracer should be timed to coincide with peak egg hatch of each generation. Frequent treatments may be necessary when crop is growing rapidly, during silking or under heavy pest pressure.

**Spray Delivery:** For control of first generation European corn borer, apply broadcast or as a directed spray into the leaf whorls. For control of second generation European corn borer, apply as a broadcast spray.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

**Chemigation:** Tracer may be applied to corn at recommended rates through sprinkler irrigation systems. Equipment should be calibrated to deliver the treatment in no more than 0.25 inch of water. Refer to the "Application by Chemigation" section in the main label for application guidelines for chemigation.

##### Restrictions:

- Do not apply more than 29 ounces (0.45 lb of spinosad) of Tracer per acre per year.
- Pre-harvest Interval: Do not apply within 1 day of harvest.

\*Trademark of Dow AgroSciences LLC

T6P / Tracer / Prop Sec 3 Suppl - Corn Grown for Seed / 03-05-00  
D06-066-XXX  
EPA Accepted: \_\_\_/\_\_\_/\_\_\_  
Initial printing.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....