

62719-267

1/15/2003

1/1

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Tracer*

EPA Reg. No. 62719-267

For Foliar Insect Control in Peanut

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Tracer* Naturalyte* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Tracer according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Tracer.

Directions for Use

Refer to product label for Tracer for General Use Precautions, Mixing and Application Instructions.

Pests and Application Rates:

| Pests | Rate (fl oz/acre) | Recommendations and Restrictions |
|--|-------------------|---|
| cabbage looper corn earworm European corn borer green cloverworm red-necked peanut worm saltmarsh caterpillar soybean looper velvetbean caterpillar | 1.5 - 3 | Rate Range: Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume. Use Restrictions: • Preharvest Interval: Do not apply within 3 days of nut harvest or 14 days of forage. • Do not apply more than a total of 9 fl oz of Tracer (0.282 lb ai of spinosad) per acre per crop or make more than 3 applications per calendar year. |
| Armyworms, including: beet armyworm fall armyworm true armyworm yellowstriped armyworm | 2 - 3 | • Minimum Treatment Interval: Do not make applications less than 7 days apart. • Grazing Restrictions: Do not allow grazing of crop residue or harvest of crop residue for hay until 14 days after the last application. |

Specific Use Directions:

Application Timing: Regularly monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Resistance Management: Do not apply Tracer more than 3 times in any 30 day period. Whenever Tracer is applied three times in succession, this should be followed by no use of Tracer for a 30 day period or rotation to another insecticide class.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and highly toxic to mollusks. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

T6P / Tracer / Sec 3 Supp Peanut / 10-11-02

D06-066-XXX

Accepted: ___/___/___

Initial printing.

ACCEPTED
JAN 15 2003

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide registered under EPA Reg. No. 62719-267.