



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

May 28, 2008

Kenneth D. Racke, Ph.D.
Regulatory Leader
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Amend previously approved supplemental labeling (add use site -sod farm-, and pests)
Tracer (EPA Reg. No. 62719-267)
Your Submission Dated February 27, 2008


Dear Dr. Racke:

We have reviewed the above referenced submission regarding registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended. The supplemental labeling, "Insect Control in Grass Forages, Grass Grown for Seed, Sod Farms, Pastures and Rangeland" is acceptable.

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate this supplemental labeling into the product's main labeling.

Enclosed, please find stamped copy of the label for your records. If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely,


George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure:

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Tracer[®]

EPA Reg. No. 62719-267

Insect Control in Grass Forages, Grass Grown for Seed, Sod Farms, Pastures and Rangeland

ATTENTION


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Tracer[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Tracer according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Tracer.

Directions for Use

Refer to product label for Tracer for General Use Precautions, Maximum Use Rates, Application Restrictions, Mixing and Application instructions.

Pests and Application Rates

| Target Pests | Rate (fl oz/acre) | Active Ingredient (lb/acre) |
|--|----------------------|--------------------------------|
| beet armyworm fall armyworm sod webworms southern armyworm true armyworm other lepidopterous species | 1 - 3 | 0.031 - 0.094 |



Application Timing

Scout at least weekly and consider the impact of both pests and beneficials. Treat when economic thresholds are exceeded, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications, but follow resistance management guidelines on the product label. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations applicable to your area.

Application Rate

Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

Resistance Management

For resistance management, do not apply Tracer more than 3 times in any 21-day period. Rotate to a different class of insect control products or use no treatment for the next 21 days. Do not apply more than 3 treatments per season. If Tracer is applied 3 times in succession, do not apply again for at least 21 days.

Specific Use Restrictions:

- **Preharvest Interval:** Do not harvest hay or fodder for 3 days after treatment. There is no preharvest interval for forage.
- Do not allow cattle to graze from treated area until spray has dried.
- Do not apply more than 6 fl oz of Tracer (0.186 lb ai spinosad) per acre per season.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R066-022

Accepted: __/__/__

Replaces R066-010.