



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 4, 2026

Terri Moss  
Regulatory Specialist  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: Label Amendment - Registration Review Mitigation for Spinosad  
Product Name: TRACER  
EPA Registration Number: 62719-267  
Case Number: 477178  
Application Dates: September 7, 2022 and January 22, 2026

Dear Terri Moss:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Spinosad Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at [carr.caleb@epa.gov](mailto:carr.caleb@epa.gov).

Sincerely,



Julie Javier, Team Leader  
Risk Mitigation and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

(Base label):

SPINOSAD	GROUP	5	INSECTICIDE
----------	-------	---	-------------

# Tracer®

## Naturalyte® Insect Control

**A Naturalyte® insect control product for control of lepidopterous larvae (worms or caterpillars), leafminers, and thrips**

### Active Ingredients:

spinosad (a mixture of spinosyn A and	
spinosyn D) .....	44.2%
Other Ingredients .....	55.8%
Total .....	100.0%

Contains 4 lb of active ingredient per gallon.  
U.S. Patent No. 5,362,634 and 5,496,931

**ACCEPTED**  
Feb 4, 2026  
Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 62719-267

### Keep Out of Reach of Children

### Precautionary Statements

#### Personal Protective Equipment (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### Environmental Hazards

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and reduce risk to these organisms. Apply this product only as specified on the label.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**(Storage and Disposal for rigid containers 5 gal or less)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**(Storage and Disposal for refillable rigid containers larger than 5 gal)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)****Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refer to label booklet for additional precautionary information and Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

**Shake Well Before Use -- Avoid Freezing**

EPA Reg. No. 62719-267

EPA Est. \_\_\_\_\_

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for:**

**Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**Net Contents** \_\_\_\_\_

(Label booklet cover):

SPINOSAD	GROUP	5	INSECTICIDE
----------	-------	---	-------------

# Tracer®

## Naturalyte® Insect Control

**A Naturalyte® insect control product for control of lepidopterous larvae (worms or caterpillars), leafminers, and thrips**

Active Ingredients:

spinosad (a mixture of spinosyn A and	
spinosyn D) .....	44.2%
Other Ingredients .....	55.8%
Total .....	100.0%

Contains 4 lb of active ingredient per gallon.

U.S. Patent No. 5,362,634 and 5,496,931

### Keep Out of Reach of Children

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information and Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

### Shake Well Before Use -- Avoid Freezing

EPA Reg. No. 62719-267

EPA Est. \_\_\_\_\_

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for:**

**Corteva Agriscience LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**Net Contents** \_\_\_\_\_

(Page 1 through end):

---

## Precautionary Statements

---

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

### Environmental Hazards

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and reduce risk to these organisms. Apply this product only as specified on the label.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Nonrefillable containers 5 gallons or less:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers 5 gallons or larger:**

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable containers 5 gallons or larger:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## General Information

Tracer® is a Naturalyte® insect control product for control of many foliage feeding pests infesting labeled crops. This product's active ingredient, spinosad, is biologically derived from the fermentation of *Saccharopolyspora spinosa*, a naturally occurring soil organism. Tracer should be mixed with water and applied as a foliar spray with aerial or ground equipment equipped for conventional insecticide spraying.

## General Use Precautions

### Integrated Pest Management (IPM) Programs

Tracer is recommended for IPM programs in labeled crops. Tracer should be applied when field scouting indicates target pest densities have reached the economic threshold, i.e., the point at which the insect population must be reduced to avoid economic losses beyond the cost of control. Other than reducing the target pest species as a food source, Tracer does not have a significant impact on certain parasitic insects or the natural predaceous arthropod complex in treated crops, including big-eyed bugs, ladybird beetles, flower bugs, lacewings, minute pirate bugs, damsel bugs, assassin bugs, predatory mites or spiders. The feeding activities of these beneficials will aid in natural control of other insects and reduce the likelihood of secondary pest outbreaks. If Tracer is tank mixed with any insecticide that reduces its selectivity in preserving beneficial predatory insects, the full benefit of Tracer in an IPM program may not be reduced.

### Insecticide Resistance Management (IRM)

For resistance management, Tracer contains spinosad, a Group 5 insecticide. Any insect population may contain individuals naturally resistant to Tracer and other Group 5 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay development of insecticide resistance, the following practices are recommended:

- Carefully follow the specific label guidelines within the use directions sections of this label, especially in regard to IRM recommendations.
- Rotate the use of Tracer or other Group 5 insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendation provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one of both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Applications should be targeted against early insect developmental stages whenever possible.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist, certified crop advisor, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

- For further information or to report suspected resistance, contact your local company representative or by calling 800-258-3033 or over internet at [www.dowagro.com](http://www.dowagro.com).

## Mixing

**Always shake well before use. Avoid freezing.**

### Application Rate Reference Table

Application Rate of Tracer (fl oz/acre)	Active Ingredient Equivalent (lb ai/acre)	Acres per Gallon of Tracer
1	0.031	128
1.4	0.045	90
1.5	0.047	85
2	0.062	64
2.14	0.067	60
2.5	0.078	51
2.9	0.089	45
3	0.094	43

**Mixing Tracer Alone:** Fill the spray tank with water to about one-half of the required spray volume. Start agitation and add the required amount of Tracer. Continue agitation while mixing and filling the spray tank to the required spray volume. Maintain sufficient agitation during application to ensure uniformity of the spray mix. Do not allow water or spray mixture to back-siphon into the water source.

**Tank Mixing:** When tank mixing Tracer with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. If foliar fertilizers are used, the jar test should be repeated with each batch of fertilizer utilizing the mixing water source. Do not use acidifying buffering agents in tank mixes with Tracer. Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

**Mixing Order for Tank Mixes:** Fill the spray tank with water to 1/4 to 1/3 of the required spray volume. Start agitation. Add different formulation types in the order indicated below, allowing time for complete dispersion and mixing after addition of each product. Allow extra dispersion and mixing time for dry flowable products.

Add different formulation types in the following order:

1. Products in water soluble packaging
2. Water dispersible granules
3. Wettable powders
4. Tracer and other aqueous suspensions

Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add:

5. Emulsifiable concentrates and water-based solutions
6. Spray adjuvants, surfactants and oils
7. Foliar fertilizers

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose.

**Premixing:** Dry and flowable formulations may be premixed with water (slurried) and added to the spray tank through a 20 to 35 mesh screen. This procedure assures good initial dispersion of these formulation types.

**Spray Tank pH:** A spray tank pH between 6.0 and 9.0 is suggested to achieve maximum performance of Tracer. If the water source is outside of this pH range, or tank mixing other pesticides, adjuvants, or foliar nutrients will cause the pH to fall outside this range, consider adjusting the spray tank pH to be between 6.0 and 9.0 before adding Tracer. To do this, add all other tank mix components first, then check the spray tank pH, adjust if desired, and then add Tracer. If you require additional information on how to adjust spray tank pH, contact your company representative.

**Use of Adjuvants:** Adjuvants may be used to improve the control of leafminers and thrips in situations where achieving uniform plant coverage is difficult (such as closed crop canopy or dense foliage), or penetration into waxy leaf surfaces is required.

- Use only adjuvant products labeled for agricultural use and follow the manufacturer's label directions. A nominal concentration of 1 to 2 quarts per 100 gallons (0.25 to 0.5% v/v) is generally sufficient.
- For leafminers and thrips, emulsified crop oils or methylated crop oil plus organosilicone combination products are recommended.
- When using adjuvants, always conduct a jar test to determine the compatibility of the various components in the mixture. Crop safety should be determined in a small area of the crop whenever there is a significant change in spray mixture ingredients or source of water for the spray mixture.
- Do not use diesel fuel or pure mineral oil.

## Application

Proper application techniques help ensure thorough spray coverage and correct dosage for optimum insect control. The following recommendations are provided for ground and aerial application of Tracer. Attention should be given to sprayer speed and calibration, wind speed, and foliar canopy to ensure adequate spray coverage.

## SPRAY DRIFT

### Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative Canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed-wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Applicators must use  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

### Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 572 (ASAE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if application are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented with the airflow in flight.

### BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

**RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**TEMPERATURE AND HUMIDITY**

When making application in hot and dry conditions, use larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

**WIND**

Drift potential generally increased with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

**Band Application:** Band application may be appropriate when the crop is small. Nozzle selection, placement, or shielding to compensate for windy conditions is critical to ensure adequate coverage.

**Chemigation Application**

Tracer may be applied through properly equipped chemigation systems for insect control in corn. Follow use directions for these crops in the Uses section of this label. Do not apply Tracer by chemigation to other labeled crops except as specified in Corteva Agriscience supplemental labeling or product bulletins.

**General Directions for Chemigation:**

Tracer may be applied through drip or overhead sprinkler irrigation systems that will apply water uniformly, including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system. Sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units are not recommended.

For continuously moving systems, the mixture containing Tracer must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. If continuously moving irrigation equipment is used, apply in no more than 0.25 inch of water. For irrigation systems that do not move during operation, apply in no more than 0.25 inch of irrigation immediately before the end of the irrigation cycle.

**Preparation:** The following use directions are to be followed when this product is applied through irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of Tracer needed to cover the desired acreage. Mix according to instructions in the Mixing section above. Continually agitate the mixture during mixing and application.

**Equipment Calibration:** In order to calibrate the irrigation system and injector to apply the mixture containing Tracer, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Calculate the amount of product required and premix; 3) Determine the irrigation rate and the number of minutes for the system to cover the intended treatment area; 4) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes (minus time to flush out) to cover the treatment area. This value equals the gallons per minute output that the injector or eductor must deliver. Convert the gallons per minute to milliliters or ounces per minute, if needed. Calibrate the injector system with the system in operation at the desired irrigation rate. It is suggested that the injection pump/system be calibrated at least twice before operation, and the system should be monitored during operation.

**Operation:** Start the water pump and irrigation system, and let the system achieve the desired pressure and speed before starting the injector. Check for leaks and uniformity and make repairs before any chemigation takes place. Start the injection system and calibrate according to manufacturer's recommendations. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

**Precautions:**

- Lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers or other experts.
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply and must be followed.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate nontarget areas.
- Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.
- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn.
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

**Specific Equipment Requirements:**

- The system must contain an air gap or approved backflow prevention device, or approved functional check valve, vacuum relief valve (including inspection port), and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations.
- The pesticide injection line must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection chemical supply.
- A pesticide injection pipe must also contain a functional interlock, e.g., mechanical or electrical, to shut off chemical supply when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection when the water pressure drops too low or water flow stops.
- Use of public water supply requires approval of a backflow prevention device or air gap (preferred) by both state and local authorities.
- Systems must use a metering pump, such as a positive displacement injection pump (or flow meter on eductor) effectively designed and constructed of materials that are compatible with pesticides and

capable of being fitted with a system interlock. An electric powered pump must meet Section 675 for "Electrically Driven or Controlled Irrigation Machines" NEC 70.

- To insure uniform mixing of the insecticide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.
- The tank holding the insecticide mixture should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injection point.

## Uses

### Corn (Field Corn, Popcorn, and Corn Grown for Seed) and Teosinte

#### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
European corn borer fall armyworm true armyworm	1 - 3
beet armyworm corn earworm southwestern corn borer western bean cutworm	2 - 3

#### Specific Use Directions:

**Application Timing:** Scout for **European corn borer** and **armyworms** with enough regularity to monitor egg laying and egg hatch. Applications of Tracer should be timed to coincide with peak egg hatch of each generation. Frequent treatments may be necessary when the crop is growing rapidly, during silking or under heavy pest pressure. For **corn earworm** control, a 1- to 2-day re-treatment schedule may be necessary at silking. For control of all other pests, a 5- to 7-day re-treatment schedule may be necessary if the crop is growing rapidly or if there is heavy pest pressure.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

**Spray Volume:** For control of first generation **European corn borer** and **armyworms**, apply broadcast or as a directed spray into the leaf whorls. For control of second generation **European corn borer**, apply as a broadcast spray. For control of **corn earworm**, apply broadcast or direct spray to ear zone. Use sufficient spray volume and nozzle pressure to ensure thorough wetting of the silks.

**Chemigation:** Tracer may be applied to corn by chemigation at labeled rates. Refer to the Chemigation Application section for application guidelines for chemigation.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

#### Restrictions:

- Do not apply more than a total of 6 fl oz of Tracer (0.188 lb ai spinosad) per acre per year.

#### Popcorn and Corn Grown for Seed

- **Preharvest Interval:** Do not apply within 1 day of grain harvest or within 3 days of forage or fodder harvest.

### Field Corn and Teosinte

- **Preharvest Interval:** Do not apply within 28 days of grain harvest or within 3 days of fodder or forage harvest.

## Cotton

### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
cotton bollworm (prebloom)	1.4 - 2.9
cotton leafperforator	
European corn borer	
tobacco budworm	
armyworms (including beet armyworm, fall armyworm)	2.14 - 2.9
cotton bollworm (post-bloom)	
leafminers	
loopers (including soybean looper, cabbage looper)	
saltmarsh caterpillar	
thrips	

### Specific Use Directions:

#### Application Timing:

**Tobacco Budworm and/or Cotton Bollworm:** For the most effective control, fields should be scouted twice per week and Tracer applied when the majority of the population is within the time of blackhead egg stage to 1/8-inch larval length. The following table illustrates the size of development of worms in relation to age and stage of development (instar) as a guide to timing treatments for optimum control:

Age (Days)	Average Size (inches)	Instar <sup>1</sup>
Hatch	1/16	1st
3	1/4	2nd
5	1/2	3rd
8	7/8	4th
10	1	5th

<sup>1</sup>**Note:** A scouting schedule of only once per week is risky since hatching worms will have grown to 3rd instar before the next scouting observation has determined the need to spray.

Heavy infestations may require repeat applications but make no more than 3 applications per 30 days or 6 applications per crop. The minimal time period between applications must be 7 days. To delay resistance selection, do not treat sequential generations with the same class of insect control products. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your company representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional area use recommendations for your area.

**Beet Armyworm:** Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply Tracer when field scouting reveals 3 or more occurrences of egg hatch or larval feeding per 100 feet of row.

**Loopers:** Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply Tracer when field scouting reveals 4 larvae per 1 foot of row or 25% defoliation.

**Application Rate:** Use a higher rate within the rate range and higher spray volume when one or more of the following is true: tobacco budworms or bollworms are more than 1/4 inch in length; target pest population is 2X above stated threshold level; or foliage canopy is tall/dense and worms are present in the lower part of the canopy. Heavy infestations may require repeat applications, but follow resistance management guidelines.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. For **tobacco budworm and/or cotton bollworm** where early season conservation of beneficial insects is practical, use Tracer to control the 1st and 3rd generation of tobacco budworm and/or cotton bollworm. Where conservation of beneficial insects is not as critical (for example, fields have received non-selective early season treatments for boll weevil or lygus bugs), use Tracer to control either the 2nd or 3rd generation of tobacco budworm and/or cotton bollworm.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 5 days apart for high rates of application.
- Do not apply more than a total of 14.4 fl oz of Tracer (0.45 lb ai spinosad) per acre per growing season.

## Grass Forages, Grass Grown for Seed, Pastures, Rangeland and Sod Farms

### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
beet armyworm	1 – 2
fall armyworm	(0.031 – 0.062 lb ai/acre)
sod webworms	
southern armyworm	
true armyworm	
other lepidopterous species	

### Specific Use Directions:

**Application Timing:** Scout at least weekly and consider the impact of both pests and beneficials. Treat when economic thresholds are exceeded, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications, but follow resistance management guidelines on the product label. Consult your company representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations applicable to your area.

**Application Rate:** Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

**Resistance Management:** Do not apply more than 3 times in any 21-day period. Whenever Tracer is applied up to 3 times in succession, this should be followed by no use of Tracer for a 21-day period or rotation to another insecticide class. Do not make more than 6 applications per season.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 3 days of harvest for hay or fodder. There is no preharvest interval for forage.
- Do not apply more than a total of 6 fl oz of Tracer (0.188 lb ai spinosad) per acre per season.
- Do not allow cattle to graze from treated area until spray has dried.

## Peanut

### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
cabbage looper	1.5 - 3
corn earworm	
European corn borer	
green cloverworm	
red-necked peanut worm	
saltmarsh caterpillar	
soybean looper	
tobacco budworm	
velvetbean caterpillar	
armyworms, including:	2 - 3
beet armyworm	
fall armyworm	
true armyworm	
yellowstriped armyworm	

### Specific Use Directions:

**Application Timing:** Regularly monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your company representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

### Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of nut harvest or within 14 days of forage.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 9 fl oz of Tracer (0.28 lb ai spinosad) per acre per crop or make more than 3 applications per calendar year.
- Do not allow grazing of crop residue or harvest of crop residue for hay until 14 days after the last application.

## Small Cereal Grains

Small cereal grains (crop group 15) including barley, buckwheat, oats, rye, triticale, wheat

### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
cereal leaf beetle	1 - 3
armyworms	1.5 - 3
corn earworm (headworm)	
grasshoppers (suppression)	
southwestern corn borer	
webworms	

### Specific Use Directions:

**Application Timing:** Scout for **armyworms** and **headworms** with enough regularity to monitor egg laying and egg hatch and treat when thresholds are reached. Applications of Tracer perform best when timed to coincide with peak egg hatch of each generation.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in the rate range for heavy infestations and/or difficult spray coverage situations.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

### Restrictions:

- **Preharvest Interval:** Do not apply within 21 days of grain or straw harvest or within 3 days of forage, fodder or hay harvest.
- Do not apply more than a total of 9 fl oz of Tracer (0.28 lb ai spinosad) per acre per year.
- Do not allow cattle to graze treated area until spray has dried.

## Sorghum, Grain Amaranth, Milo, Pearl Millet, and Proso Millet

### Pests and Application Rates:

Pests	Tracer (fl oz/acre)
armyworms	1.5 - 3
corn earworm (headworm)	
southwestern corn borer	
webworms	

### Specific Use Directions:

**Application Timing:** Scout for **armyworms** and **headworms** with enough regularity to monitor egg laying and egg hatch and treat when thresholds are reached. Applications of Tracer perform best when timed to coincide with peak egg hatch of each generation.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in the rate range for heavy infestations and/or difficult spray coverage situations.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult

your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 7 days of grain or fodder harvest or within 14 days of forage harvest.
- Do not apply more than a total of 14.4 fl oz of Tracer (0.45 lb ai spinosad) per acre per year.
- Do not allow cattle to graze treated area until spray has dried.

## Soybean

**Pests and Application Rates:**

Pests	Tracer (fl oz/acre)
green clover worm	1 - 2
soybean looper	
true armyworm	
velvet bean caterpillar	
armyworms (such as fall armyworm, yellowstriped armyworm, beet armyworm)	1.5 - 2
corn earworm (podworm)	
saltmarsh caterpillar	

**Specific Use Directions:**

**Application Timing:** Treat when field counts or crop injury indicates damaging pest populations are present or developing. Time applications to treat small larvae and use sufficient spray volume to ensure good coverage.

**Application Rate:** Use a higher rate in the rate range for heavy infestations and/or difficult spray coverage situations.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- Do not apply more than a total of 6 fl oz of Tracer (0.188 lb ai spinosad) per acre per year.
- Do not feed treated forage or hay to meat or dairy animals.

## Tobacco

**Pests and Application Rates:**

Pests	Tracer (fl oz/acre)
thrips	1.4 – 2.9
tobacco budworm	
tobacco hornworm	

**Specific Use Directions:**

**Application Timing:** Scout for lepidopterous larvae with enough regularity to monitor egg laying and egg hatch and treat when thresholds are reached. Applications of Tracer perform best when timed to coincide with peak egg hatch of each generation.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in the rate range for heavy infestations and/or difficult spray coverage situations.

**Spray Volume:** Use a minimum of 20 gallons of water per acre to obtain full coverage of foliage, increasing volume and nozzles per row as necessary with crop maturity.

**Resistance Management:** Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local company representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.4 fl oz of Tracer (0.45 lb ai spinosad) per acre per year or make more than 3 applications per 30 days or 6 applications per crop.

---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

**Warranty Disclaimer**

---

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks use shall be assumed by buyer and/or user.

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™®Trademarks of Corteva Agriscience and its affiliated companies

EPA accepted \_\_\_/\_\_\_/\_\_\_