

PM 25

62719-275

12-22-97

10/16

Please read instructions on reverse before completing form.

Form Approved OMB No. 2970-0060. Approval expires 05-31-98



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

253281

Application for Pesticide - Section I

1. Company/Product Number DowElanco/62719- 275 275	2. EPA Product Manager James Thompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DowElanco/ Frontrow	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) DowElanco 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review in accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in Composition and labeling to: EPA Reg. No. Product Name	

Section - II

NOTIFICATION

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed label in response to Agency letter dated
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

DEC 22 1997

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)
Notification of main labeling change per PR Notice 95-2.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Chlor-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
*Certification must be submitted		If "Yes" Unit Packaging wgt. container	No. per container	If "Yes" Package wgt	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Other _____		
			Lithograph Paper glued Stenciled		

Section - IV

1. Contact Point /Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name John J. Jachetta, Ph.D.	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4672
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowing false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		8. Data Application Received (\$tamped)
2. Signature 	3. Title Regulatory Manager	
4. Typed Name John J. Jachetta, Ph.D.	5. Date December 18, 1997	

P4D/PropSec3/12-16-97
file: FrontrowSec3 Dec97d.doc

Frontrow*
EPA Reg. No. 62719-XXX

A Co-pack Product For Postemergence Broadleaf Weed Control In Soybeans.


FirstRate*
EPA Reg. No. 62719-275

Flumetsulam 80 WDG
EPA Reg. No. 62719-277

Registration Notes:

Proposed labeling for Section 3 registration.

Herbicide
*Trademark of DowElanco



Keep Out of Reach of Children**CAUTION PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation • Harmful If Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Flumetsulam has been identified in groundwater sampling from a field research site under vulnerable conditions. There is the possibility that flumetsulam may leach through soil to groundwater, especially, where soils are coarse and groundwater is near the surface.

This chemical and its transformation products demonstrate the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

This chemical can contaminate surface water through spray drift.

Under some conditions, this chemical, and/or its transformation products, may have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several weeks post-application. Vulnerable conditions include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

60716

(Label Booklet cover)

(Logo) DowElanco

Frontrow*

**A Co-pack Product For Postemergence Broadleaf Weed Control In Soybeans
Containing FirstRate and Flumetsulam 80 WDG Herbicides.**

EPA Reg. No. 62719-XXX

FirstRate*

Active Ingredient:

cloransulam-methyl: N-(2-carbomethoxy-6-chlorophenyl)-5-ethoxy-7-fluoro(1,2,4)triazolo-[1,5-c]pyrimidine-2-sulfonamide84%

Inert Ingredients16%

Total100%

Contains 0.84 pounds of active ingredient per pound of product.

US Patent Number: 5,163,995

EPA Reg. No. 62719-275

Net Wt - 1, 1.5 oz Water Soluble Packet

Flumetsulam 80 WDG

Active Ingredient:

flumetsulam: N-(2,6-difluorophenyl)-5-methyl-1,2,4-triazolo-[1,5a]-pyrimidine-2-sulfonamide80.0%

Inert ingredients20.0%

Total100.0%

Contains 0.8 pounds of active ingredient per pound of product.

US Patent Numbers: 4,554,163 and 4,818,273

EPA Reg. No. 62719-277

Net Wt - 1, .6 oz Water Soluble Packet

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE) and Directions for Use including Storage and Disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" elsewhere on this label.

70416

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.


EPA Reg. No. 62719-XXX

EPA Est. _____

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268 USA

Herbicide

Net Weight _____



80116

(page one through end)

Precautionary Statements

Hazards to Humans and Domestic Animals**Causes Eye Irritation • Harmful if Absorbed Through Skin****Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.****Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Flumetsulam has been identified in groundwater sampling from a field research site under vulnerable conditions. There is the possibility that flumetsulam may leach through soil to groundwater, especially, where soils are coarse and groundwater is near the surface.

This chemical and its transformation products demonstrate the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

This chemical can contaminate surface water through spray drift.

Under some conditions, this chemical, and/or its transformation products, may have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several weeks post-application. Vulnerable conditions include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

90416

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Disposal: Wastes resulting from the use of this product may be disposed of on site according to label use directions or at an approved waste disposal facility.

Container Disposal: Dispose of empty package or overpacks in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

Frontrow® herbicide is a co-pack product, for postemergence broadleaf weed control in soybeans. Each co-pack (overpack) of Frontrow contains 1.5 oz of FirstRate® herbicide and .6 oz of Flumetsulam 80 WDG herbicide in separate water soluble packets. Frontrow is formulated as water dispersible granules. The individual components of the Frontrow co-pack must be used together or a reduction in weed control will result. Do not mix and apply separately.

When applications are made under adverse (dry or cold) conditions or when large weeds or less susceptible species are treated, weed suppression may be observed. Weed suppression is a visual reduction in weed competition (reduced population, size, and/or vigor) as compared to an untreated area. Degree of control can be increased by applying Frontrow under good growing conditions (i.e., adequate moisture and temperature).

10/2/16

Safe Pesticide Handling Procedures

Handling Precautions for Product Packaged in Water Soluble Packets: Do not remove water soluble packet from overpack except for immediate use. Do not allow water soluble packet to come into contact with water prior to use. Do not handle water soluble packet with wet hands or wet gloves. Carefully reseal overpack containing unopened water soluble packets and protect package from moisture.

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid over filling the spray tank.
- Do not discharge excess material on soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess spray solution and apply on approved crop site.

• Sprayer Cleanup:

To avoid injury or exposure to nontarget crops, thoroughly clean all mixing and spray equipment on approved rinse pads or on the field site where an approved crop is to be grown. Do this immediately after application and prior to use on crops other than those registered for use on this label. Spray equipment should be cleaned at the end of each day by the following procedure:

1. Drain any remaining Frontrow from the spray tank and dispose of according to label disposal instructions.
2. Hose down the interior surfaces of the tank and fill the tank 1/2 full with water. Recirculate for 5 minutes. Spray part of the mixture through the hoses, boom and nozzles and drain the tank. All rinse material should be collected in a suitable container and disposed of according to label directions (applied to cropland to be planted to crops recommended in this label) or in a locally approved manner (see "Storage and Disposal" section).
3. Remove the nozzles and screens and clean separately.
4. If the spray equipment will be used on crops other than those registered for use on this label, repeat steps 1 and 2 again and thoroughly wash the spray mixture from the outside of spray tank and the boom.

Placement of mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

General Use Precautions

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involved mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the

11/2/16

largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Do not apply this product through any type of irrigation system.

Do not use flood irrigation to apply or incorporate this product.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

- Make only one application per crop growing season.
- Do not apply more than 0.42 oz per acre of Frontrow (0.3 oz FirstRate + 0.12 oz Flumetsulam 80 WDG).
- Do not graze or feed treated soybean forage, hay or straw to livestock.
- Do not apply when air temperature is near freezing or when freezing conditions are expected for several days following application.
- Do not aerially apply Frontrow. Apply using ground equipment only.
- Avoid all direct or indirect contact with non-target plants. Do not apply near desirable vegetation and allow adequate distance between target area and desirable plants.
- Do not apply when weather conditions favor drift to non target sites. To minimize spray drift to non target areas:
 - Use low pressure application equipment capable of producing a large droplet spray.
 - Do not use nozzles that produce a fine-droplet spray.
 - Minimize drift by using sufficient spray volume to ensure adequate coverage with large droplet size sprays.
 - Keep ground-driven spray boom as low as possible above the target surface.
 - Spray when conditions are calm or wind speed is low. Do not spray when wind is gusting or steady wind speed is greater than 10 mph.
- Follow all applicable directions, precautions and restrictions on labeling for other products used in combination with Frontrow.

Rotational Crop Restrictions:

- When tank mixing with other herbicides, follow crop rotation guidelines on the label of each product used.
- The following rotational crops may be planted at the indicated interval following application of Frontrow.

Crop	Rotation Interval ¹ (Months)
Small grains	4 1/2
Corn, Sorghum, Cotton, Peanuts	9
Sugar beets, Sunflowers, Tobacco	26 ¹¹

¹Minimum number of months that must pass before planting other crops after application of Frontrow.

¹¹Note: Rotation to sugar beets, sunflowers, tobacco, canola and all other crops requires a 26 month rotation interval and a successful field blossomy.

20/16

Field Bloassay Instructions: Using typical tillage, seeding practices, and timings for the particular crop, plant several strips of the desired crop variety across the field previously treated with Frontrow. Plant the strips perpendicular to the direction Frontrow was applied. The strips should also be located so that different field conditions are encountered, including differences in soil texture, pH, and drainage. If the crop does not show visible symptoms of injury, stand reduction, or yield reduction, the field can be seeded with the test crop in the growing season following the bioassay. If visible injury, stand reduction, or yield reduction occurs, the test crop should not be seeded, and the bioassay must be repeated the next growing season.

Mixing and Application

Spray Volume and Application: Apply in a total spray volume of 10 to 20 gallons per acre. Use sufficient spray volume to provide uniform coverage. Sufficient agitation should be maintained during mixing and spraying to ensure a uniform spray mixture. Apply with ground equipment using standard low pressure (20 to 40 pounds per square inch) herbicide sprayers equipped with flat fan nozzles. Do not apply using liquid fertilizer as carrier.

Mixing - Frontrow Alone

Water Soluble Packaging: Each Frontrow co-pack (overpack) contains two water soluble packets which must be removed from the overpack and added to the spray tank at the same time. Refer to Application Rates below to determine the number of co-pack units (each containing two water soluble packets) that must be added to the spray tank for the acreage to be treated. Thorough mixing is required.

Mixing Instructions:

1. Fill the tank with 1/2 of the total amount of water or liquid fertilizer required.
2. Start agitation system.
3. Add any adjuvant materials, and then add the required number of co-pack units (each containing two water soluble packets) to the spray tank.

Note: Handling water soluble packets with hands should be minimized. Do not handle if hands or gloves are wet. Water soluble packets must be removed from the overpack before adding to the spray tank. This can be done by opening the overpack and adding the two water soluble packets (product in transparent film) directly to the spray tank. Do not open water soluble packets. Water soluble packets will float on the surface until the soluble film dissolves and the product is released. Overpacks are not water soluble and should not be placed in spray tank.

4. Continue agitation and complete filling the tank while the packets dissolve.
5. Before spraying, make sure packets have completely disintegrated and product is thoroughly mixed with water. Depending on the water temperature and the degree of agitation, the packet and its contents should be completely dispersed within 5 minutes from the time they were added to the spray tank.

Screens in spray tank lines should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

To insure a uniform spray mixture continuous agitation is required during mixing and spraying. Apply within 24 hours after mixing. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying.

Pre-mixing (other products): If pre-mixing is required for other dry or flowable products applied in tank mix combination with Frontrow, follow recommendations for pre-mixing of such products provided in their respective product labels.

130416

Mixing - Frontrow In Tank Mix Combination

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture. To prevent foaming during filling, keep end of fill pipe below the surface of the liquid in the spray tank.

Compatibility Testing: When tank mixing Frontrow with other products, a compatibility test (jar test) using relative proportions of tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. Use a compatibility agent if the need for one is indicated by the jar test (ingredients separate readily and are not easily redispersed).

Mixing Order for Tank Mixes:

1. Fill the spray tank to 1/4 to 1/3 of the total spray volume required with water.
2. Start agitation system.
3. Add different formulation types in the following order: (1) Compatibility agent, if required; (2) Frontrow and other dry flowables; (3) wettable powders; (4) aqueous suspensions, flowables and liquids. Maintain agitation and fill spray tank to 3/4 of total spray volume and add: (5) emulsifiable concentrates and (6) solutions. Allow time for complete mixing and dispersion after addition of each product.
4. Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application.

If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Application Rates

Each co-pack (overpack) of Frontrow, containing 1.5 oz of FirstRate and .6 oz of Flumetsulam 80 WDG, will treat 10 acres. The following table provides per acre rates of Frontrow when applied according to label directions. Note: Open the overpack and add both of the water soluble packets directly to the spray tank. The overpack is NOT water soluble.

Application Rates Per Acre:

One Co-pack of Frontrow contains: FirstRate (3.0 oz packet) and Flumetsulam 80 WDG (1.2 oz packet)	FirstRate* (oz/acre)	Flumetsulam 80 WDG (oz/acre)
One overpack treats 10 acres	1.5	.6

Rate Formula: Acres to be treated + # of acres per co-pack = Number of co-packs
One co-pack will treat 10 acres

Sample Calculation: Acres to treat = 27
27 acres + 10 acres per co-packs = 2.7 co-packs

When the number of overpacks calculated does not equal a whole number, round to the nearest whole number of overpacks.

140/16

Approved Uses

SOYBEANS**Postemergence Application**

Apply Frontrow at a rate of 1 co-pack (containing 1.5 oz of FirstRate and .6 oz of Fiumetsulam 80 WDG) per 10 acres. Apply as a broadcast spray when weeds are actively growing and before weeds exceed the growth stages (number of true leaves per plant) indicated in the table below. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control. Application timing generally corresponds to soybean growth stages of unifoliate to three to four expanded trifoliate leaves. Frontrow may be applied alone or in tank mix combination with other postemergence herbicides. Applications of Frontrow must include either a crop oil concentrate, a nonionic surfactant or a nonionic surfactant plus urea ammonium nitrate.

When applied as directed Frontrow will control the following weeds:

Weeds Controlled	Application Stage of Growth (Number of true leaves per plant)
cocklebur	2-4
jimsonweed	2-4
prickly sida	1-2
momingglory (annual) entireleaf ivyleaf palmleaf pitted red smallflower tall	2-4
ragweed common giant	2-4
sicklepod ¹	cotyledon - 1 (< 2 inches)
velvetleaf	2-4
venice mallow	2-4
wild sunflower	2-4

¹Reduced control will result if applications are made sicklepod at 2 leaf stage or larger. Additional herbicide treatment may be required to control sicklepod which germinates after application.

150416

When applied as directed Frontrow will suppress the following weeds:

Weeds Controlled	Application Stage of Growth (Number of true leaves per plant)
hemp sesbania	Cotyledon - 1
pigweed species redroot smooth spiny	1-2

Spray Adjuvants

Either a crop oil concentrate, a nonionic surfactant or a nonionic surfactant plus ammonium nitrate solution must be included in the spray solution.

Crop Oil Concentrate

Under hot, dry conditions, a crop oil concentrate may be used in place of the nonionic surfactant plus ammonium nitrate solution to enhance weed control. Crop oil concentrate is especially helpful in controlling sicklepod and prickly sida. Apply the crop oil concentrate at 1.2% v/v (volume/volume) (4 3/4 qt per 100 gal of spray solution). Use a good-quality, petroleum-based or methylated seed oil-based crop oil concentrate with at least 14% emulsifiers and 80% oil.

Note: Crop oil concentrate may increase the potential for crop injury in soybeans.

Nonionic Surfactant or Nonionic Surfactant Plus Ammonium Nitrate Solution

Use the nonionic surfactant in the spray solution at a rate (concentration) of 0.25% v/v (volume/volume) (1 qt per 100 gal of spray solution). Use only products that contain at least 80% nonionic surfactant as the active ingredient. When used in combination with a urea ammonium nitrate solution use the nonionic surfactant in the spray solution at a rate (concentration) of 0.125% v/v (1 pt per 100 gal of spray solution) and the urea ammonium nitrate solution at a concentration of 2.5% v/v.

Tank Mix Applications

When applied in tank mix combination with other herbicides, follow applicable use instructions, including application rates, precautions and restrictions for each product used in the tank mixture, including use of adjuvants.

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including adverse conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by the buyer.

16 of 16

Limitation of Remedies

The exclusive remedy for losses or damages resulting from the use of this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of DowElanco

EPA-accepted / /