

62719-275

06/20/2012

13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington D C 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

JUN 20 2012

Cindy Loy  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis IN 46268 1054

Subject Supplemental Label Increased Postemergence Application Rate in Soybean  
Firstrate  
EPA Reg No 62719 275  
Date Submitted June 12 2012

Dear Ms Loy

The supplemental labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act as amended is acceptable

This supplemental label expires on June 12 2015 and must not be used or distributed after that date Within eighteen (18) months of the date of this letter you must incorporate this supplemental labeling into the main product labeling A stamped copy of your label is enclosed for your records If you have any questions please contact Maggie Rudick at (703) 347 0257 or [rudick.maggie@epa.gov](mailto:rudick.maggie@epa.gov)

Sincerely

Kable Bo Davis  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

M2A / FirstRate / Sec 3 Soybean Amend / 06 11 12  
file R034 004 FirstRate Sec 3 Soybean 11Jun12d doc

**FirstRate<sup>®</sup>**

EPA Reg No 62719 275


**Registration Notes**

**Source Section 3 supplemental label** based on EPA accepted text dated August 8 2008

**Proposed changes by amendment**

- 1 Added Michigan and Ohio to list of states in which the product may be used
- 2 Added This supplemental labeling expires on June 12 2015 and must not be used or distributed after that date

<sup>®</sup>Trademark of Dow AgroSciences LLC



3/3

ACCEPTED  
JUN 20 2012  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 62719-275

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## FirstRate®

EPA Reg No 62719 275

(For Distribution and Use Only in the States of Alabama Arkansas Connecticut Delaware Georgia Florida Kentucky Louisiana Maine Maryland Michigan Mississippi Missouri (Bootheel) New Jersey New York North Carolina Ohio Oklahoma Pennsylvania South Carolina Tennessee Texas Virginia)

This supplemental labeling expires on June 12 2015 and must not be used or distributed after that date

### Increased Postemergence Application Rate in Soybean

**ATTENTION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling  
This labeling must be in the possession of the user at the time of application  
Read the label affixed to the container for FirstRate<sup>®</sup> herbicide before applying Carefully follow all precautionary statements and applicable use directions  
Use of FirstRate according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for FirstRate

### Directions for Use

Refer to the main product label for FirstRate for Use Precautions and Restrictions and Mixing Instructions

Use FirstRate as a postemergence broadcast spray at a rate of 0.3 oz per acre (see container label for a list of weeds controlled) A second application of up to 0.3 oz of FirstRate per acre may be applied to later germinating weeds For especially heavy weed infestations or added residual control FirstRate may be applied at up to 0.6 oz per acre Applications of 0.3 to 0.6 oz per acre may be made prior to maximum leaf stage or height Applications can be made alone or in combination with other labeled products such as glyphosate Refer to labels for additional instructions and recommendations pertaining to tank mixes

If sicklepod is the primary target weed apply 0.3 oz of FirstRate per acre when sicklepod is in the cotyledon to 1 leaf stage Make a second application of 0.3 oz of FirstRate per acre 10 to 14 days after the first application

### Restrictions

**Preharvest Interval** When more than 0.3 oz of FirstRate per acre is applied postemergence the preharvest interval for soybeans is 70 days and the preharvest interval for forage or hay is 25 days  
**Maximum Use Rates** Do not exceed 1.05 oz per acre in a single growing season If a soil application is made no more than 0.3 oz per acre may be applied postemergence When more than one postemergence application is made do not exceed a total of 0.6 oz per acre

® Trademark of Dow AgroSciences LLC

R034 004  
Accepted \_\_\_/\_\_\_/\_\_\_  
Replaces R034 003