

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Python* WDG

EPA Reg. No. 62719-277

Postemergence Control of Teaweed (Prickly Sida) in Soybeans

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Python WDG before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as described in this supplemental labeling, use of Python WDG is subject to all precautions and limitations imposed by the label affixed to the container for Python WDG.

Directions for Use

Python* WDG herbicide may be applied postemergence at 0.125 oz/acre to control prickly sida in soybeans.

If a postemergence application of Python WDG is made following a previous preemergence application of Python WDG herbicide, check to make sure that the cumulative rate of 0.07 pounds per acre of flumetsulam per year is not exceeded. One ounce of Python WDG contains 0.050 lb of flumetsulam. A postemergence application of Python WDG at 0.125 oz per acre contains 0.00625 lb of flumetsulam."

Note: Do not use liquid fertilizer as total carrier for postemergence application.

Weed Control Advisory

Apply to actively growing weeds. Unfavorable conditions such as drought, or near freezing temperatures before, at or following application may result in reduced weed control. Degree of control will depend on application rate used, weed susceptibility, and spray adjuvant as well as growing conditions at the time of treatment.

Cultivation: For best results, do not cultivate within 10 days before or after application.

Application Timing: Applications may be made to soybeans from the first to the fifth trifoliolate leaf stage of growth. Application should be made when teaweed has no more than two true leaves (2-inch maximum height). Weeds too large for optimum control will be suppressed, but may recover after 2 to 3 weeks. Spraying at the cotyledon stage is not recommended. Do not apply if rainfall is expected within 6 hours' after application.

Transient leaf yellowing and/or growth retardation (stunt) of soybeans may occur following application of Python WDG. These effects will be evident for 5 to 7 days after application to soybeans under stress. Under favorable growing conditions the crop will quickly recover.

ACCEPTED

JUL 22 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under

62719-277

Application Rate: Python WDG at 0.125 oz/acre (single application only).

Refer to the rate table in Application and Mixing section of product label to determine the number of water soluble packets for acreage to be treated.

Use of Surfactants: All postemergence applications of Python WDG in soybeans must include a non-ionic surfactant at 0.25% volume/volume (1qt/100 gal). Use a good quality surfactant with at least 80% active ingredient (of which at least 50% is actual non-ionic surfactant). Under extremely dry growing conditions, use of an agriculturally approved sprayable liquid fertilizer together with the non-ionic surfactant, may enhance control. Use 28%, 30%, or 32% urea ammonium nitrate at 2.5% volume/volume (2.5 gal/100 gal). **Note: Do not use liquid fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur.** Use only EPA approved surfactants for use on food crops.

Tank Mixing: Python WDG may be applied alone or in tank mix combination with other herbicides registered for postemergence application in soybeans unless tank mixing is specifically prohibited on the Python WDG label or on the label of the tank mix product. When Python WDG is tank-mixed with a companion herbicide, follow all applicable use directions, including precautions, restrictions and limitation listed on the manufacturer's label.

Note: Python WDG is not recommended as a postemergence application tank mix partner with Classic due to risk of crop injury.

Python WDG tank mixes with products for postemergence grass control such as Assure, Fusilade 2000 or Poast will not affect the performance of Python WDG, however, the performance of the grass control product may be adversely affected through herbicide antagonism. For best results, application of postemergence grass control products should be delayed for 3 days after applying Python WDG.

*Trademark of Dow AgroSciences LLC

EPA-accepted: 11-13-97