

62719-280

09/11/2008

1/41



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SEP 11 2008

Ms. Kimberly Gilbert
Regulatory Leader
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

RE: Notification of Minor Label Change per PR Notice 98-10 and PRN 2008-1
EPA Registration Number: 62719-280
Date of Submission: August 4, 2008

Dear Ms. Gilbert:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated August 4, 2008, for the product, T&O Fertilizer-Contains Gallery Plus Team. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington", is written above the typed name.

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

2/41

Application for Pesticide - Section I

1. Company/Product Number Dow AgroSciences LLC/62719-280	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/T&O Fert-Contains Gallerv plus Tean	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>SEP 11 2008</u> Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and PR Notice 2008-1 and EPA regulations at 40 CFR 152.46, 40 CFR 156.10, and 156.80 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and PR Notice 2008-1 and 40 CFR 152.46, 40 CFR 156.10, and 156.80 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container.	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Kimberly S. Gilbert		Title Regulatory Manager	
		Telephone No. (Include Area Code) (317) 337-4605 (fax: 317-337-4649)	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory Manager	
4. Typed Name Kimberly S. Gilbert (ksgilbert@dow.com)		5. Date August 4, 2008	

308/2E
August 4, 2008



Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U. S. Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

T&O FERTILIZER – CONTAINS GALLERY PLUS TEAM (A.I. ISOXABEN, BENEFIN, TRIFLURALIN)
EPA REGISTRATION NUMBER: 62719-280
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICES 98-10 AND 2008-1

Enclosed please find labeling for the notification action of T&O Fertilizer – Contains Gallery® plus Team® specialty herbicide. The following changes have been made by notification:

Sub Label A

1. Sale copy: To comply with Florida fertilizer law 5E-1.003, added "Covers up to XX,000 sq ft (XXX' x XX') and/or This bag covers XX,000 sq ft."
2. Established Turfgrass: To comply with Florida fertilizer law 5E-1.003, added new bullet point to Specific Use Restrictions be included for those products registered in the state of Florida.

Sub Label B

3. Sale copy: To comply with Florida fertilizer law 5E-1.003, added (1) "In the state of Florida, do not apply near water. . . onto your lawn/garden."; (2) "Covers up to XX,000 sq ft (XXX' x XX') and/or This bag covers XX,000 sq ft."
4. Environmental Hazards: Per PR Notice 2008-1, revised statement.
5. Established Home Lawns: To comply with Florida fertilizer law 5E-1.003, added new bullet point to Specific Use Restrictions to be included for those products registered in the state of Florida.

This notification is consistent with the provisions of PR Notice 98-10 and PR Notice 2008-1 and EPA regulations at 40 CFR 152.46, 40 CFR 156.10, and 156.80 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and PR Notice 2008-1 and 40 CFR 152.46, 40 CFR 156.10, and 156.80 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Contents of Submission

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled T&O Fertilizer-Contains Gallery plus Team (Q8B/T&O Fertilizer - Contains Gallery Plus Team/MSTR Notif/08-04-08) (36 Pages plus Registration Notes) (5 Copies)

T&O Fertilizer - Contains Gallery[®] plus Team[®]

EPA Reg. No. 62719-280

The master label this product consists of separate, self-contained sub labels for professional use and residential use, respectively.

Registration Notes:

Source label text based on EPA accepted copy dated September 10, 2007 and notifications coded "Q8B / T&O Fertilizer – Contains Gallery plus Team / MSTR Notif / 10-05-07" and "Q8B / T&O Fertilizer – Contains Gallery Plus Team / MSTR Notif / 04-09-08."

Changes by notification:


Sub Label A

1. Sale copy: To comply with Florida fertilizer law 5E-1.003, added "Covers up to XX,000 sq ft (XXX' x XX') and/or This bag covers XX,000 sq ft."
2. Established Turfgrass: To comply with Florida fertilizer law 5E-1.003, added new bullet point to Specific Use Restrictions to be included for those products registered in the state of Florida.

Sub Label B

3. Sale copy: To comply with Florida fertilizer law 5E-1.003, added (1) "In the state of Florida, do not apply near water. . . onto your lawn/garden."; and (2) "Covers up to XX,000 sq ft (XXX' x XX') and/or This bag covers XX,000 sq ft."
4. Environmental Hazards: Per PR Notice 2008-1, revised statement.
5. Established Home Lawns: To comply with Florida fertilizer law 5E-1.003, added new bullet point to Specific Use Restrictions to be included for those products registered in the state of Florida.

[®]Trademark of Dow AgroSciences LLC



[Sub Label A: Commercial Uses]

(Base label for rigid containers 5 gal or less):

T&O Fertilizer - Contains Gallery® plus Team®**Specialty Herbicide**

A fertilizer containing Gallery and Team, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals.

For professional use only.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or


For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]



and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide
and isomers 0.38%

trifluralin: α, α, α -trifluoro-2,6-dinitro-*N*,
N-dipropyl-*p*-toluidine 0.39%

benefin: *N*-butyl-*N*-ethyl- α, α, α -trifluoro-2,6-dinitro-*p*-toluidine 0.76%

Other Ingredients 98.47%

Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P_2O_5)	X.XX%
Soluble Potash (K_2O)	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

Indianapolis, Indiana 46268

Net Weight _____

(Base label for nonrigid containers any size):

T&O Fertilizer - Contains Gallery[®] plus Team[®]

Specialty Herbicide

A fertilizer containing Gallery and Team, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals.

For professional use only.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegrass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

11/41

and/or
University tested [and researched]
 and/or
Labeled for use around a [wide] variety of trees [and shrubs]
 and/or
Never see crabgrass again
 and/or
Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-
 5-isoxazolyl]-2,6-dimethoxybenzamide
 and isomers 0.38%
 trifluralin: α,α,α -trifluoro-2,6-dinitro-*N*,
N-dipropyl-*p*-toluidine 0.39%
 benefin: *N*-butyl-*N*-ethyl- α,α,α -
 trifluoro-2,6-dinitro-*p*-toluidine 0.76%
 Other Ingredients 98.47%
 Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

Keep Out of Reach of Children

CAUTION**Precautionary Statements****Hazards to Humans and Domestic Animals**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

12/41

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight _____

13/41

(Base label for refillable rigid containers greater than 5 gal):

T&O Fertilizer - Contains Gallery® plus Team®

Specialty Herbicide

A fertilizer containing Gallery and Team, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals.

For professional use only.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control[/prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds**Active Ingredients:**

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-
5-isoxazoly]-2,6-dimethoxybenzamide
and isomers..... 0.38%

trifluralin: α,α,α -trifluoro-2,6-dinitro-*N*,
N-dipropyl-*p*-toluidine 0.39%

benefin: *N*-butyl-*N*-ethyl- α,α,α -
trifluoro-2,6-dinitro-*p*-toluidine 0.76%

Other Ingredients 98.47%

Total..... 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P_2O_5)	X.XX%
Soluble Potash (K_2O)	X.XX%

Keep Out of Reach of Children**CAUTION****Precautionary Statements****Hazards to Humans and Domestic Animals**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

15/41

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

EPA Est. _____

16/41

Net Weight _____

17/41

(Base label for nonrefillable rigid containers larger than 5 gal):

T&O Fertilizer - Contains Gallery® plus Team®

Specialty Herbicide

A fertilizer containing Gallery and Team, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals.

For professional use only.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or


Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]



A 10x10 grid of 100 small, stylized human figures arranged in a structured pattern, representing a population.

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.38%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.39%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.76%
Other Ingredients	98.47%
Total	100.00%

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

CAUTION

Hazards to Humans and Domestic Animals

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

19/41

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

Indianapolis, Indiana 46268

Net Weight

 $20/41$

21/41

(Label booklet cover):

T&O Fertilizer - Contains Gallery® plus Team®

Specialty Herbicide

A fertilizer containing Gallery and Team, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals.

For professional use only.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegrass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or


Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

22/11

and/or
University tested [and researched]
 and/or
Labeled for use around a [wide] variety of trees [and shrubs]
 and/or
Never see crabgrass again
 and/or
Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-
 5-isoxazolyl]-2,6-dimethoxybenzamide
 and isomers 0.38%
 trifluralin: α, α, α -trifluoro-2,6-dinitro-*N*,
N-dipropyl-*p*-toluidine 0.39%
 benefin: *N*-butyl-*N*-ethyl- α, α, α -
 trifluoro-2,6-dinitro-*p*-toluidine 0.76%
 Other Ingredients 98.47%
 Total 100.00%



[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight _____

23/11

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons (or pets), either directly or through drift. Only protected handlers may be in the area during application.

24/41

Entry Restrictions: Do not enter or allow other people (or pets) to enter the treated areas until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this

25/41

procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established turfgrass and landscape ornamentals (trees, shrubs and groundcovers) for control of certain broadleaf weeds and annual grasses. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliarly applied postemergence herbicide either before or after application of this product.

General Use Precautions and Restrictions

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turfgrass is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Apply this product only to established turfgrass. Apply to newly seeded turfgrass (including overseeded turfgrass) only after seedlings are established (three leaf stage and tillering) and well rooted (generally after 3 mowings). Do not overseed established turfgrass sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 175 lb of product per acre unless injury can be tolerated, or 12 to 16 weeks after a single application of 265 lb of product per acre unless injury can be tolerated.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. Herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turfgrass that is weakened due to severe winter weather or is not well established.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turfgrass injury may occur.

Treatment of Turfgrass or Ornamental Species Not Listed on the Label for T&O Fertilizer - Contains Gallery Plus Team

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass

Common Name	Scientific Name
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

26/11

Established Warm Season Turfgrass

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum</i> var. <i>Saurae parodi</i>
bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name	Scientific Name
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
birch, paper	<i>Betula papyrifera</i>
cherry	<i>Prunus</i> spp.
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
honeylocust	<i>Gleditsia triacanthos</i>
maple, flowering	<i>Abutilon hybridum</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
oak, pin	<i>Quercus palustris</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, scotch	<i>Pinus sylvestris</i>
plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>

Ornamental Shrubs

Common Name	Scientific Name
arborvitae	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cinquefoil	<i>Potentilla fruticosa</i>
cleysera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
elaeagnus, silverberry	<i>Elaeagnus pungens</i>
euonymus, winged	<i>Euonymus alata</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>

27/41

forsythia	<i>Forsythia</i> spp.
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
juniper, creeping	<i>Juniperus horizontalis</i>
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
myrtle, wax	<i>Myrica cerifera</i>
privet	<i>Ligustrum</i> spp.
pyracantha	<i>Pyracantha fortuneana</i>
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spirea, vanhoutte	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

Ornamental Groundcover Plantings

Common Name	Scientific Name
ceanothus	<i>Ceanothus</i> spp.
coyote brush	<i>Baccharis pilularis</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
gazania	<i>Gazania</i> spp.
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lilyturf, bigblue	<i>Liriope muscari</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rosemary	<i>Rosmarinus officinalis</i>
snow-in-summer	<i>Cerastium tomentosum</i>
St. Johnswort	<i>Hypericum coris</i>

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus</i> spp.
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
large crabgrass	
smooth crabgrass	
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria</i> spp.
bottlegrass	
bristlegrass	
giant foxtail	
green foxtail	
pigeongrass	
robust foxtail	

geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds controlled by a repeat application when applied prior to weed germination:

Common Name	Scientific Name
clover spp. ¹	<i>Trifolium</i> spp.
clover, white ¹	<i>Trifolium repens</i>
dandelion ¹	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
plantain, slender	<i>Plantago elongata</i>

¹From seed only

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	175	XXX	XXX	XXX	265
XXXXXXX	XX	XX	XX	XX	XX

Uses

Established Turfgrass

Application Timing

Apply in late summer or early fall, approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft). A minimum of 175 lb per acre (4 lb per 1000 sq ft) is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

29/41

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** Cool season turfgrass – 175 lb of T&O Fertilizer – Contains Gallery plus Team per acre; warm season turfgrass – 265 lb of T&O Fertilizer – Contains Gallery plus Team per acre.
- **Maximum Yearly Rate:** Cool season turfgrass – 350 lb of T&O Fertilizer – Contains Gallery plus Team per acre; warm season turfgrass – 530 lb of T&O Fertilizer – Contains Gallery plus Team per acre.
- **Minimum Re-Treatment Interval:** Cool season turfgrass – 56 to 70 days; warm season turfgrass – 70 to 84 days.
- This product must be watered in as soon as possible after application.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.
- Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, and groundcovers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing

Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds. Do not apply this product to plantings when the foliage is wet.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft) as a spring application or 265 lb per acre (6 lb per 1000 sq ft) as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks

following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 265 lb of T&O Fertilizer – Contains Gallery plus Team per acre.
 - **Maximum Yearly Rate:** 530 lb of T&O Fertilizer – Contains Gallery plus Team per acre.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 265 lb of T&O Fertilizer – Contains Gallery plus Team per acre.
 - **Maximum Yearly Rate:** 530 lb of T&O Fertilizer – Contains Gallery plus Team per acre.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

[Sub Label B: Residential Uses]

(Base label):

T&O Fertilizer - Contains Gallery® plus Team®**Specialty Herbicide**

A fertilizer containing Gallery and Team, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals.

For residential use only.

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegrass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-
5-isoxazolyl]-2,6-dimethoxybenzamide
and isomers 0.38%
trifluralin: α,α,α -trifluoro-2,6-dinitro-*N*,
N-dipropyl-*p*-toluidine 0.39%
benefin: *N*-butyl-*N*-ethyl- α,α,α -
trifluoro-2,6-dinitro-*p*-toluidine 0.76%
Other Ingredients 98.47%
Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P_2O_5)	X.XX%
Soluble Potash (K_2O)	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

33/41

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight _____

34/41

(Label booklet cover):

T&O Fertilizer - Contains Gallery[®] plus Team[®]

Specialty Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals.

For residential use only.

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control[/prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

35/41

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as]**[crabgrass] [goosegrass] [barnyardgrass]**

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds**Active Ingredients:**isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-

5-isoxazolyl]-2,6-dimethoxybenzamide

and isomers 0.38%

trifluralin: α,α,α -trifluoro-2,6-dinitro-*N*,*N*-dipropyl-*p*-toluidine 0.39%benefin: *N*-butyl-*N*-ethyl- α,α,α -trifluoro-2,6-dinitro-*p*-toluidine 0.76%

Other Ingredients 98.47%

Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**Guaranteed Fertilizer Analysis**

Total Nitrogen (N) X.XX%

Available Phosphoric Acid (P_2O_5) X.XX%Soluble Potash (K_2O) X.XX%**Keep Out of Reach of Children****CAUTION****Refer to inside of label booklet for additional precautionary information including Directions for Use.****Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for**Dow AgroSciences LLC****9330 Zionsville Road****Indianapolis, Indiana 46268****Net Weight** _____

36/41

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

How This Product Works

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established home lawns and other turfgrass areas and landscape ornamentals (trees, shrubs and groundcovers) for control of certain broadleaf weeds and annual grasses. This product also feeds the lawn with a fertilizer designed to provide prolonged, even greening without excess top growth.

This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliarly applied postemergence herbicide either before or after application of this product.

General Use Precautions and Restrictions

- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turfgrass is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Apply this product only to established turfgrass. Apply to newly seeded turfgrass (including overseeded turfgrass) only after seedlings are established and well rooted (generally after 3 mowings). Do not overseed established turfgrass sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 4 lb of product per 1000 sq ft unless injury can be tolerated, or 12 to 16 weeks after a single application of 6 lb of product per 1000 sq ft unless injury can be tolerated.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. Herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turfgrass that is weakened due to severe winter weather or is not well established.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turfgrass injury may occur.

Treatment of Turfgrass or Ornamental Species Not Listed on the Label for T&O Fertilizer – Contains Gallery Plus Team

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass	Established Warm Season Turfgrass
bentgrass, colonial	bahiagrass
bentgrass, creeping	Bermudagrass
bluegrass, Kentucky	centipedegrass
fescue, chewings	fescue, tall
fescue, creeping red	St. Augustinegrass
fescue, sheep	zoysiagrass
fescue, tall	
ryegrass, perennial	

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

38/41

Ornamental Trees

almond	peach
apple, crabapple	pine, Austrian
apricot	pine, eastern white
birch, paper	pine, Japanese black
cherry	pine, scotch
cottonwood	plum
dogwood, flowering	redbud, eastern
dogwood, kousa	spruce, Colorado
honeylocust	spruce, Norway
maple, flowering	sweetgum
maple, red	sycamore
maple, silver	walnut, black
oak, pin	

Ornamental Shrubs

arborvitae	honeysuckle
azalea	juniper, creeping
boxwood, littleleaf	laurel, mountain
camellia, Japanese	lilac, common
camellia, sasanqua	myrtle, wax
cinquefoil	privet
cleyera, Japanese	pyracantha
cotoneaster, cranberry	redcedar, eastern
elaeagnus, silverberry	rhododendron
euonymus, winged	spiraea, vanhoutte
euonymus, wintercreeper	viburnum
forsythia	yew, Japanese
holly	yewpine

Ornamental Groundcover Plantings

ceanothus	ivy, English
coyote brush	lilyturf, bigblue
daisy trailing African	plumbago, dwarf
gazania	rosemary
ice plant, largeleaf	snow-in-summer
ivy, Algerian	St. Johnswort

Weeds Controlled**Weeds controlled when applied prior to weed germination:**

annual bluegrass	geranium, Carolina
barnyardgrass	goosegrass
burweed, lawn	henbit
buttercup	knotweed, prostrate
carpetweed	mallow, little
celery, wild	parsley - piert
chickweed, common	pennywort, lawn
chickweed, mouseear	pineappleweed
crabgrass	speedwell, thymeleaf
large crabgrass	spurge, prostrate
smooth crabgrass	spurge, spotted
foolsparsley	spurry, corn

39/41

foxtails	woodsorrel, creeping
bottlegrass	woodsorrel, yellow
bristlegrass	
giant foxtail	
green foxtail	
pigeongrass	
robust foxtail	

Weeds controlled by a repeat application when applied prior to weed germination:

clover spp. ¹	medic, black
clover, white ¹	plantain, broadleaf
dandelion ¹	plantain, buckhorn
groundsel, common	plantain, slender

¹From seed only**How to Apply**

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	4	XX	XX	XX	6
XXXXXXXX	XX	XX	XX	XX	XX

Uses**Established Home Lawns****When to Apply**

Apply in late summer or early fall, approximately 3 weeks prior to germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer.

How Much to Apply

Apply this product at a rate of 4 to 6 lb per 1000 sq ft. A minimum of 4 lb per 1000 sq ft is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 6 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Maximum Yearly Rate:** 12 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turfgrass – 56 to 70 days; warm season turfgrass – 70 to 84 days.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

40/41

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, and groundcovers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

When to Apply

Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds. Do not apply this product to plantings when the foliage is wet.

How Much to Apply

Apply this product at a rate of 4 to 6 lb per 1000 sq ft as a spring application or 6 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 6 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.
 - **Maximum Yearly Rate:** 12 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 6 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.
 - **Maximum Yearly Rate:** 12 lb of T&O Fertilizer – Contains Gallery plus Team per 1000 sq ft.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent

41/41

risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™Trademark of Dow AgroSciences LLC
EPA accepted 09/10/07