

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

62719-289

02/10/2006

119

P4A / Turf Fertilizer Contains Team Pro X.XX% / MSTR Amend / 12-12-05

[Sub Label A: Commercial Uses]

(Product Label)

**Turf Fertilizer - Contains Team™ Pro {X.XX<sup>(a)</sup>}%**

A granular fertilizer containing Team Pro herbicide for the preemergence control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grass and broadleaf weeds in established turfgrass and established landscape ornamentals.

**Herbicide**

Active ingredient(s):

benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine ..... {X.XX (b)}%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine ..... {X.XX (c)}%

Inert ingredients ..... {X.XX (d)}%

Total ..... 100.00%

**ACCEPTED**

FEB -2 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-289

Contains [X.XX]<sup>††</sup> pounds active ingredient per [XX] pound bag.

[<sup>††</sup>Editor's note: Bag size and calculated amount of active ingredient per bag provided by supplemental distributor.]

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N) .....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O) .....	X.X%

Keep Out of Reach of Children

**WARNING**

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Net Weight - \_\_\_\_\_

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## WARNING

**Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin**

Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

---

### Directions for Use

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated area (except those involved in the watering) until the watering-in is complete and the surface is dry.

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**General Information**

This fertilizer product contains Team™ Pro, a preemergence herbicide for control of annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

**Weeds Controlled**

<b>Common Name</b>	<b>Scientific Name</b>
<b>Annual Grasses</b>	
annual bluegrass	<i>Poa annua</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
crabgrass, large or hairy	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
goosegrass (silver crabgrass or crowfootgrass)	<i>Eleusine indica</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria glauca</i>

<b>Broadleaf Weeds</b>	
chickweed, common	<i>Stellaria media</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
pigweed	<i>Amaranthus spp.</i>
purslane, common	<i>Portulaca oleracea</i>

**Weeds Suppressed**

<b>Common Name</b>	<b>Scientific Name</b>
creeping spurge	<i>Euphorbia serpens</i>
creeping woodsorrel	<i>Oxalis corniculata</i>
panicum, Texas	<i>Panicum texanum</i>
prostrate spurge	<i>Euphorbia humistrata</i>

pusley, Florida  
spotted soursue  
woodsorrel

*Richardia scabra*  
*Euphorbia maculata*  
*Oxalis* spp.

**Uses**

**Established Turfgrass**

**When to Apply:** For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons or for lawn care operator programs.

**How to Apply:** Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**Activation:** Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

**Important Use Precautions:** Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application. This product will not harm most nearby ornamentals.

This product may be applied to the following turfgrass species:

<b>Established Cool Season Turfgrasses</b>	<b>Established Warm Season Turfgrasses</b>
bentgrass fairways	bahiagrass
fescues	Bermudagrass
Kentucky bluegrass	centipedegrass
perennial ryegrass	fescue, tall
	St. Augustinegrass
	zoysiagrass

**Broadcast Application Rates**

Application Timing	Application Rate	Specific Use Directions
<b>Spring:</b> To control spring-to-summer germinating annual weeds. <b>Fall:</b> To control fall germinating weeds.	{XXX(e)} to {XXX(g)} lb/acre ({X.X(h)} to {X.X(j)} lb/1000 sq ft)	In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination.
<b>Split Application:</b> For extended control of spring and summer annual weeds.	{XXX(e)} lb/acre followed by {XXX(e)} lb/acre ({X.X(h)} followed by {X.X(h)} lb/1000 sq ft)	In cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later.

	{XXX(f)} lb/acre followed by {XXX(f)} lb/acre ({X.X(i)} followed by {X.X(i)} lb/1000 sq ft)	In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.
--	---	---

**Reseeding Precautions**

Team Pro, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding following the last application of this product should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of Product Applied Per Acre	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of {XXX(e)} lb/acre ({XXX(h)} lb/1000 sq ft)	8
Single application {XXX(g)} lb/acre ({XXX(i)} lb/1000 sq ft)	12 to 16
Split application of {XXX(e)} lb/acre ({XXX(h)} lb/1000 sq ft)	8 to 10
Split application of {XXX(f)} lb/acre ({XXX(i)} lb/1000 sq ft)	10 to 12

**[Spreader Settings]**

[Editors note: Formulator label may include a table of recommended spreader equipment and spreader settings for the fertilizer blend used.]

**Established Landscape Ornamental Plants**

This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers.

**When to Apply:** Apply one to two weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas.

**Soil Preparation:** This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

**How to Apply:** See paragraph in Established Turfgrass section.

**Activation:** Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall within a few hours after application.

**Broadcast Application Rates**

Season of Application	Pounds per 1000 sq ft
spring application (summer annuals)	{X.X(h)} to {X.X(i)} lb/1000 sq ft
fall application (winter annuals)	{X.X(h)} to {X.X(i)} lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application.

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

### Ornamental Trees

<b>Common Name</b>	<b>Scientific Name</b>
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
ash, white	<i>Fraxinus americana</i>
bald cypress	<i>Taxodium distichum</i>
birch, European white	<i>Betula pendula</i>
blackgum	<i>Nyssa sylvatica</i>
cherry	<i>Prunus</i> spp.
chestnut, Chinese	<i>Castanea mollissima</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
douglas fir	<i>Pseudotsuga menziesii</i>
fir, balsam	<i>Abies balsamea</i>
hemlock, Canada	<i>Tsuga canadensis</i>
honeylocust	<i>Gleditsia triacanthos</i>
larch, Japanese	<i>Larix kaempferi</i>
locust, black	<i>Robinia pseudoacacia</i>
maple, Norway	<i>Acer platanoides</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
maple, sugar	<i>Acer saccharum</i>
oak, pin	<i>Quercus palustris</i>
oak, scarlet	<i>Quercus coccinea</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, loblolly	<i>Pinus taeda</i>
pine, red	<i>Pinus resinosa</i>
pine, scotch	<i>Pinus sylvestris</i>
planetree, London	<i>Platanus acerifolia</i>
plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
spruce, white	<i>Picea glauca</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
tuliptree	<i>Liriodendron tulipifera</i>
walnut, black	<i>Juglans nigra</i>

### Ornamental Shrubs

<b>Common Name</b>	<b>Scientific Name</b>
andromeda, Japanese	<i>Pieris japonica</i>
arborvitae, American	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
barberry, Japanese	<i>Berberis thunbergii</i>
barberry, mento	<i>Berberis mentorensis</i>

boxwood, common	<i>Buxus sempervirens</i>
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cherrylaurel, American	<i>Prunus caroliniana</i>
cinquefoil	<i>Potentilla</i> spp.
cleyera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
cotoneaster, zabel	<i>Cotoneaster zabelii</i>
deutzia	<i>Deutzia</i> spp.
elaeanthus, silverberry	<i>Elaeagnus pungens</i>
euonymus, spreading	<i>Euonymus kiautschovica</i>
euonymus, winged	<i>Euonymus alatus</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
firethorn	<i>Pyracantha</i> spp.
forsythia	<i>Forsythia</i> spp.
guava, pineapple	<i>Feijoa sellowiana</i>
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
India nawthorn	<i>Raphiolepis indica</i>
juniper	<i>Juniperus</i> spp.
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
mockorange	<i>Philadelphus</i> spp.
pittosporum, Japanese	<i>Pittosporum tobira</i>
privet	<i>Ligustrum</i> spp.
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spiraea, vanhoutte	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
weigela	<i>Weigela</i> spp.
willow	<i>Salix</i> spp.
yew, anglojap	<i>Taxus media</i>
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

### Ornamental Groundcover Plantings

Common Name	Scientific Name
aaronsbeard	<i>Hypericum calycinum</i>
bellflower, adriatic	<i>Campanula elatines</i>
bellflower, poscharsky	<i>Campanula poscharskyana</i>
ceanothus	<i>Ceanothus</i> spp.
coreopsis	<i>Coreopsis</i> spp.
cotoneaster	<i>Cotoneaster</i> spp.
coyote brush	<i>Baccharis pilularis</i>
crown vetch	<i>Coronilla vana</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
fern, asparagus	<i>Asparagus densiflorus</i>
gazania	<i>Gazania</i> spp.
germander	<i>Teucrium chamaedrys</i>
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lily-of-the-nile	<i>Agapanthus</i> spp.
lilyturf, bigblue	<i>Liriope muscari</i>



marigold	<i>Tagetes</i> spp.
myoporum	<i>Myoporum laetum</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rockrose	<i>Cistus</i> spp.
rosemary	<i>Rosmarinus officinalis</i>
rupturewort	<i>Hemiaria glabra</i>
snow-in-summer	<i>Cerastium tomentosum</i>
speedwell	<i>Veronica</i> spp.
St. Johnswort	<i>Hypericum coris</i>
stonecrop (sedum)	<i>Sedum</i> spp.
strawberry, beach	<i>Fragaria chiloensis</i>
thrift	<i>Armeria maritima</i>
verbena	<i>Verbena</i> spp.
wirevine, creeping	<i>Muehlenbeckia axillaris</i>
yarrow, wooly	<i>Achillea tomentosa</i>

### Roses and Other Established Flowers

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

---

### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

### Limitation of Remedies

---

9/9

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™ Trademark of Dow AgroSciences LLC  
EPA-accepted \_\_\_/\_\_\_/\_\_\_

**NEXT**

**LABEL**

62719-289

02/02/2006

1/8

P4A / Turf Fertilizer Contains Team Pro X.XX% / MSTR Amend / 12-12-05

[Sub Label B: Residential Uses]

(Product Label)

**Turf Fertilizer - Contains Team™ Pro {X.XX (a)}%**

A granular fertilizer containing Team Pro herbicide for the preemergence control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grass and broadleaf weeds in established home lawns and landscape ornamentals.

**Herbicide**

ONE STEP – extended grassy weed control

Active Ingredient(s):

benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine ..... {X.XX (b)}%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine ..... {X.XX (c)}%

Inert Ingredients ..... {X.XX (d)}%

Total ..... 100.00%

**ACCEPTED**

FEB - 2 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-289

Contains [X.XX]<sup>††</sup> pounds active ingredient per [XX] pound bag.

[<sup>††</sup>Editor's note: Bag size and calculated amount of active ingredient per bag provided by supplemental distributor.]

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O).....	X.X%

**Keep Out of Reach of Children**

**WARNING**

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

2/8

P4A / Turf Fertilizer Contains Team Pro X.XX% / MSTR Amend / 12-12-05

**Net Weight - \_\_\_\_\_**

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals**

**WARNING**

**Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin**

Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, chemical-resistant gloves, shoes plus socks and protective eyewear.

**User Safety Recommendations**

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

---

**Directions for Use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

**Application Restrictions:** Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

**Entry Restrictions:** Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

### **Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

### **General Information**

This fertilizer product contains Team™ Pro, a preemergence herbicide for control of annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.

### **Weeds Controlled**

#### **Annual Grasses**

annual bluegrass  
barnyardgrass (watergrass)  
crabgrass, large or hairy  
crabgrass, smooth  
goosegrass (silver crabgrass  
or crowfootgrass)  
green foxtail  
yellow foxtail

#### **Broadleaf Weeds**

chickweed, common  
knotweed, prostrate  
lambsquarters, common  
pigweed  
purslane, common

### **Weeds Suppressed**

creeping spurge  
creeping woodsorrel  
panicum, Texas  
prostrate spurge  
pusley, Florida  
spotted spurge  
woodsorrel

### **Uses**

#### **Established Home Lawns**

**When to Apply:** For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons.

**How to Apply:** Apply this product evenly over the lawn using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands.

576

**Activation:** Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

**Important Use Precautions:** Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application. This product will not harm most nearby ornamentals.

This product may be applied to the following turfgrass species:

<b>Established Cool Season Turfgrasses</b>	<b>Established Warm Season Turfgrasses</b>
bentgrass fairways	bahiagrass
fescues	Bermudagrass
Kentucky bluegrass	centipedegrass
perennial ryegrass	fescue, tall
	St. Augustinegrass
	zoysiagrass

**Broadcast Application Rates**

Application Timing	Application Rate	Specific Use Directions
<b>Spring:</b> To control spring-to-summer germinating annual weeds. <b>Fall:</b> To control fall germinating weeds.	{X.X(h)} to {X.X(j)} lb/1000 sq ft	In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination.
<b>Split Application:</b> For extended control of spring and summer annual weeds.	{X.X(h)} per 1000 sq ft plus a second application of {X.X(h)} lb per 1000 sq ft	In cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later.
	{X.X(j)} lb per 1000 sq ft plus a second application of {X.X(j)} lb per 1000 sq ft	In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.

**Reseeding Precautions**

Team Pro, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding following the last application of this product should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of Product Applied (Pounds per 1000 sq ft)	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of {XXX(h)} lb/1000 sq ft	8
Single application {XXX(j)} lb/1000 sq ft	12 to 16
Split application of {XXX(h)} lb/1000 sq ft	8 to 10
Split application of {XXX(j)} lb/1000 sq ft	10 to 12



**[Spreader Settings]**

[Editors note: Formulator label may include a table of recommended spreader equipment and spreader settings for the fertilizer blend used.]

**Established Landscape Ornamental Plants**

This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers.

**When to Apply:** Apply one to two weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas.

**Soil Preparation:** This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

**How to Apply:** See paragraph in Established Home Lawns section.

**Activation:** Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall within a few hours after application.

**Broadcast Application Rates**

Season of Application	Pounds per 1000 sq ft
spring application (summer annuals)	{X.X(h)} to {X.X(j)} lb/1000 sq ft
fall application (winter annuals)	{X.X(h)} to {X.X(j)} lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application.

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

**Ornamental Trees**

almond	hemlock, Canada	pine, loblolly
apple, crabapple	honeylocust	pine, red
apricot	larch, Japanese	pine, scotch
ash, white	locust, black	planetree, London
bald cypress	maple, Norway	plum
birch, European white	maple, red	redbud, eastern
blackgum	maple, silver	spruce, Colorado
cherry	maple, sugar	spruce, Norway
chestnut, Chinese	oak, pin	spruce, white
cottonwood	oak, scarlet	sweetgum
dogwood, flowering	peach	sycamore
dogwood, kousa	pine, Austrian	tuliptree
douglas fir	pine, eastern white	walnut, black
fir, balsam	pine, Japanese black	

**Ornamental Shrubs**

andromeda, Japanese	forsythia
arborvitae, American	guava, pineapple
azalea	holly
barberry, Japanese	honeysuckle

- |                         |                       |
|-------------------------|-----------------------|
| barberry, mento         | India hawthorn        |
| boxwood, common         | juniper               |
| boxwood, littleleaf     | laurel, mountain      |
| camellia, Japanese      | lilac, common         |
| camellia, sasanqua      | mockorange            |
| cherrylaurel, American  | pittosporum, Japanese |
| cinquefoil              | privet                |
| cleyera, Japanese       | redcedar, eastern     |
| cotoneaster, cranberry  | rhododendron          |
| cotoneaster, zabel      | spiraea, vanhoutte    |
| deutzia                 | viburnum              |
| elaeanthus, silverberry | weigela               |
| euonymus, spreading     | willow                |
| euonymus, winged        | yew, anglojap         |
| euonymus, wintercreeper | yew, Japanese         |
| firethorn               | yewpine               |

**Ornamental Groundcover Plantings**

- |                        |                    |
|------------------------|--------------------|
| aaronsbeard            | lilyturf, bigblue  |
| bellflower, adriatic   | marigold           |
| bellflower, poscharsky | myoporum           |
| ceanothus              | plumbago, dwarf    |
| coreopsis              | rockrose           |
| cotoneaster            | rosemary           |
| coyote brush           | rupturewort        |
| crown vetch            | snow-in-summer     |
| daisy trailing African | speedwell          |
| fern, asparagus        | St. Johnswort      |
| gazania                | stonecrop (sedum)  |
| germander              | strawberry, beach  |
| ice plant, largeleaf   | thrift             |
| ivy, Algerian          | verbena            |
| ivy, English           | wirevine, creeping |
| lily-of-the-nile       | yarrow, wooly      |

**Roses and Other Established Flowers**

- |                   |               |                   |                      |
|-------------------|---------------|-------------------|----------------------|
| African daisy     | dianthus      | marigold          | shasta daisy         |
| aster (perennial) | dusty miller  | marigold, cape    | snapdragon           |
| balsam            | floss flower  | morningglory      | snow-on-the-mountain |
| blackeyed susan   | forget-me-not | nasturtium        | stock                |
| calendula         | four o'clock  | nicotiana         | sunflower            |
| carnation         | gaillardia    | petunia           | sweet alyssum        |
| centaurea, velvet | gladiolus     | phlox             | sweet pea            |
| chrysanthemum     | golden glow   | pincushion flower | sweet sultan         |
| coreopsis         | impatiens     | poppy, California | sweet william        |
| cornflower        | ixora         | portulaca         | vinca                |
| cosmos            | lobelia       | rose              | yarrow               |
| dahlia            | lupine        | salvia            | zinnia               |

**Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

878

---

### **Warranty Disclaimer**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

### **Limitation of Remedies**

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™Trademark of Dow AgroSciences LLC  
EPA-accepted \_\_\_/\_\_\_/\_\_\_