

62719-289

12-10-2007

1/27



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number: 62719-289  
Date of Issuance: 12-10-07

Term of Issuance:

Name of Pesticide Product:  
Turf Fertilizer Contains Team  
Pro 0.86% Herbicide

Name and Address of Registrant (include ZIP Code):

Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, Indiana  
46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

Commercial Label


1. Change the Signal Word to "Warning".
2. To the First Aid section add "If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice."
3. Change the Hazards to Humans and Domestic Animals statements to "Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."

2/27

4. Change the PPE section to "All loaders, applicators, and other handlers must wear: Long-sleeved shirt and long pants, Shoes plus socks, Chemical-resistant gloves (such as or made of any waterproof material), and Goggles, face shield or safety glasses."
5. The entry and application restrictions must include "or pets" (Do not apply... worker or other persons or pets... and Do not enter or allow people or pets...).
6. To the Application instructions add "Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing: This product must be watered in as soon as possible. Do not enter or allow other (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry."
7. To the Storage and Disposal section add a "Pesticide Disposal" heading.
8. To the Warranty section change "Seller makes no" to "To the extent consistent with applicable law, Seller makes no" and "to the extent permitted by law" to "to the extent consistent with applicable law".

Homeowner Use Label

9. Change the Signal Word to "Warning".
10. To the First Aid section add "If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice."
11. Change the Hazards to Humans and Domestic Animals statements to "Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
12. To the Storage and Disposal section add a "Pesticide Disposal" heading.
13. To the Warranty section change "Seller makes no" to "To the extent consistent with applicable law, Seller makes no" and "to the extent permitted by law" to "to the extent consistent with applicable law".

<p>Signature of Approving Official:</p>  <p>Kathryn V. Montague, Acting Product Manager (23) Herbicide Branch, Registration Division (7505P)</p>	<p>Date:</p> <p>12-10-07</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

3/27

[Sub Label A: Commercial Uses]

(Base label for rigid containers 5 gal or less):

**Turf Fertilizer - Contains Team™ Pro 0.86%**

**Herbicide**

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

12-10-07

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-289

Active Ingredients:

benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl- <i>p</i> -toluidine.....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis**

Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O) .....	X.X%

**Keep Out of Reach of Children**

**CAUTION**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

4/27

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose of as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

5/27

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

<sup>TM</sup>Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

**Net Weight** \_\_\_\_\_

6/27

(Base label for nonrigid containers any size):

# Turf Fertilizer - Contains Team™ Pro 0.86%

## Herbicide

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals**

**Active Ingredients:**

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O).....	X.X%

**Keep Out of Reach of Children**  
**CAUTION**

---

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

7/27

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose of as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Net Weight \_\_\_\_\_

8/27

(Base label for rigid containers greater than 5 gal):

# Turf Fertilizer - Contains Team™ Pro 0.86%

## Herbicide

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals**

### Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O) .....	X.X%

**Keep Out of Reach of Children**

## CAUTION

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.**

#### Personal Protective Equipment (PPE)

**All loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.



9/27

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose of as waste.

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Net Weight \_\_\_\_\_

10/27

(Label booklet cover):

# Turf Fertilizer - Contains Team™ Pro 0.86%

## Herbicide

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals**

**Active Ingredients:**

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O) .....	X.X%

**Keep Out of Reach of Children**

## CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

**Net Weight** \_\_\_\_\_

11/27

(Page 1 through end):

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION****Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

---

**Directions for Use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

12/27

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

### **Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

#### **Nonrefillable rigid containers 5 gallons or less:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### **Nonrefillable nonrigid containers:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

#### **Refillable rigid containers larger than 5 gal:**

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### **General Information**

This product is a granular fertilizer containing Team™, a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

### **General Use Precautions and Restrictions**

- Do not apply this product to golf course putting greens.

13/07

- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

**Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 349 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

**This product is recommended for use on the following turfgrass species:**

**Established Cool Season Turfgrass**

bentgrass fairways  
fescues  
Kentucky bluegrass  
perennial ryegrass

**Established Warm Season Turfgrass**

bahiagrass  
Bermudagrass  
centipedegrass  
fescue, tall  
St. Augustinegrass  
zoysiagrass

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

**Ornamental Trees**

**Common Name**

**Scientific Name**

almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
ash, white	<i>Fraxinus americana</i>
bald cypress	<i>Taxodium distichum</i>
birch, European white	<i>Betula pendula</i>
blackgum	<i>Nyssa sylvatica</i>
cherry	<i>Prunus</i> spp.
chestnut, Chinese	<i>Castanea mollissima</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
douglas fir	<i>Pseudotsuga menziesii</i>
fir, balsam	<i>Abies balsamea</i>
hemlock, Canada	<i>Tsuga canadensis</i>
honeylocust	<i>Gleditsia triacanthos</i>
larch, Japanese	<i>Larix kaempferi</i>
locust, black	<i>Robinia pseudoacacia</i>
maple, Norway	<i>Acer platanoides</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
maple, sugar	<i>Acer saccharum</i>
oak, pin	<i>Quercus palustris</i>
oak, scarlet	<i>Quercus coccinea</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>

14/27

pine, loblolly	<i>Pinus taeda</i>
pine, red	<i>Pinus resinosa</i>
pine, scotch	<i>Pinus sylvestris</i>
planetree, London	<i>Platanus acerifolia</i>
plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
spruce, white	<i>Picea glauca</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
tuliptree	<i>Liriodendron tulipifera</i>
walnut, black	<i>Juglans nigra</i>

### Ornamental Shrubs

#### Common Name

andromeda, Japanese  
 arborvitae, American  
 azalea  
 barberry, Japanese  
 barberry, mento  
 boxwood, common  
 boxwood, littleleaf  
 camellia, Japanese  
 camellia, sasanqua  
 cherrylaurel, American  
 cinquefoil  
 cleyera, Japanese  
 cotoneaster, cranberry  
 cotoneaster, zabel  
 deutzia  
 elaeagnus, silverberry  
 euonymus, spreading  
 euonymus, winged  
 euonymus,  
 wintercreeper  
 firethorn  
 forsythia  
 guava, pineapple  
 holly  
 honeysuckle  
 India hawthorn  
 juniper  
 laurel, mountain  
 lilac, common  
 mockorange  
 pittosporum, Japanese  
 privet  
 redcedar, eastern  
 rhododendron  
 spiraea, vanhoutte  
 viburnum  
 weigela  
 willow  
 yew, anglojap

#### Scientific Name

*Pieris japonica*  
*Thuja occidentalis*  
*Rhododendron* spp.  
*Berberis thunbergii*  
*Berberis mentorensis*  
*Buxus sempervirens*  
*Buxus microphylla*  
*Camellia japonica*  
*Camellia sasanqua*  
*Prunus caroliniana*  
*Potentilla* spp.  
*Cleyera japonica*  
*Cotoneaster apiculata*  
*Cotoneaster zabelii*  
*Deutzia* spp.  
*Elaeagnus pungens*  
*Euonymus kiautschovica*  
*Euonymus alatus*  
*Euonymus fortunei*  
  
*Pyracantha* spp.  
*Forsythia* spp.  
*Feijoa sellowiana*  
*Ilex* spp.  
*Lonicera* spp.  
*Raphiolepis indica*  
*Juniperus* spp.  
*Kalmia latifolia*  
*Syringa vulgaris*  
*Philadelphus* spp.  
*Pittosporum tobira*  
*Ligustrum* spp.  
*Juniperus virginiana*  
*Rhododendron* spp.  
*Spiraea vanhouttei*  
*Viburnum* spp.  
*Weigela* spp.  
*Salix* spp.  
*Taxus media*

15/07

yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

**Ornamental Groundcover Plantings**

<b>Common Name</b>	<b>Scientific Name</b>
aaronsbeard	<i>Hypericum calycinum</i>
bellflower, adriatic	<i>Campanula elatines</i>
bellflower, poscharsky	<i>Campanula poscharskyana</i>
ceanothus	<i>Ceanothus</i> spp.
coreopsis	<i>Coreopsis</i> spp.
cotoneaster	<i>Cotoneaster</i> spp.
coyote brush	<i>Baccharis pilularis</i>
crown vetch	<i>Coronilla vana</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
fern, asparagus	<i>Asparagus densiflorus</i>
gazania	<i>Gazania</i> spp.
germander	<i>Teucrium chamaedrys</i>
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lily-of-the-nile	<i>Agapanthus</i> spp.
lilyturf, bigblue	<i>Liriope muscari</i>
marigold	<i>Tagetes</i> spp.
myoporum	<i>Myoporum laetum</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rockrose	<i>Cistus</i> spp.
rosemary	<i>Rosmarinus officinalis</i>
rupturewort	<i>Herniaria glabra</i>
snow-in-summer	<i>Cerastium tomentosum</i>
speedwell	<i>Veronica</i> spp.
St. Johnswort	<i>Hypericum coris</i>
stonecrop (sedum)	<i>Sedum</i> spp.
strawberry, beach	<i>Fragaria chiloensis</i>
thrift	<i>Armeria maritima</i>
verbena	<i>Verbena</i> spp.
wirevine, creeping	<i>Muehlenbeckia axillaris</i>
yarrow, wooly	<i>Achillea tomentosa</i>

**Roses and Other Established Flowers**

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

**Weeds Controlled**

11/6/27

**Weeds controlled when applied prior to weed germination:**

<b>Common Name</b>	<b>Scientific Name</b>
<b>Annual Grasses</b>	
annual bluegrass	<i>Poa annua</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
crabgrass, large or hairy	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
goosegrass (silver crabgrass or crowfoot)	<i>Eleusine indica</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria glauca</i>

<b>Broadleaf Weeds</b>	
chickweed, common	<i>Stellaria media</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambquarters, common	<i>Chenopodium album</i>
pigweed	<i>Amaranthus</i> spp.
purslane, common	<i>Portulaca oleracea</i>

**Weeds Suppressed**

**Weeds suppressed when applied prior to weed germination:**

<b>Common Name</b>	<b>Scientific Name</b>
creeping spurge	<i>Euphorbia serpens</i>
creeping woodsorrel	<i>Oxalis corniculata</i>
panicum, Texas	<i>Panicum texanum</i>
prostrate spurge	<i>Euphorbia humistrata</i>
pusley, Florida	<i>Richardia scabra</i>
spotted spurge	<i>Euphorbia maculata</i>
woodsorrel	<i>Oxalis</i> spp.

**Application Directions**

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**Spreader Settings as a Guide for Calibration**

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	XXX	XXX	XXX	XXX	XXX
XXXXXXXX	XX	XX	XX	XX	XX

**Uses**

**Established Turfgrass**

**Application Timing**



17/27

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

#### Broadcast Application Rates

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	174 to 349lb/acre 4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)
winter annual grass weeds (fall application)	174 to 349 lb/acre 4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 233 lb plus 233 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate.

#### Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 349 lb of Turf Fertilizer – Contains Team Pro 0.86% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

#### Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

#### Application Timing

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

#### Broadcast Application Rates

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

#### Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
  - **Maximum Single Application Rate:** 8 lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
  - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**

18/27

- o **Maximum Single Application Rate:** 8-lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

---

### **Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### **Warranty Disclaimer**

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

### **Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™Trademark of Dow AgroSciences LLC  
EPA accepted    /    /

19/07

**[Sub Label B: Residential Uses]**

(Base label):

**Turf Fertilizer - Contains Team™ Pro 0.86%**  
**Herbicide**

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.**

ONE STEP – extended grassy weed control

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O).....	X.X%

**Keep Out of Reach of Children**

**CAUTION**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, and shoes plus socks.**

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

20/27

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg: No. 62719-289

EPA Est. No. \_\_\_\_\_

™ Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

**Net Weight** \_\_\_\_\_

21/07

(Label booklet cover):

# Turf Fertilizer - Contains Team™ Pro 0.86%

## Herbicide

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.**

ONE STEP – extended grassy weed control

**Active Ingredients:**

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.43%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.43%
Other Ingredients .....	99.14%
Total .....	100.00%

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N).....	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.X%
Soluble Potash (K <sub>2</sub> O).....	X.X%

### Keep Out of Reach of Children

## CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC  
Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

**Net Weight** \_\_\_\_\_

22/27

(Page 1 through end):

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION****Causes Moderate Eye Irritation • Harmful If Swallowed**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, and shoes plus socks.**

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

---

**Directions for Use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

**Entry Restrictions:** Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

23/07

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

## How This Product Works

This product is a granular fertilizer containing Team™, a preemergence herbicide for application to established home lawns and landscape ornamentals (trees, shrubs, groundcovers and flowers) for control of annual grasses and broadleaf weeds.

## General Use Precautions and Restrictions

- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

**Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 4 lb of product per 1000 sq ft, or 12 to 16 weeks after a single application of 8 lb of product per 1000 sq ft unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 4 lb of product per 1000 sq ft rate is applied, or 10 to 12 weeks after the last 5.34 lb of product per 1000 sq ft rate is applied unless injury can be tolerated.

**This product is recommended for use on the following turfgrass species:**

### Established Cool Season Turfgrass

bentgrass fairways  
fescues  
Kentucky bluegrass  
perennial ryegrass

### Established Warm Season Turfgrass

bahiagrass  
Bermudagrass  
centipedegrass  
fescue, tall  
St. Augustinegrass  
zoysiagrass

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

### Ornamental Trees

almond	hemlock, Canada	pine, loblolly
apple, crabapple	honeylocust	pine, red
apricot	larch, Japanese	pine, scotch
ash, white	locust, black	planetree, London
bald cypress	maple, Norway	plum
birch, European white	maple, red	redbud, eastern
blackgum	maple, silver	spruce, Colorado
cherry	maple, sugar	spruce, Norway
chestnut, Chinese	oak, pin	spruce, white
cottonwood	oak, scarlet	sweetgum
dogwood, flowering	peach	sycamore
dogwood, kousa	pine, Austrian	tuliptree
douglas fir	pine, eastern white	walnut, black
fir, balsam	pine, Japanese black	

### Ornamental Shrubs

24/97

andromeda, Japanese	forsythia
arborvitae, American	guava, pineapple
azalea	holly
barberry, Japanese	honeysuckle
barberry, mento	India hawthorn
boxwood, common	juniper
boxwood, littleleaf	laurel, mountain
camellia, Japanese	lilac, common
camellia, sasanqua	mockorange
cherrylaurel, American	pittosporum, Japanese
cinquefoil	privet
cleystera, Japanese	redcedar, eastern
cotoneaster, cranberry	rhododendron
cotoneaster, zabel	spiraea, vanhoutte
deutzia	viburnum
elaeagnus, silverberry	weigela
euonymus, spreading	willow
euonymus, winged	yew, anglojap
euonymus, wintercreeper	yew, Japanese
firethorn	yewpine

**Ornamental Groundcover Plantings**

aaronsbeard	lilyturf, bigblue
bellflower, adriatic	marigold
bellflower, poscharsky	myoporum
ceanothus	plumbago, dwarf
coreopsis	rockrose
cotoneaster	rosemary
coyote brush	rupturewort
crown vetch	snow-in-summer
daisy trailing African	speedwell
fern, asparagus	St. Johnswort
gazania	stonecrop (sedum)
germander	strawberry, beach
ice plant, largeleaf	thrift
ivy, Algerian	verbena
ivy, English	wirevine, creeping
lily-of-the-nile	yarrow, wooly

**Roses and Other Established Flowers**

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

**Weeds Controlled**



25/27

**Weeds controlled when applied prior to weed germination:**

<b>Annual Grasses</b>	<b>Broadleaf Weeds</b>
annual bluegrass	chickweed, common
barnyardgrass (watergrass)	knotweed, prostrate
crabgrass, large or hairy	lambsquarters, common
crabgrass, smooth	pigweed
goosegrass (silver crabgrass or crowfoot)	purslane, common
green foxtail	
yellow foxtail	

**Weeds Suppressed**

**Weeds suppressed when applied prior to weed germination:**

- creeping spurge
- creeping woodsorrel
- panicum, Texas
- prostrate spurge
- pusley, Florida
- spotted spurge
- woodsorrel

**How to Apply**

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate the application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or spinkler irrigation of 0.5 inch or more is required to activate this product.

**Spreader Settings as a Guide for Calibration**

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	XX	XX	XX	XX	XX
XXXXXXXX	XX	XX	XX	XX	XX

**Uses**

**Established Home Lawns**

**When to Apply**

Apply this product 1 to 2 weeks prior to germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

**How Much to Apply**

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	4 to 8 lb/1000 sq ft	8 lb/1000 sq ft

26/07

winter annual grass weeds (fall application)	4 to 8 lb/1000 sq ft	8 lb/1000 sq ft
-------------------------------------------------	----------------------	-----------------

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 4 lb plus 4 lb per 1000 sq ft for cool season turfgrass and in the northern half of the warm season turfgrass area, or 5.34 lb plus 5.34 lb per 1000 sq ft in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 4 lb per 1000 sq ft rate and 10 to 12 weeks following the 5.34 lb per 1000 sq ft rate.

#### Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

#### Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

#### When to Apply

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

#### How to Apply

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

#### Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
  - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
  - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
  - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

---

#### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

#### Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF

27/07

MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™Trademark of Dow AgroSciences LLC  
EPA accepted   /  /