

62719-289

03/30/2009

1/31



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Kimberly S. Gilbert
Product Registration
Dow AgroScience LLC
9330 Zionsville Road
Indianapolis, IN 46268

MAR 30 2009

SUBJECT: Application for Pesticide Notification (PRN 98-10 and 2008-1)
Request General Label Change and Environmental Hazard Statement
EPA Reg. No. 62719-289
Application Dated March 3, 2009

Dear Ms. Gilbert:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 and 2008-1 dated 03/03/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and 2008-1 and finds that the action requested fall within the scope of PRN 98-10 and 2008-1. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number Dow AgroSciences LLC/62719-289	2. EPA Product Manager Joanne I. Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/Turf Fert-Contains Team Pro 0.86%	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

NOTIFICATION
MAR 30 2009

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and PR Notice 2008-1 and the requirements of EPA regulations at 40 CFR 152.46, 40 CFR 156.10, 156.80, 156.140, 156.144, 156.146, and 156.156 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and PR Notice 2008-1 and 40 CFR 152.46, 40 CFR 156.10, 156.80, 156.140, 156.144, 156.146, and 156.156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Kimberly S. Gilbert	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4685 (fax: 317-337-4649)	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Manager		
4. Typed Name Kimberly S. Gilbert (ksgilbert@dow.com)	5. Date March 3, 2009		

Turf Fertilizer - Contains Team[®] Pro 0.86%

EPA Reg. No. 62719-289

The master label consists of separate, self-contained sub labels for commercial use and residential use, respectively.


Registration Notes:

Source label text based on EPA accepted copy dated December 10, 2007 and notifications coded "P4A / Turf Fert Contains Team Pro 0.86% / MSTR Notif / 12-19-07" and P4A / Turf Fert Contains Team Pro 0.86% / MSTR Notif / 04-09-08."

Changes by notification:

1. Professional Use Sub Label A, Sale copy: To comply with Florida fertilizer law 5E-1.003, added "Covers up to XX,000 sq ft (XXX' x XX') and/or This bag covers XX,000 sq ft."
2. Established Turfgrass, Specific Use Restrictions: To comply with Florida fertilizer law 5E-1.003, added new bullet point to be included for those products registered in the state of Florida.
3. Residential Use Sub Label B, Sale copy: To comply with Florida fertilizer law 5E-1.003, added "In the state of Florida, . . . covers XX,000 sq ft."
4. Residential Use Sub Label B, Environmental Hazards: Per PR Notice 2008-1, revised statement.

[®]Trademark of Dow AgroSciences LLC



5/31

[Sub Label A: Commercial Uses]

(Base label):

Turf Fertilizer - Contains Team[®] Pro 0.86%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals

Covers up to XX,000 sq ft (XXX' x XX')
and/or

This bag covers XX,000 sq ft

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]


and/or

University tested [and researched]

and/or

NOTIFICATION

MAR 30 2009



6/31

**Labeled for use around a [wide] variety of trees and shrubs
and/or
Never see crabgrass again**

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro- 2,6-dinitro- <i>p</i> -toluidine	0.43%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.43%
Other Ingredients	99.14%
Total	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

**Keep Out of Reach of Children
WARNING**

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles, faceshield or safety glasses

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

7/31

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrigid containers any size)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers greater than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

8/31

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. _____

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

9/31

Net Weight ____

10/31

(Label booklet cover):

Turf Fertilizer - Contains Team[®] Pro 0.86%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

11/31

Never see crabgrass again

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro- 2,6-dinitro- <i>p</i> -toluidine	0.43%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.43%
Other Ingredients	99.14%
Total	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

Keep Out of Reach of Children

WARNING

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268**

Net Weight _____

12/31

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING**

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles, faceshield or safety glasses

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

13/31

Application Restrictions: Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow people or pets to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

14/31

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Information

This product is a granular fertilizer containing Team[®], a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

General Use Precautions and Restrictions

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 349 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass	Established Warm Season Turfgrass
bentgrass fairways	bahiagrass
fescues	Bermudagrass
Kentucky bluegrass	centipedegrass
perennial ryegrass	fescue, tall
	St. Augustinegrass
	zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name	Scientific Name
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
ash, white	<i>Fraxinus americana</i>
bald cypress	<i>Taxodium distichum</i>
birch, European white	<i>Betula pendula</i>

15/31

blackgum	<i>Nyssa sylvatica</i>
cherry	<i>Prunus</i> spp.
chestnut, Chinese	<i>Castanea mollissima</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
douglas fir	<i>Pseudotsuga menziesii</i>
fir, balsam	<i>Abies balsamea</i>
hemlock, Canada	<i>Tsuga canadensis</i>
honeylocust	<i>Gleditsia triacanthos</i>
larch, Japanese	<i>Larix kaempferi</i>
locust, black	<i>Robinia pseudoacacia</i>
maple, Norway	<i>Acer platanoides</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
maple, sugar	<i>Acer saccharum</i>
oak, pin	<i>Quercus palustris</i>
oak, scarlet	<i>Quercus coccinea</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, loblolly	<i>Pinus taeda</i>
pine, red	<i>Pinus resinosa</i>
pine, scotch	<i>Pinus sylvestris</i>
planetree, London	<i>Platanus acerifolia</i>
plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
spruce, white	<i>Picea glauca</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
tuliptree	<i>Liriodendron tulipifera</i>
walnut, black	<i>Juglans nigra</i>

Ornamental Shrubs

Common Name	Scientific Name
andromeda, Japanese	<i>Pieris japonica</i>
arborvitae, American	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
barberry, Japanese	<i>Berberis thunbergii</i>
barberry, mento	<i>Berberis mentorensis</i>
boxwood, common	<i>Buxus sempervirens</i>
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cherry laurel, American	<i>Prunus caroliniana</i>
cinquefoil	<i>Potentilla</i> spp.
cleysera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
cotoneaster, zabel	<i>Cotoneaster zabelii</i>
deutzia	<i>Deutzia</i> spp.
elaegnus, silverberry	<i>Elaeagnus pungens</i>
euonymus, spreading	<i>Euonymus kiautschovica</i>

16/31

euonymus, winged	<i>Euonymus alatus</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
firethorn	<i>Pyracantha</i> spp.
forsythia	<i>Forsythia</i> spp.
guava, pineapple	<i>Feijoa sellowiana</i>
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
India hawthorn	<i>Raphiolepis indica</i>
juniper	<i>Juniperus</i> spp.
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
mockorange	<i>Philadelphus</i> spp.
pittosporum, Japanese	<i>Pittosporum tobira</i>
privet	<i>Ligustrum</i> spp.
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spiraea, vanhoutte	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
weigela	<i>Weigela</i> spp.
willow	<i>Salix</i> spp.
yew, anglojap	<i>Taxus media</i>
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

Ornamental Groundcover Plantings

Common Name	Scientific Name
aaronsbeard	<i>Hypericum calycinum</i>
bellflower, adriatic	<i>Campanula elatines</i>
bellflower, poscharsky	<i>Campanula poscharskyana</i>
ceanothus	<i>Ceanothus</i> spp.
coreopsis	<i>Coreopsis</i> spp.
cotoneaster	<i>Cotoneaster</i> spp.
coyote brush	<i>Baccharis pilularis</i>
crown vetch	<i>Coronilla vana</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
fern, asparagus	<i>Asparagus densiflorus</i>
gazania	<i>Gazania</i> spp.
germander	<i>Teucrium chamaedrys</i>
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lily-of-the-nile	<i>Agapanthus</i> spp.
lilyturf, bigblue	<i>Liriope muscari</i>
marigold	<i>Tagetes</i> spp.
myoporum	<i>Myoporum laetum</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rockrose	<i>Cistus</i> spp.
rosemary	<i>Rosmarinus officinalis</i>
rupturewort	<i>Herniaria glabra</i>
snow-in-summer	<i>Cerastium tomentosum</i>
speedwell	<i>Veronica</i> spp.
St. Johnswort	<i>Hypericum coris</i>
stonecrop (sedum)	<i>Sedum</i> spp.
strawberry, beach	<i>Fragaria chiloensis</i>

thrift	<i>Armeria maritima</i>
verbena	<i>Verbena</i> spp.
wirevine, creeping	<i>Muehlenbeckia axillaris</i>
yarrow, wooly	<i>Achillea tomentosa</i>

Roses and Other Established Flowers

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Common Name	Scientific Name
Annual Grasses	
annual bluegrass	<i>Poa annua</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
crabgrass, large or hairy	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
goosegrass (silver crabgrass or crowfoot)	<i>Eleusine indica</i>
green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria glauca</i>
Broadleaf Weeds	
chickweed, common	<i>Stellaria media</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
pigweed	<i>Amaranthus</i> spp.
purslane, common	<i>Portulaca oleracea</i>

Weeds Suppressed

Weeds suppressed when applied prior to weed germination:

Common Name	Scientific Name
creeping spurge	<i>Euphorbia serpens</i>
creeping woodsorrel	<i>Oxalis corniculata</i>
panicum, Texas	<i>Panicum texanum</i>
prostrate spurge	<i>Euphorbia humistrata</i>
pusley, Florida	<i>Richardia scabra</i>
spotted spurge	<i>Euphorbia maculata</i>
woodsorrel	<i>Oxalis</i> spp.

Application Directions

18/31

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	XXX	XXX	XXX	XXX	XXX
XXXXXXXX	XX	XX	XX	XX	XX

Uses

Established Turfgrass

Application Timing

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	174 to 349lb/acre 4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)
winter annual grass weeds (fall application)	174 to 349 lb/acre 4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 233 lb plus 233 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 349 lb of Turf Fertilizer – Contains Team Pro 0.86% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.

19/31

- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/gfbmp07.pdf>.
- Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

Broadcast Application Rates

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 8 lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 8 lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

20/31

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™® Trademark of Dow AgroSciences LLC
EPA accepted 12/10/07

[Sub Label B: Residential Uses]

(Base label):

Turf Fertilizer - Contains Team[®] Pro 0.86%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

and/or

ONE STEP – extended grassy weed control

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

22/31

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.43%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.43%
Other Ingredients	99.14%
Total	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

Keep Out of Reach of Children

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, shoes plus socks and goggles or safety glasses.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. _____

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight _____

(Label booklet cover):

Turf Fertilizer - Contains Team[®] Pro 0.86%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

and/or

ONE STEP – extended grassy weed control

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

25/31

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass] and/or

University tested [and researched] and/or

Labeled for use around a [wide] variety of trees and shrubs and/or

Never see crabgrass again

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.43%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.43%
Other Ingredients	99.14%
Total	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.X%
X.X% Nitrate Nitrogen	
X.X% Water Soluble Nitrogen	
X.X% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	X.X%
Soluble Potash (K ₂ O)	X.X%

Keep Out of Reach of Children

WARNING

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-289

EPA Est. No. _____

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268

Net Weight _____

26/31

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING**

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, shoes plus socks and goggles or safety glasses.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic

27/31

modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

How This Product Works

This product is a granular fertilizer containing Team[®], a preemergence herbicide for application to established home lawns and landscape ornamentals (trees, shrubs, groundcovers and flowers) for control of annual grasses and broadleaf weeds.

General Use Precautions and Restrictions

- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 4 lb of product per 1000 sq ft, or 12 to 16 weeks after a single application of 8 lb of product per 1000 sq ft unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 4 lb of product per 1000 sq ft rate is applied, or 10 to 12 weeks after the last 5.34 lb of product per 1000 sq ft rate is applied unless injury can be tolerated.

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass

bentgrass fairways
fescues
Kentucky bluegrass
perennial ryegrass

Established Warm Season Turfgrass

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass
zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

almond	hemlock, Canada	pine, loblolly
apple, crabapple	honeylocust	pine, red
apricot	larch, Japanese	pine, scotch
ash, white	locust, black	planetree, London
bald cypress	maple, Norway	plum
birch, European white	maple, red	redbud, eastern
blackgum	maple, silver	spruce, Colorado

28/31

cherry	maple, sugar	spruce, Norway
chestnut, Chinese	oak, pin	spruce, white
cottonwood	oak, scarlet	sweetgum
dogwood, flowering	peach	sycamore
dogwood, kousa	pine, Austrian	tuliptree
douglas fir	pine, eastern white	walnut, black
fir, balsam	pine, Japanese black	

Ornamental Shrubs

andromeda, Japanese	forsythia
arborvitae, American	guava, pineapple
azalea	holly
barberry, Japanese	honeysuckle
barberry, mento	India hawthorn
boxwood, common	juniper
boxwood, littleleaf	laurel, mountain
camellia, Japanese	lilac, common
camellia, sasanqua	mockorange
cherrylaurel, American	pittosporum, Japanese
cinquefoil	privet
cleyera, Japanese	redcedar, eastern
cotoneaster, cranberry	rhododendron
cotoneaster, zabel	spirea, vanhoutte
deutzia	viburnum
elaeagnus, silverberry	weigela
euonymus, spreading	willow
euonymus, winged	yew, anglojap
euonymus, wintercreeper	yew, Japanese
firethorn	yewpine

Ornamental Groundcover Plantings

aaronsbeard	lilyturf, bigblue
bellflower, adriatic	marigold
bellflower, poscharsky	myoporum
ceanothus	plumbago, dwarf
coreopsis	rockrose
cotoneaster	rosemary
coyote brush	rupturewort
crown vetch	snow-in-summer
daisy trailing African	speedwell
fern, asparagus	St. Johnswort
gazania	stonecrop (sedum)
germander	strawberry, beach
ice plant, largeleaf	thrift
ivy, Algerian	verbena
ivy, English	wirevine, creeping
lily-of-the-nile	yarrow, wooly

Roses and Other Established Flowers

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum

29/31

centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Annual Grasses	Broadleaf Weeds
annual bluegrass	chickweed, common
barnyardgrass (watergrass)	knotweed, prostrate
crabgrass, large or hairy	lambsquarters, common
crabgrass, smooth	pigweed
goosegrass (silver crabgrass or crowfoot)	purslane, common
green foxtail	
yellow foxtail	

Weeds Suppressed

Weeds suppressed when applied prior to weed germination:

creeping spurge
 creeping woodsorrel
 panicum, Texas
 prostrate spurge
 pusley, Florida
 spotted spurge
 woodsorrel

How to Apply

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate the application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or spinkler irrigation of 0.5 inch or more is required to activate this product.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	XX	XX	XX	XX	XX
XXXXXXXX	XX	XX	XX	XX	XX

Uses

Established Home Lawns

When to Apply

Apply this product 1 to 2 weeks prior to germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

How Much to Apply

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	4 to 8 lb/1000 sq ft	8 lb/1000 sq ft
winter annual grass weeds (fall application)	4 to 8 lb/1000 sq ft	8 lb/1000 sq ft

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 4 lb plus 4 lb per 1000 sq ft for cool season turfgrass and in the northern half of the warm season turfgrass area, or 5.34 lb plus 5.34 lb per 1000 sq ft in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 4 lb per 1000 sq ft rate and 10 to 12 weeks following the 5.34 lb per 1000 sq ft rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
 - **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- [Editor’s note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

When to Apply

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

How to Apply

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control

31/31

performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™® Trademark of Dow AgroSciences LLC
EPA accepted 12/10/07

NOTIFICATION

MAR 30 2009