

62719-290

03/20/2003

1/13

P4A/Turf Fertilizer - Contains Team/ Amend With Edits / 12-09-02

Page 1

(Product Label)

(Logo) Dow AgroSciences

## Turf Fertilizer - Contains Team\* {X.XX<sup>(a)</sup>}%

**A Granular Fertilizer Containing Team for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, *Poa annua* and Other Annual Grass and Broadleaf Weeds in Established Turfgrasses [and established landscape ornamentals].**

ONE STEP - extended grassy weed control.


### Active Ingredient(s):

benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine ..... {X.XX (b)}%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine ..... {X.XX (c)}%

Inert Ingredients ..... {X.XX (d)}%

Total ..... 100.00%



### Guaranteed Fertilizer Analysis†

Total Nitrogen (N) ..... {X.XX}%

Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) ..... {X.XX}%

Soluble Potash (K<sub>2</sub>O) ..... {X.XX}%

[†Editor's note: Specific analysis of the fertilizer blend used to be provided by supplemental distributor.]

Contains [X.XX]†† pounds active ingredient per [XX] pound bag.

[††Editor's note: Bag size and calculated amount of active ingredient per bag provided by supplemental distributor.]

**Keep Out of Reach of Children**

## **CAUTION-WARNING**

See back or container/side panel for additional precautionary statements.

**Notice:** Read the entire label. Use only according to label directions. **Before buying using this product, read Limited Warranty and Disclaimer Section elsewhere on this label Inherent Risks of Use and Limitation of Remedies at end of Directions for Use section. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-290

EPA Est. No. \_\_\_\_\_

\*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

**Herbicide**

**Net Weight - [XX] Pounds**

## Precautionary Statements

### Hazards to Humans and Domestic Animals

## CAUTION-WARNING

~~Causes Moderate Eye Irritation~~ ~~Causes Substantial But Temporary Eye Injury~~ • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

~~Avoid breathing dust and contact with eyes, skin or clothing. Do not get in eyes or on clothing. Avoid contact with skin. The granular formulation may cause skin sensitization reactions in certain individuals. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Use protective clothing such as coveralls, a long sleeved shirt and long pants, shoes and socks, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.~~

### Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists. ~~Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.~~

If swallowed: Do not induce vomiting. Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol.

If on skin or clothing: Wash with plenty of soap and water. Get medical attention if irritation develops. Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth to mouth) and get medical attention immediately.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

## Environmental Hazards

~~This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sties. Do not contaminate water when disposing of equipment washwaters.~~

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

## Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
Read all directions carefully before applying this product.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated area (except those involved in the watering) until the watering-in is complete and the surface is dry.

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## General Information

This turf fertilizer product contains Team\*, a preemergence herbicide for control of annual grasses and broadleaf weeds in established perennial turfgrasses [and established landscape ornamentals].

## Weeds Controlled

### Annual Grasses:

#### Common Name

annual bluegrass

crabgrass, smooth

crabgrass, large or hairy

goosegrass

(silver crabgrass or crowfootgrass)

barnyardgrass (watergrass)

#### Scientific Name

*Poa annua*

*Digitaria ischaemum*

*Digitaria sanguinalis*

*Eleusine indica*

*Echinochloa crus-galli*

green foxtail	<i>Setaria viridis</i>
yellow foxtail	<i>Setaria glauca</i>

**Broadleaf Weeds:**

Common Name	Scientific Name
chickweed, common	<i>Stellaria media</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
pigweed	<i>Amaranthus spp.</i>
purslane, common	<i>Portulaca oleracea</i>

**Weeds Suppressed:**

Common Name	Scientific Name
panicum, Texas	<i>Panicum texanum</i>
pusley, Florida	<i>Richardia scabra</i>
prostrate spurge	<i>Euphorbia humistrata</i>
spotted spurge	<i>Euphorbia maculata</i>
creeping spurge	<i>Euphorbia serpens</i>
creeping woodsorrel	<i>Oxalis corniculata</i>
woodsorrel	<i>Oxalis spp.</i>

---

## Approved Uses

---

### Established Turfgrasses

**When to Apply:** For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons.

**How to Apply:** Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands.

**Activation:** Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

**Important Use Precautions:** Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

**This Product may be applied to the following established turfgrass species:**

**Established Cool Season**

**Turfgrasses**

Bentgrass fairways  
Kentucky bluegrass  
Perennial ryegrass  
Fescues

**Established Warm Season**

**Turfgrasses**

Bahiagrass  
Bermudagrass  
Centipedegrass  
Fescue, tall  
St. Augustinegrass  
Zoysiagrass

**Broadcast Application Rates:**

Application Timing	Application Rates	Specific Use Directions
<b>Spring:</b> To control spring-to-summer germinating annual weeds. <b>Fall:</b> To control fall-germinating weeds.	{X.X (e)} to {X.X(g)} lb per 1000 sq ft	In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination.
<b>Split Application:</b> For extended control of spring and summer annual weeds.	{X.X (e)} lb per 1000 sq ft plus a second application of {X.X (e)} lb per 1000 sq ft	In Cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later.
	{X.X (f)} lb per 1000 sq ft plus a second application of {X.X (f)} lb per 1000 sq ft	In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.

**Reseeding Precautions**

Team herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application of this product, should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of Product Applied (Pounds per 1000 sq ft)	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of {X.X(e)} lb/1000 sq ft	8
Single application {X.X(g)} lb/1000 sq ft	12 to 16
Split application of {X.X(e)} lb/1000 sq ft	8 to 10
Split application of {X.X(f)} lb/1000 sq ft	10 to 12

**[Spreader Settings]**

**[Editors note: Formulator label may include a table of recommended spreader equipment and spreader settings for the fertilizer blend used.]**

**[Editors note: The following section is optional if the product is to be marketed strictly as a turf fertilizer.]**

### **Established Ornamental Plants in Landscape Settings**

This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers.

**Application timing:** Apply one to two weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet.

**Soil Preparation:** This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

**Application Equipment and Application Techniques:** See General Information section.

**Activation:** Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall, within a few hours after application.

#### **Broadcast Application Rates:**

Season of Application	Pounds per 1000 sq ft
Spring Application (Summer annuals)	{X.X(e)} to {X.X(g)} lb/1000 sq ft
Fall Application (Winter annuals)	{X.X(e)} to {X.X(g)} lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 - 12 weeks following the initial application.

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

#### **Ornamental Trees:**

Common Name	Scientific Name
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
ash, white	<i>Fraxinus americana</i>
bald cypress	<i>Taxodium distichum</i>
birch, European white	<i>Betula pendula</i>
blackgum	<i>Nyssa sylvatica</i>
cherry	<i>Prunus</i> spp.
chestnut, Chinese	<i>Castanea mollissima</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
douglasfir	<i>Pseudotsuga menziesii</i>
fir, balsam	<i>Abies balsamea</i>
hemlock, Canada	<i>Tsuga canadensis</i>
honeylocust	<i>Gleditsia triacanthos</i>
larch, Japanese	<i>Larix kaempferi</i>
locust, black	<i>Robinia pseudoacacia</i>
maple, Norway	<i>Acer platanoides</i>

maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
maple, sugar	<i>Acer saccharum</i>
oak, pin	<i>Quercus palustris</i>
oak, scarlet	<i>Quercus coccinea</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, loblolly	<i>Pinus taeda</i>
pine, red	<i>Pinus resinosa</i>
pine, scotch	<i>Pinus sylvestris</i>
planetree, London	<i>Platanus acerifolia</i>
plum	<i>Prunus spp.</i>
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
spruce, white	<i>Picea glauca</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
tuliptree	<i>Liriodendron tulipifera</i>
walnut, black	<i>Juglans nigra</i>

### Ornamental Shrubs:

Common Name	Scientific Name
andromeda, Japanese	<i>Pieris japonica</i>
arborvitae, American	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron spp.</i>
barberry, Japanese	<i>Berberis thunbergii</i>
barberry, mento	<i>Berberis mentorensis</i>
boxwood, common	<i>Buxus sempervirens</i>
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cherry laurel, American	<i>Prunus caroliniana</i>
cinquefoil	<i>Potentilla spp.</i>
cleysera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
cotoneaster, zabel	<i>Cotoneaster zabelii</i>
deutzia	<i>Deutzia spp.</i>
elaeagnus, silverberry	<i>Elaeagnus pungens</i>
euonymus, spreading	<i>Euonymus kiautschovica</i>
euonymus, winged	<i>Euonymus alatus</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
firethorn	<i>Pyracantha spp.</i>
forsythia	<i>Forsythia spp.</i>
guava, pineapple	<i>Feijoa sellowiana</i>
holly	<i>Ilex spp.</i>
honeysuckle	<i>Lonicera spp.</i>
India hawthorn	<i>Raphiolepis indica</i>
juniper	<i>Juniperus spp.</i>
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>



9/13

mockorange	<i>Philadelphus</i> spp.
pittosporum, Japanese	<i>Pittosporum tobira</i>
privet	<i>Ligustrum</i> spp.
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spirea, vanhoutte	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
weigela	<i>Weigela</i> spp.
willow	<i>Salix</i> spp.
yew, anglojap	<i>Taxus media</i>
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

### Ornamental Groundcover Plantings:

Common Name	Scientific Name
aaronsbeard	<i>Hypericum calycinum</i>
bellflower, adriatic	<i>Campanula elatines</i>
bellflower, poscharsky	<i>Campanula poscharskyana</i>
ceanothus	<i>Ceanothus</i> spp.
coreopsis	<i>Coreopsis</i> spp.
cotoneaster	<i>Cotoneaster</i> spp.
coyote brush	<i>Baccharis pilularis</i>
crown vetch	<i>Coronilla vana</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
fern, asparagus	<i>Asparagus densiflorus</i>
gazania	<i>Gazania</i> spp.
germander	<i>Teucrium chamaedrys</i>
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lily-of-the-nile	<i>Agapanthus</i> spp.
lilyturf, bigblue	<i>Liriope muscari</i>
marigold	<i>Tagetes</i> spp.
myoporum	<i>Myoporum laetum</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rockrose	<i>Cistus</i> spp.
rosemary	<i>Rosmarinus officinalis</i>
rupturewort	<i>Herniaria glabra</i>
snow-in-summer	<i>Cerastium tomentosum</i>
speedwell	<i>Veronica</i> spp.
st. johnswort	<i>Hypericum coris</i>
stonecrop (sedum)	<i>Sedum</i> spp.
strawberry, beach	<i>Fragaria chiloensis</i>
thrift	<i>Armeria maritima</i>
verbena	<i>Verbena</i> spp.
wirevine, creeping	<i>Muehlenbeckia axillaris</i>
yarrow, woolly	<i>Achillea tomentosa</i>

### Roses and Other Established Flowers:

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain

blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

---

### **Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### **Warranty Disclaimer**

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC

EPA-accepted   /  /

## Appendix 1.

**Appendix 1 and the product CSF provide information necessary for preparation of final printed labeling and formulation of end-use product as follows:**

**Table 1** provides the appropriate numbers to use in the label for a selected end use product. Each column has been referenced with letters (a through g) which correspond to the areas on the label where these numbers should be inserted. Inserting these numbers provides the correct percent concentration for the active ingredients benefin and trifluralin, and application rates of the formulated turf fertilizer/herbicide product for each alternative end-use formulation.

### Example:

- A formulator desires to create a product containing a turf fertilizer that could be applied at 4 to 8 pounds of formulated product per 1000 square feet. By utilizing the table in appendix 1 the formulator would select a concentration of 0.86% (column a) which would deliver 1.5 pounds of Team per acre at the 4 pounds of formulated product per 1000 square feet rate (column e) or 3 pounds of Team per acre at 8 pounds of formulated product per 1000 square feet (column g). If a different rate of formulated product per acre or per 1000 square feet had been required then a different line on the table would be used.
- Using the line for Team at a total active ingredient concentration of 0.86%, the appropriate table values from columns a, b, c, d, e, f, and g would be inserted into final printed labeling as indicated on this label. Replacement occurs wherever "X.XX" appears in upper case brackets "{-}".
- Values that are provided and inserted into final printed labeling by the supplemental distributor are indicated by Editor's Notes and appear within lower case brackets "{-}".

**Confidential Statement of Formula (CSF):** Provides standard information (amounts, percent w/w, and certified limits) for preparation of each alternative end-use pesticide/turf fertilizer formulation.

**Table 1.** Active ingredient concentrations and application rates for formulated product corresponding to specified active ingredient application rates (a.i./acre) for Turf Fertilizer Contains Team\* herbicide.

Team Herbicide (%)	Benefin (%)	Trifluralin (%)	Granular Fertilizer (%)	Total (%)	Pounds of Formulated Product Per 1000 sq ft		
					Low Rate	Medium Rate	High Rate
					1.5 lb active/acre	2.0 lb active/acre	3 lb active/acre
(a)	(b)	(c)	(d)	---	(e)	(f)	(g)
1.25%	0.83%	0.42%	98.75%	100.00%	2.75	3.67	5.51
1.24%	0.83%	0.41%	98.76%	100.00%	2.78	3.70	5.55
1.23%	0.82%	0.41%	98.77%	100.00%	2.80	3.73	5.60
1.22%	0.82%	0.40%	98.78%	100.00%	2.82	3.76	5.65
1.21%	0.81%	0.40%	98.79%	100.00%	2.85	3.79	5.69
1.20%	0.80%	0.40%	98.80%	100.00%	2.87	3.83	5.74
1.19%	0.80%	0.39%	98.81%	100.00%	2.89	3.86	5.79
1.18%	0.79%	0.39%	98.82%	100.00%	2.92	3.89	5.84
1.17%	0.78%	0.39%	98.83%	100.00%	2.94	3.92	5.89
1.16%	0.78%	0.38%	98.84%	100.00%	2.97	3.96	5.94
1.15%	0.77%	0.38%	98.85%	100.00%	2.99	3.99	5.99

(Table 1 Continued)

Team Herbicide (%)	Benefin (%)	Trifluralin (%)	Granular Fertilizer (%)	Total (%)	Pounds of Formulated Product Per 1000 sq ft		
					Low Rate	Medium Rate	High Rate
					1.5 lb active/acre	2.0 lb active/acre	3 lb active/acre
(a)	(b)	(c)	(d)	---	(e)	(f)	(g)
1.14%	0.76%	0.38%	98.86%	100.00%	3.02	4.03	6.04
1.13%	0.76%	0.37%	98.87%	100.00%	3.05	4.06	6.09
1.12%	0.75%	0.37%	98.88%	100.00%	3.07	4.10	6.15
1.11%	0.74%	0.37%	98.89%	100.00%	3.10	4.14	6.20
1.10%	0.74%	0.36%	98.90%	100.00%	3.13	4.17	6.26
1.09%	0.73%	0.36%	98.91%	100.00%	3.16	4.21	6.32
1.08%	0.72%	0.36%	98.92%	100.00%	3.19	4.25	6.38
1.07%	0.72%	0.35%	98.93%	100.00%	3.22	4.29	6.44
1.06%	0.71%	0.35%	98.94%	100.00%	3.25	4.33	6.50
1.05%	0.70%	0.35%	98.95%	100.00%	3.28	4.37	6.56
1.04%	0.70%	0.34%	98.96%	100.00%	3.31	4.41	6.62
1.03%	0.69%	0.34%	98.97%	100.00%	3.34	4.46	6.69
1.02%	0.68%	0.34%	98.98%	100.00%	3.38	4.50	6.75
1.01%	0.68%	0.33%	98.99%	100.00%	3.41	4.55	6.82
1.00%	0.67%	0.33%	99.00%	100.00%	3.44	4.59	6.89
0.99%	0.66%	0.33%	99.01%	100.00%	3.48	4.64	6.96
0.98%	0.66%	0.32%	99.02%	100.00%	3.51	4.69	7.03
0.97%	0.65%	0.32%	99.03%	100.00%	3.55	4.73	7.10
0.96%	0.64%	0.32%	99.04%	100.00%	3.59	4.78	7.17
0.95%	0.64%	0.31%	99.05%	100.00%	3.62	4.83	7.25
0.94%	0.63%	0.31%	99.06%	100.00%	3.66	4.88	7.33
0.93%	0.62%	0.31%	99.07%	100.00%	3.70	4.94	7.41
0.92%	0.62%	0.30%	99.08%	100.00%	3.74	4.99	7.49
0.91%	0.61%	0.30%	99.09%	100.00%	3.78	5.05	7.57
0.90%	0.60%	0.30%	99.10%	100.00%	3.83	5.10	7.65
0.89%	0.60%	0.29%	99.11%	100.00%	3.87	5.16	7.74
0.88%	0.59%	0.29%	99.12%	100.00%	3.91	5.22	7.83
0.87%	0.58%	0.29%	99.13%	100.00%	3.96	5.28	7.92
0.86%	0.58%	0.28%	99.14%	100.00%	4.00	5.34	8.01
0.85%	0.57%	0.28%	99.15%	100.00%	4.05	5.40	8.10
0.84%	0.56%	0.28%	99.16%	100.00%	4.10	5.47	8.20
0.83%	0.56%	0.27%	99.17%	100.00%	4.15	5.53	8.30
0.82%	0.55%	0.27%	99.18%	100.00%	4.20	5.60	8.40
0.81%	0.54%	0.27%	99.19%	100.00%	4.25	5.67	8.50
0.80%	0.54%	0.26%	99.20%	100.00%	4.30	5.74	8.61
0.79%	0.53%	0.26%	99.21%	100.00%	4.36	5.81	8.72
0.78%	0.52%	0.26%	99.22%	100.00%	4.41	5.89	8.83
0.77%	0.52%	0.25%	99.23%	100.00%	4.47	5.96	8.94
0.76%	0.51%	0.25%	99.24%	100.00%	4.53	6.04	9.06
0.75%	0.50%	0.25%	99.25%	100.00%	4.59	6.12	9.18

13/13

(Table 1 Continued)

Team Herbicide (%)	Benefin (%)	Trifluralin (%)	Granular Fertilizer (%)	Total (%)	Pounds of Formulated Product Per 1000 sq ft		
					Low Rate	Medium Rate	High Rate
					1.5 lb active/acre	2.0 lb active/acre	3 lb active/acre
(a)	(b)	(c)	(d)	—	(e)	(f)	(g)
0.74%	0.50%	0.24%	99.26%	100.00%	4.65	6.20	9.31
0.73%	0.49%	0.24%	99.27%	100.00%	4.72	6.29	9.43
0.72%	0.48%	0.24%	99.28%	100.00%	4.78	6.38	9.57
0.71%	0.48%	0.23%	99.29%	100.00%	4.85	6.47	9.70
0.70%	0.47%	0.23%	99.30%	100.00%	4.92	6.56	9.84
0.69%	0.46%	0.23%	99.31%	100.00%	4.99	6.65	9.98
0.68%	0.46%	0.22%	99.32%	100.00%	5.06	6.75	10.13
0.67%	0.45%	0.22%	99.33%	100.00%	5.14	6.85	10.28
0.66%	0.44%	0.22%	99.34%	100.00%	5.22	6.96	10.43
0.65%	0.44%	0.21%	99.35%	100.00%	5.30	7.06	10.60
0.64%	0.43%	0.21%	99.36%	100.00%	5.38	7.17	10.76
0.63%	0.42%	0.21%	99.37%	100.00%	5.47	7.29	10.93
0.62%	0.42%	0.20%	99.38%	100.00%	5.55	7.41	11.11
0.61%	0.41%	0.20%	99.39%	100.00%	5.65	7.53	11.29
0.60%	0.40%	0.20%	99.40%	100.00%	5.74	7.65	11.48
0.59%	0.40%	0.19%	99.41%	100.00%	5.84	7.78	11.67
0.58%	0.39%	0.19%	99.42%	100.00%	5.94	7.92	11.87
0.57%	0.38%	0.19%	99.43%	100.00%	6.04	8.06	12.08
0.56%	0.38%	0.18%	99.44%	100.00%	6.15	8.20	12.30
0.55%	0.37%	0.18%	99.45%	100.00%	6.26	8.35	12.52
0.54%	0.36%	0.18%	99.46%	100.00%	6.38	8.50	12.75
0.53%	0.36%	0.17%	99.47%	100.00%	6.50	8.66	12.99
0.52%	0.35%	0.17%	99.48%	100.00%	6.62	8.83	13.24
0.51%	0.34%	0.17%	99.49%	100.00%	6.75	9.00	13.50
0.50%	0.33%	0.17%	99.50%	100.00%	6.89	9.18	13.77

\*Trademark of Dow AgroSciences LLC