


62719-290

12-10-2007

1/23

| | | |
|--|--------------------------------------|--------------------------------------|
|  U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460 | EPA Reg. Number: 62719-290 | Date of Issuance: 12-10-07 |
| | Term of Issuance: | |
| Name of Pesticide Product: Turf Fertilizer 0.86% | | |
| NOTICE OF PESTICIDE: <input type="checkbox"/> Registration <input checked="" type="checkbox"/> Reregistration (under FIFRA, as amended) | | |
| Name and Address of Registrant (include ZIP Code): Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 | | |
| Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number. | | |
| <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:</p> <ol style="list-style-type: none">1. Change the Signal Word to "Warning".2. To the First Aid section add "If On Skin Or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice."3. Change the Hazards to Humans and Domestic Animals statements to "Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing." | | |

Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.”

4. Change the PPE language to “All loaders, applicators, and other handlers must wear: Long sleeved shirt and long pants, Shoes plus socks, Chemical resistant gloves (such as or made of any waterproof material) and Goggles, face shield or safety glasses.”.
5. Change the Entry restriction language from “Do not enter or allow others to enter...” to “Do not enter or allow people or pets to enter...”.
6. To the label add “Watering in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing: This product must be watered in as soon as possible. Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering in is complete and the surface is dry.”.
7. The rate of “81 lbs of product per 1000 square feet” exceeds the maximum single application rates for benfluralin (see page 75 of the RED).
8. Delete “recommended for use on” from the label.
9. To the Warranty section change “Seller makes no” to “To the extent consistent with applicable law, seller makes no”, “in no case shall Seller” to “To the extent consistent with applicable law, in no case shall Seller” and “to the extent permitted by law” to “To the extent consistent with applicable law”.

Signature of Approving Official:



Kathryn V. Montague, Acting Product Manager (23)
Herbicide Branch, Registration Division (7505P)

Date:

12-10-07

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

3/23

(Base label for rigid containers 5 gal or less):

Turf Fertilizer - Contains Team^{TM*} ~~{X.XX}~~^(a) 0.86%**Herbicide**

A granular fertilizer containing Team_a for the preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

ONE STEP - extended grassy weed control.

Active Ingredients:

| | |
|---|---|
| benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine | {X.XX} ^(b) 0.57% |
| trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine | {X.XX} ^(c) 0.29% |
| Inert Other Ingredients | {X.XX} ^(d) 99.14% |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

| | |
|--|------|
| Total Nitrogen (N) | X.X% |
| Available Phosphoric Acid (P ₂ O ₅) | X.X% |
| Soluble Potash (K ₂ O) | X.X% |

Keep Out of Reach of Children**WARNING CAUTION****Precautionary Statements****Hazards to Humans and Domestic Animals**

Causes Moderate Eye Irritation Substantial But Temporary Eye Injury • Harmful If Swallowed
Absorbed Through The Skin

Avoid contact with eyes or clothing. Do not get in eyes or on clothing. Avoid contact with skin.
Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

ACCEPTED
 with COMMENTS
 in EPA Letter Dated

12-10-07

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

62719-290

4/23

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands ~~after handling and~~ before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

See Refer to label booklet back or container/side panel for additional precautionary statements Directions for Use.

5/23

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of Directions for Use section elsewhere on this label.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-290

EPA Est. No. _____

*TMTrademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Herbicide

Net Weight _____

6/23

(Base label for nonrigid containers any size):

Turf Fertilizer - Contains Team^{TM*} ~~{X.XX}~~^(a) 0.86%**Herbicide**

A granular fertilizer containing Team, a for the preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

ONE STEP - extended grassy weed control.

Active Ingredients:

| | |
|---|---|
| benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine | {X.XX} ^(b) 0.57% |
| trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine | {X.XX} ^(c) 0.29% |
| Other Ingredients | {X.XX} ^(d) 99.14% |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

| | |
|--|------|
| Total Nitrogen (N) | X.X% |
| Available Phosphoric Acid (P_2O_5) | X.X% |
| Soluble Potash (K_2O) | X.X% |

Keep Out of Reach of Children**WARNING CAUTION****Precautionary Statements****Hazards to Humans and Domestic Animals**

Causes Moderate Eye Irritation Substantial But Temporary Eye Injury • Harmful If Swallowed Absorbed Through The Skin

Avoid contact with eyes or clothing. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

7/23

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands ~~after handling and~~ before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

See Refer to label booklet back or container/side panel for additional precautionary statements Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of Directions for Use section elsewhere on this label.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-290

EPA Est. No. _____

*TM Trademark of Dow AgroSciences LLC

8/23

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Herbicide

Net Weight _____

9/23

(Base label for rigid containers greater than 5 gal):

Turf Fertilizer - Contains Team^{TM*} ~~{X.XX^(a)}~~ 0.86%**Herbicide**

A granular fertilizer containing Team, a for the preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

ONE STEP - extended grassy weed control.

Active Ingredients:

| | |
|--|--|
| benefin: N-butyl-N-ethyl- α,α,α -trifluoro- | |
| 2,6-dinitro-p-toluidine | {X.XX^(b)} 0.57% |
| trifluralin: α,α,α -trifluoro-2,6-dinitro- | |
| N,N-dipropyl-p-toluidine | {X.XX^(c)} 0.29% |
| Inert Other Ingredients | {X.XX^(d)} 99.14% |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

| | |
|--|------|
| Total Nitrogen (N) | X.X% |
| Available Phosphoric Acid (P ₂ O ₅) | X.X% |
| Soluble Potash (K ₂ O) | X.X% |

Keep Out of Reach of Children**WARNING CAUTION****Precautionary Statements****Hazards to Humans and Domestic Animals**

Causes Moderate Eye Irritation Substantial But Temporary Eye Injury • Harmful If Swallowed
Absorbed Through The Skin

Avoid contact with eyes or clothing. Do not get in eyes or on clothing. Avoid contact with skin.
Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or
using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

10/23

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands ~~after handling and~~ before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

See Refer to label booklet back or container/side panel for additional precautionary statements
Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of Directions for Use section elsewhere on this label.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

11/23

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-290

EPA Est. No. _____

[®][™] Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Herbicide

Net Weight _____

(Label booklet cover):

Turf Fertilizer - Contains Team^{TM*} ~~{X.XX}~~^(a) 0.86%

Herbicide

A granular fertilizer containing Team, a for the preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

ONE STEP - extended grassy weed control.

Active Ingredients:

| | |
|--|--|
| benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine | {X.XX} ^(b) <u>0.57%</u> |
| trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl- <i>p</i> -toluidine | {X.XX} ^(c) <u>0.29%</u> |
| Inert Other Ingredients | {X.XX} ^(d) <u>99.14%</u> |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

| | |
|--|------|
| Total Nitrogen (N) | X.X% |
| Available Phosphoric Acid (P ₂ O ₅) | X.X% |
| Soluble Potash (K ₂ O) | X.X% |

Keep Out of Reach of Children

WARNING CAUTION

See Refer to inside of label booklet back or container/side panel for additional precautionary information including statements Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of Directions for Use section elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-290

EPA Est. No. _____

*TMTrademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Herbicide

Net Weight _____

13/p3

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Causes Moderate Eye Irritation Substantial But Temporary Eye Injury • Harmful If Swallowed
Absorbed Through The Skin

Avoid contact with eyes or clothing. Do not get in eyes or on clothing. Avoid contact with skin.
Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or
using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- ~~Chemical-resistant gloves~~
- Shoes plus socks
- ~~Protective eyewear~~

~~User Safety Requirements:~~ Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands ~~after handling and before~~ eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when

14/03

disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, watering-in is required after application, do not enter or allow others to enter the treated area (except those involved in the watering) until the watering-in is complete and the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

15/23

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a granular turf fertilizer product containing Team[®]™, a preemergence herbicide for application to established perennial turfgrasses and established landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

General Use Precautions and Restrictions

Important Use Precautions:

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

Reseeding Precautions: Do not reseed treated areas for at least 8 weeks after a single application of 4 lb of product per 1000 sq ft, or 12 to 16 weeks after a single application of 8 lb of product per 1000 sq ft unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 4 lb of product per 1000 sq ft rate is applied, or 10 to 12 weeks after the last 5.34 lb of product per 1000 sq ft rate is applied unless injury can be tolerated. Team herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application of this product, should not be attempted until the applicable minimum time delay in the following table has been met:

This product is recommended for use on may be applied to the following established turfgrass species:

Established Cool Season Turfgrasses

bentgrass fairways
fescues
Kentucky bluegrass
perennial ryegrass

Established Warm Season Turfgrasses

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass
zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name

almond
apple, crabapple
apricot
ash, white
bald cypress
birch, European white
blackgum
cherry

Scientific Name

Prunus dulcis
Malus spp.
Prunus armeniaca
Fraxinus americana
Taxodium distichum
Betula pendula
Nyssa sylvatica
Prunus spp.

16/23

| | |
|----------------------|--------------------------------|
| chestnut, Chinese | <i>Castanea mollissima</i> |
| cottonwood | <i>Populus deltoides</i> |
| dogwood, flowering | <i>Cornus florida</i> |
| dogwood, kousa | <i>Cornus kousa</i> |
| douglas fir | <i>Pseudotsuga menziesii</i> |
| fir, balsam | <i>Abies balsamea</i> |
| hemlock, Canada | <i>Tsuga canadensis</i> |
| honeylocust | <i>Gleditsia triacanthos</i> |
| larch, Japanese | <i>Larix kaempferi</i> |
| locust, black | <i>Robinia pseudoacacia</i> |
| maple, Norway | <i>Acer platanoides</i> |
| maple, red | <i>Acer rubrum</i> |
| maple, silver | <i>Acer saccharinum</i> |
| maple, sugar | <i>Acer saccharum</i> |
| oak, pin | <i>Quercus palustris</i> |
| oak, scarlet | <i>Quercus coccinea</i> |
| peach | <i>Prunus persica</i> |
| pine, Austrian | <i>Pinus nigra</i> |
| pine, eastern white | <i>Pinus strobus</i> |
| pine, Japanese black | <i>Pinus thunbergiana</i> |
| pine, loblolly | <i>Pinus taeda</i> |
| pine, red | <i>Pinus resinosa</i> |
| pine, Scotch | <i>Pinus sylvestris</i> |
| planetree, London | <i>Platanus acerifolia</i> |
| plum | <i>Prunus</i> spp. |
| redbud, eastern | <i>Cercis canadensis</i> |
| spruce, Colorado | <i>Picea pungens</i> |
| spruce, Norway | <i>Picea abies</i> |
| spruce, white | <i>Picea glauca</i> |
| sweetgum | <i>Liquidambar styraciflua</i> |
| sycamore | <i>Platanus occidentalis</i> |
| tuliptree | <i>Liriodendron tulipifera</i> |
| walnut, black | <i>Juglans nigra</i> |

Ornamental Shrubs

| Common Name | Scientific Name |
|-------------------------|-------------------------------|
| andromeda, Japanese | <i>Pieris japonica</i> |
| arborvitae, American | <i>Thuja occidentalis</i> |
| azalea | <i>Rhododendron</i> spp. |
| barberry, Japanese | <i>Berberis thunbergii</i> |
| barberry, mento | <i>Berberis mentorensis</i> |
| boxwood, common | <i>Buxus sempervirens</i> |
| boxwood, littleleaf | <i>Buxus microphylla</i> |
| camellia, Japanese | <i>Camellia japonica</i> |
| camellia, sasanqua | <i>Camellia sasanqua</i> |
| cherrylaurel, American | <i>Prunus caroliniana</i> |
| cinquefoil | <i>Potentilla</i> spp. |
| cleyera, Japanese | <i>Cleyera japonica</i> |
| cotoneaster, cranberry | <i>Cotoneaster apiculata</i> |
| cotoneaster, zabel | <i>Cotoneaster zabelii</i> |
| deutzia | <i>Deutzia</i> spp. |
| elaeagnus, silverberry | <i>Elaeagnus pungens</i> |
| euonymus, spreading | <i>Euonymus kiautschovica</i> |
| euonymus, winged | <i>Euonymus alatus</i> |
| euonymus, wintercreeper | <i>Euonymus fortunei</i> |

17/23

| | |
|-----------------------|--------------------------------|
| firethorn | <i>Pyracantha</i> spp. |
| forsythia | <i>Forsythia</i> spp. |
| guava, pineapple | <i>Feijoa sellowiana</i> |
| holly | <i>Ilex</i> spp. |
| honeysuckle | <i>Lonicera</i> spp. |
| India hawthorn | <i>Raphiolepis indica</i> |
| juniper | <i>Juniperus</i> spp. |
| laurel, mountain | <i>Kalmia latifolia</i> |
| lilac, common | <i>Syringa vulgaris</i> |
| mockorange | <i>Philadelphus</i> spp. |
| pittosporum, Japanese | <i>Pittosporum tobira</i> |
| privet | <i>Ligustrum</i> spp. |
| redcedar, eastern | <i>Juniperus virginiana</i> |
| rhododendron | <i>Rhododendron</i> spp. |
| spiraea, vanhoutte | <i>Spiraea vanhouttei</i> |
| viburnum | <i>Viburnum</i> spp. |
| weigela | <i>Weigela</i> spp. |
| willow | <i>Salix</i> spp. |
| yew, anglojap | <i>Taxus media</i> |
| yew, Japanese | <i>Taxus cuspidata</i> |
| yewpine | <i>Podocarpus macrophyllus</i> |

Ornamental Groundcover Plantings

| Common Name | Scientific Name |
|------------------------|----------------------------------|
| aaronsbeard | <i>Hypericum calycinum</i> |
| bellflower, adriatic | <i>Campanula elatines</i> |
| bellflower, poscharsky | <i>Campanula poscharskyana</i> |
| ceanothus | <i>Ceanothus</i> spp. |
| coreopsis | <i>Coreopsis</i> spp. |
| cotoneaster | <i>Cotoneaster</i> spp. |
| coyote brush | <i>Baccharis pilularis</i> |
| crown vetch | <i>Coronilla vana</i> |
| daisy trailing African | <i>Osteospermum fruticosum</i> |
| fern, asparagus | <i>Asparagus densiflorus</i> |
| gazania | <i>Gazania</i> spp. |
| germander | <i>Teucrium chamaedrys</i> |
| ice plant, largeleaf | <i>Carpobrotus edulis</i> |
| ivy, Algerian | <i>Hedera canariensis</i> |
| ivy, English | <i>Hedera helix</i> |
| lily-of-the-nile | <i>Agapanthus</i> spp. |
| lilyturf, bigblue | <i>Liriope muscari</i> |
| marigold | <i>Tagetes</i> spp. |
| myoporum | <i>Myoporum laetum</i> |
| plumbago, dwarf | <i>Ceratostigma umbaginoides</i> |
| rockrose | <i>Cistus</i> spp. |
| rosemary | <i>Rosmarinus officinalis</i> |
| rupturewort | <i>Herniaria glabra</i> |
| snow-in-summer | <i>Cerastium tomentosum</i> |
| speedwell | <i>Veronica</i> spp. |
| St. Johnswort | <i>Hypericum coris</i> |
| stonecrop (sedum) | <i>Sedum</i> spp. |
| strawberry, beach | <i>Fragaria chiloensis</i> |
| thrift | <i>Armeria maritima</i> |
| verbena | <i>Verbena</i> spp. |
| wirevine, creeping | <i>Muehlenbeckia axillaris</i> |

18/23

yarrow, wooly

*Achillea tomentosa***Roses and Other Established Flowers**

| | | | |
|-------------------|---------------|-------------------|----------------------|
| African daisy | dianthus | marigold | shasta daisy |
| aster (perennial) | dusty miller | marigold, cape | snapdragon |
| balsam | floss flower | morningglory | snow-on-the-mountain |
| blackeyed susan | forget-me-not | nasturtium | stock |
| calendula | four o'clock | nicotiana | sunflower |
| carnation | gaillardia | petunia | sweet alyssum |
| centaurea, velvet | gladiolus | phlox | sweet pea |
| chrysanthemum | golden glow | pincushion flower | sweet sultan |
| coreopsis | impatiens | poppy, California | sweet william |
| cornflower | ixora | portulaca | vinca |
| cosmos | lobelia | rose | yarrow |
| dahlia | lupine | salvia | zinnia |

Weeds Controlled**Weeds controlled when applied prior to weed germination:**

| Common Name | Scientific Name |
|--|-------------------------------|
| Annual Grasses | |
| annual bluegrass | <i>Poa annua</i> |
| barnyardgrass (watergrass) | <i>Echinochloa crus-galli</i> |
| crabgrass, large or hairy | <i>Digitaria sanguinalis</i> |
| crabgrass, smooth | <i>Digitaria ischaemum</i> |
| goosegrass (silver crabgrass or crowfootgrass) | <i>Eleusine indica</i> |
| green foxtail | <i>Setaria viridis</i> |
| yellow foxtail | <i>Setaria glauca</i> |
| Broadleaf Weeds | |
| chickweed, common | <i>Stellaria media</i> |
| knotweed, prostrate | <i>Polygonum aviculare</i> |
| lambsquarters, common | <i>Chenopodium album</i> |
| pigweed | <i>Amaranthus</i> spp. |
| purslane, common | <i>Portulaca oleracea</i> |

Weeds Suppressed**Weeds suppressed when applied prior to weed germination:**

| Common Name | Scientific Name |
|---------------------|-----------------------------|
| creeping spurge | <i>Euphorbia serpens</i> |
| creeping woodsorrel | <i>Oxalis corniculata</i> |
| panicum, Texas | <i>Panicum texanum</i> |
| prostrate spurge | <i>Euphorbia humistrata</i> |
| pusley, Florida | <i>Richardia scabra</i> |
| spotted spurge | <i>Euphorbia maculata</i> |
| woodsorrel | <i>Oxalis</i> spp. |

Application Directions

How to Apply: Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment the spreader prior to use according to the manufacturer's directions. Check frequently to be make sure equipment is

19/23

working properly and distributing granules uniformly. ~~Avoid~~ Do not use of spreaders that apply material in narrow concentrated bands. Avoid skips or ~~and excessive~~ overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

| Brand and Model of Spreader | Calibration Settings at Specified Rate (lb/acre) | | | | |
|-----------------------------|--|-----|-----|-----|-----|
| | XXX | XXX | XXX | XXX | XXX |
| XXXXXXXX | XX | XX | XX | XX | XX |

Approved Uses

Established Turfgrass

Application Timing

When to Apply: Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. ~~For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons~~

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates

| | Cool Season Turfgrass | Warm Season Turfgrass |
|--|-----------------------------|------------------------|
| <u>summer annual grass weeds</u> (spring application) | <u>4 to 8 lb/1000 sq ft</u> | <u>8 lb/1000 sq ft</u> |
| <u>winter annual grass weeds</u> (fall application) | <u>4 to 8 lb/1000 sq ft</u> | <u>8 lb/1000 sq ft</u> |

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 4 lb plus 4 lb per 1000 sq ft for cool season turfgrass and in the northern half of the warm season turfgrass area, or 5.34 lb plus 5.34 lb per 1000 sq ft in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 4 lb per 1000 sq ft rate and 10 to 12 weeks following the 5.34 lb per 1000 sq ft rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 81 lb of Turf Fertilizer – Contains Team 0.86% per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

Activation: Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

| Application Timing | Application Rates | Specific Use Directions |
|--------------------|-------------------|-------------------------|
|--------------------|-------------------|-------------------------|

20/23

| | | |
|---|--|---|
| Spring: To control spring-to-summer germinating annual weeds. Fall: To control fall-germinating weeds. | {X.X (e)} to {X.X (g)} lb per 1000 sq ft | In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination. |
| Split Application: For extended control of spring and summer annual weeds. | {X.X (e)} lb per 1000 sq ft plus a second application of {X.X (e)} lb per 1000 sq ft | In cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later. |
| | {X.X (f)} lb per 1000 sq ft plus a second application of {X.X (f)} lb per 1000 sq ft | In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later. |

| Amount of Product Applied (Pounds per 1000 sq ft) | Minimum Time Delay Following Last Application Before Reseeding (Weeks) |
|--|--|
| Single application of {X.X (e)} lb/1000 sq ft | 8 |
| Single application {X.X (g)} lb/1000 sq ft | 12 to 16 |
| Split application of {X.X (e)} lb/1000 sq ft | 8 to 10 |
| Split application of {X.X (f)} lb/1000 sq ft | 10 to 12 |

Established Ornamental Plants in Landscape Settings

This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers. Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing

Apply 1 to 2 weeks prior to the expected germination time of target annual weeds seeds or to clean cultivated and hoe to remove existing weeds before applying this product weed free areas. Do not apply this product to plantings when the foliage is wet. **Soil Preparation:** This product can be surface applied around established plants or on ornamental beds. Apply to the a smooth soil surface (no large clods) to assure ensure uniform distribution around established plants or on ornamental beds. Clean-cultivate and hoe to remove existing weeds before applying this product.

Application Equipment and Application Techniques: See General Information section.

Broadcast Application Rates

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. **Activation:** Optimum weed control performance will be obtained when an application is followed by sprinkler overhead irrigation or rainfall within a few hours after application.

| Season of Application | Pounds per 1000 sq ft |
|-------------------------------------|--------------------------------------|
| Spring Application (Summer annuals) | {X.X (e)} to {X.X (g)} lb/1000 sq ft |
| Fall Application (Winter annuals) | {X.X (e)} to {X.X (g)} lb/1000 sq ft |

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.

21/23

- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team 0.86% per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team 0.86% per 1000 sq ft.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, and Inherent Risks of Use, above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*™ Trademark of Dow AgroSciences LLC
EPA accepted ___/___/___

22/p3

(Table 1 Continued)

| Team Herbicide (%) | Benefin (%) | Trifluralin (%) | Granular Fertilizer (%) | Total (%) | Pounds of Formulated Product Per 1000 sq ft | | |
|--------------------------|----------------|--------------------|-------------------------------|-----------|--|-----------------------|---------------------|
| | | | | | Low Rate | Medium Rate | High Rate |
| | | | | | 1.5 lb active/acre | 2.0 lb active/acre | 3 lb active/acre |
| (a) | (b) | (c) | (d) | — | (e) | (f) | (g) |
| 1.14% | 0.76% | 0.38% | 98.86% | 100.00% | 3.02 | 4.03 | 6.04 |
| 1.13% | 0.76% | 0.37% | 98.87% | 100.00% | 3.05 | 4.06 | 6.09 |
| 1.12% | 0.75% | 0.37% | 98.88% | 100.00% | 3.07 | 4.10 | 6.15 |
| 1.11% | 0.74% | 0.37% | 98.89% | 100.00% | 3.10 | 4.14 | 6.20 |
| 1.10% | 0.74% | 0.36% | 98.90% | 100.00% | 3.13 | 4.17 | 6.26 |
| 1.09% | 0.73% | 0.36% | 98.91% | 100.00% | 3.16 | 4.21 | 6.32 |
| 1.08% | 0.72% | 0.36% | 98.92% | 100.00% | 3.19 | 4.25 | 6.38 |
| 1.07% | 0.72% | 0.35% | 98.93% | 100.00% | 3.22 | 4.29 | 6.44 |
| 1.06% | 0.71% | 0.35% | 98.94% | 100.00% | 3.25 | 4.33 | 6.50 |
| 1.05% | 0.70% | 0.35% | 98.95% | 100.00% | 3.28 | 4.37 | 6.56 |
| 1.04% | 0.70% | 0.34% | 98.96% | 100.00% | 3.31 | 4.41 | 6.62 |
| 1.03% | 0.69% | 0.34% | 98.97% | 100.00% | 3.34 | 4.46 | 6.69 |
| 1.02% | 0.68% | 0.34% | 98.98% | 100.00% | 3.38 | 4.50 | 6.75 |
| 1.01% | 0.68% | 0.33% | 98.99% | 100.00% | 3.41 | 4.55 | 6.82 |
| 1.00% | 0.67% | 0.33% | 99.00% | 100.00% | 3.44 | 4.59 | 6.89 |
| 0.99% | 0.66% | 0.33% | 99.01% | 100.00% | 3.48 | 4.64 | 6.96 |
| 0.98% | 0.66% | 0.32% | 99.02% | 100.00% | 3.51 | 4.69 | 7.03 |
| 0.97% | 0.65% | 0.32% | 99.03% | 100.00% | 3.55 | 4.73 | 7.10 |
| 0.96% | 0.64% | 0.32% | 99.04% | 100.00% | 3.59 | 4.78 | 7.17 |
| 0.95% | 0.64% | 0.31% | 99.05% | 100.00% | 3.62 | 4.83 | 7.25 |
| 0.94% | 0.63% | 0.31% | 99.06% | 100.00% | 3.66 | 4.88 | 7.33 |
| 0.93% | 0.62% | 0.31% | 99.07% | 100.00% | 3.70 | 4.94 | 7.41 |
| 0.92% | 0.62% | 0.30% | 99.08% | 100.00% | 3.74 | 4.99 | 7.49 |
| 0.91% | 0.61% | 0.30% | 99.09% | 100.00% | 3.78 | 5.05 | 7.57 |
| 0.90% | 0.60% | 0.30% | 99.10% | 100.00% | 3.83 | 5.10 | 7.65 |
| 0.89% | 0.60% | 0.29% | 99.11% | 100.00% | 3.87 | 5.16 | 7.74 |
| 0.88% | 0.59% | 0.29% | 99.12% | 100.00% | 3.91 | 5.22 | 7.83 |
| 0.87% | 0.58% | 0.29% | 99.13% | 100.00% | 3.96 | 5.28 | 7.92 |
| 0.86% | 0.58% | 0.28% | 99.14% | 100.00% | 4.00 | 5.34 | 8.01 |
| 0.85% | 0.57% | 0.28% | 99.15% | 100.00% | 4.05 | 5.40 | 8.10 |
| 0.84% | 0.56% | 0.28% | 99.16% | 100.00% | 4.10 | 5.47 | 8.20 |
| 0.83% | 0.56% | 0.27% | 99.17% | 100.00% | 4.15 | 5.53 | 8.30 |
| 0.82% | 0.55% | 0.27% | 99.18% | 100.00% | 4.20 | 5.60 | 8.40 |
| 0.81% | 0.54% | 0.27% | 99.19% | 100.00% | 4.25 | 5.67 | 8.50 |
| 0.80% | 0.54% | 0.26% | 99.20% | 100.00% | 4.30 | 5.74 | 8.61 |
| 0.79% | 0.53% | 0.26% | 99.21% | 100.00% | 4.36 | 5.81 | 8.72 |
| 0.78% | 0.52% | 0.26% | 99.22% | 100.00% | 4.41 | 5.89 | 8.83 |
| 0.77% | 0.52% | 0.25% | 99.23% | 100.00% | 4.47 | 5.96 | 8.94 |
| 0.76% | 0.51% | 0.25% | 99.24% | 100.00% | 4.53 | 6.04 | 9.06 |
| 0.75% | 0.50% | 0.25% | 99.25% | 100.00% | 4.59 | 6.12 | 9.18 |

23/23

(Table 1 Continued)

| Team Herbicide (%) | Benefin (%) | Trifluralin (%) | Granular Fertilizer (%) | Total (%) | Pounds of Formulated Product Per 1000 sq ft | | |
|--------------------------|----------------|--------------------|-------------------------------|-----------|--|-----------------------|---------------------|
| | | | | | Low Rate | Medium Rate | High Rate |
| | | | | | 1.5 lb active/acre | 2.0 lb active/acre | 3 lb active/acre |
| (a) | (b) | (c) | (d) | --- | (e) | (f) | (g) |
| 0.74% | 0.50% | 0.24% | 99.26% | 100.00% | 4.65 | 6.20 | 9.34 |
| 0.73% | 0.49% | 0.24% | 99.27% | 100.00% | 4.72 | 6.29 | 9.43 |
| 0.72% | 0.48% | 0.24% | 99.28% | 100.00% | 4.78 | 6.38 | 9.57 |
| 0.71% | 0.48% | 0.23% | 99.29% | 100.00% | 4.85 | 6.47 | 9.70 |
| 0.70% | 0.47% | 0.23% | 99.30% | 100.00% | 4.92 | 6.56 | 9.84 |
| 0.69% | 0.46% | 0.23% | 99.31% | 100.00% | 4.99 | 6.65 | 9.98 |
| 0.68% | 0.46% | 0.22% | 99.32% | 100.00% | 5.06 | 6.75 | 10.13 |
| 0.67% | 0.45% | 0.22% | 99.33% | 100.00% | 5.14 | 6.85 | 10.28 |
| 0.66% | 0.44% | 0.22% | 99.34% | 100.00% | 5.22 | 6.96 | 10.43 |
| 0.65% | 0.44% | 0.21% | 99.35% | 100.00% | 5.30 | 7.06 | 10.60 |
| 0.64% | 0.43% | 0.21% | 99.36% | 100.00% | 5.38 | 7.17 | 10.76 |
| 0.63% | 0.42% | 0.21% | 99.37% | 100.00% | 5.47 | 7.29 | 10.93 |
| 0.62% | 0.42% | 0.20% | 99.38% | 100.00% | 5.55 | 7.41 | 11.11 |
| 0.61% | 0.41% | 0.20% | 99.39% | 100.00% | 5.65 | 7.53 | 11.29 |
| 0.60% | 0.40% | 0.20% | 99.40% | 100.00% | 5.74 | 7.65 | 11.48 |
| 0.59% | 0.40% | 0.19% | 99.41% | 100.00% | 5.84 | 7.78 | 11.67 |
| 0.58% | 0.39% | 0.19% | 99.42% | 100.00% | 5.94 | 7.92 | 11.87 |
| 0.57% | 0.38% | 0.19% | 99.43% | 100.00% | 6.04 | 8.06 | 12.08 |
| 0.56% | 0.38% | 0.18% | 99.44% | 100.00% | 6.15 | 8.20 | 12.30 |
| 0.55% | 0.37% | 0.18% | 99.45% | 100.00% | 6.26 | 8.35 | 12.52 |
| 0.54% | 0.36% | 0.18% | 99.46% | 100.00% | 6.38 | 8.50 | 12.75 |
| 0.53% | 0.36% | 0.17% | 99.47% | 100.00% | 6.50 | 8.66 | 12.99 |
| 0.52% | 0.35% | 0.17% | 99.48% | 100.00% | 6.62 | 8.83 | 13.24 |
| 0.51% | 0.34% | 0.17% | 99.49% | 100.00% | 6.75 | 9.00 | 13.50 |
| 0.50% | 0.33% | 0.17% | 99.50% | 100.00% | 6.89 | 9.18 | 13.77 |

*Trademark of Dow AgroSciences LLC