

62719-292

03/08/2000

1/2

T6P / Success / Sec 3 Suppl Dried Beans & Peas / 01-04-00

**Success\***

EPA Reg. No. 62719-292

Final printed labeling based on EPA accepted copy dated December 20, 1999.

Section 3 supplemental labeling for foliar insect control in dried beans and peas.

ACCEPTED

MAR 8 2000

62719-292

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Success\*

EPA Reg. No. 62719-292

### For Foliar Insect Control in Dried Beans and Peas

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success\* Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

#### Directions for Use

Dried cultivars of *Lupinus* spp., *Phaseolus* spp. and *Pisum* spp., including but not limited to: lupins, kidney bean, lima bean, navy bean, pinto bean, tepary bean, adzuki bean, blackeyed pea, cowpea, crowder pea, chickpea, mungbean, field pea, pigeon pea and lentil.

#### Pests and Application Rates:

Pests	Application Rate <sup>†</sup>		
	Active Ingredient (lb/acre)	Product (fl oz/acre)	Acres per Gallon of Product
European corn borer (eggs and larvae)	0.047 - 0.094	3 - 6	43 - 21
Armyworms, corn earworm, loopers	0.062 - 0.094	4 - 6	32 - 21
Thrips, leafminers	0.094	6	21

<sup>†</sup>Use the higher end of the rate range for larger larvae or high infestations.

#### Specific Use Directions:

**Application Timing:** To determine when to apply, scout with enough regularity to monitor populations of labeled pests. Treat when a labeled pest infestation first appears, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but make no more than 6 applications per crop. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations for your area.

#### Restrictions:

- Do not apply more than 12 fl oz ounces (0.186 lb of spinosad) of Success per acre per season.
- Do not feed treated forage or hay to meat or dairy animals.
- **Pre-harvest Interval:** Do not apply within 28 days of harvest.

\*Trademark of Dow AgroSciences LLC

T6P / Success / Sec 3 Suppl Dried Beans & Peas / 01-04-00

D06-065-009

EPA-accepted 12/20/1999

Initial printing.