


## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

62719-292

## Success\*

EPA Reg. No. 62719-292

### For Control of Lepidoptera Larvae Infesting Caneberries

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success\* Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

#### Directions for Use

Success may be used to control lepidoptera larvae infesting caneberries including, but not limited to, blackberry, loganberry, red and black raspberry, and cultivars and/or hybrids of these.

Refer to product label for Success for General Use Precautions, Mixing and Application Instructions.

#### Pests and Application Rates:

Pests	Rate (fl oz/acre)	Recommendations and Restrictions
beet armyworm bertha armyworm leafrollers green fruitworm western raspberry fruitworm looper sawfly	4 - 6	<p><b>Rate Range:</b> Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.</p> <p><b>Use Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>Pre-harvest Interval:</b> Do not apply within 1 day of harvest.</li> <li>• Do not apply more than a total of 29 fl oz of Success (0.45 lb ai of spinosad) per acre per crop or make more than 6 applications per calendar year.</li> <li>• <b>Minimum Treatment Interval:</b> Do not make applications less than 5 days apart.</li> </ul>

#### Specific Use Directions:

**Application Timing:** Treat when pests appear, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Resistance Management:** Do not apply Success more than 3 times in any 30 day period. Whenever Success is applied three times in succession, this should be followed by no use of Success for a 30 day period or rotation to another insecticide class.

**Note:** This product is toxic to bees exposed to treatment for 3 hours following treatment and highly toxic to mollusks. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

\*Trademark of Dow AgroSciences LLC

T6P / Success / Prop Sec 3 Supp IR-4 Caneberries / 02-04-02  
D06-065-XXX  
EPA-accepted \_\_\_/\_\_\_/\_\_\_  
Initial printing.

**NEXT**

**LABEL**

62719-292

09/27/2002

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

SEP 27 2002

A. Sterett Robertson  
Regulatory Manager  
Regulatory Success-Americas  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: Supplemental Labeling for Foliar Insect Control in Root Vegetables  
Success (A.I. SPINOSAD)  
EPA Registration No. 62719-292  
Your submission dated July 2, 2002

Dear: Mr. Robertson:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable, subject to the comments listed below.. A stamped copy of the label is enclosed for your records.

1. Delete all references to dry bulb onion, garlic, and shallots. Tolerances have not been established for dry bulb onions

If you have any questions, please call Dr. William Sproat of my team at 703-308-8587.

Sincerely,

A handwritten signature in black ink, appearing to read "George T. LaRocca".

George T. LaRocca  
Product Manager 13  
Insecticide Branch  
Registration Division (7505C)

Enclosure

SEP 27 2002

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Success\*

EPA Reg. No. 62719-292

### Foliar Insect Control in Root Vegetables (Subgroup 1A<sup>†</sup>) and Dry Bulb Onion

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success\* Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

### Directions for Use

<sup>†</sup>Root vegetables (subgroup 1A) including, but not limited to: Black Salsify, Carrot, Celeriac, Chicory, Edible Burdock, Ginseng, Horseradish, Parsnip, Radish, Oriental Radish, Rutabaga, Salsify, Skirret, Spanish Salsify, Turnip, Turnip-Rooted Chervil, and Turnip-Rooted Parsley.

Refer to product label for Success for General Use Precautions, Mixing and Application instructions.

#### Pests and Application Rates:

Crop	Pests	Success (fl oz/acre)	Specific Use Restrictions
black salsify carrot chicory ginseng horseradish parsnip salsify skirret Spanish salsify turnip-rooted chervil turnip-rooted parsley	armyworms dipteran leafminers European corn borer fleabeetle loopers thrips <sup>†</sup>	3 - 6	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 21 fl oz of Success (0.33 lb ai of spinosad) per acre per crop or make more than 4 applications per calendar year.</li> </ul>
celeriac edible burdock Oriental radish radish rutabaga turnip			<ul style="list-style-type: none"> <li>• Do not apply more than a total of 18 fl oz of Success (0.28 lb ai of spinosad) per acre per crop or make more than 3 applications per calendar year.</li> </ul>
dry bulb onion garlic shallots (dry bulbs only)	armyworms thrips <sup>†</sup> (suppression only)		<ul style="list-style-type: none"> <li>• Do not apply more than a total of 21 fl oz of Success (0.33 lb ai of spinosad) per acre per crop or make more than 4 applications per calendar year.</li> </ul>

<sup>†</sup>Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing on the product label. Control in onions has been variable; high pressure directed sprays can assist leaf penetration of onion.

#### Specific Use Directions:

**Application Timing:** For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae.

Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in specified rate range for larger larvae or heavier infestations.

**General Use Restrictions:**

- **Minimum Treatment Interval:** Do not make applications less than 5 days apart.
- **Pre-harvest Interval:** Do not apply within 3 days of harvest.

**Note:** This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

\*Trademark of Dow AgroSciences LLC

T6P / Success / Prop Sec 3 IR-4 Root Veggies-Dry Bulb Onion / 06-21-02

D06-065-XXX

EPA-accepted   /  /  

Initial printing.

**NEXT**

**LABEL**

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Success\*

EPA Reg. No. 62719-292

### For Foliar Insect Control in Herbs and Spices

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success\* Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

#### Directions for Use

##### Herbs and Spices


**Including but not limited to:** Angelica, balm, basil, borage, burnet, camomile, catnip, chervil (dried), chive, chive (Chinese), clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dillweed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram, nasturtium, parsley (dried) pennyroyal, rosemary, rue, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood.

Refer to product label for Success for General Use Precautions, Mixing and Application instructions..

##### Pests and Application Rates:

Pests	Application Rate †	
	Active Ingredient (lb/acre)	Product (fl oz/acre)
loopers armyworms	0.062 - 0.094	4 - 6

† Use the higher end of the rate range for larger larvae or high infestations.



##### Specific Use Directions:

**Application Timing:** For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae.

Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Apply Success as a foliar spray at the rate indicated to control target pests. Heavy infestations may require repeat applications, but follow resistance management guidelines.


**Resistance Management:** For resistance management purposes, do not apply more than 3 times in any 30-day period. Rotate to a different class of insect control products or use no treatment for the next 30 days. Do not make more than 4 applications per crop.



**Restrictions:**

- Do not apply more than 5 applications per crop or a total of 30 ounces (0.47 lb ai of spinosad) of Success per acre per crop.
- **Pre-harvest Interval:** Do not apply within 1 days of harvest.

<p>*Trademark of Dow AgroSciences LLC</p> <p>T6P / Success / Prop Sec 3 Suppl Herbs and Spices / 09-25-01</p> <p>D06-065-XXX</p> <p>EPA-accepted __/__/__</p> <p>Initial printing.</p>	
--	--



**NEXT**

**LABEL**

62719-292

09/27/2002

1/1

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Success\*

EPA Reg. No. 62719-292

### For Control of Lepidoptera Larvae and Thrips Infesting Grapes

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success\* Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application Instructions.

#### Pests and Application Rates:

Pests	Rate (fl oz/acre)	Recommendations and Restrictions
cutworm grape leaffolder grape leaf skeletonizer omnivorous leafroller orange tortrix thrips	4 - 8	<p><b>Rate Range:</b> Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.</p> <p><b>Use Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>Pre-harvest Interval:</b> Do not apply within 7 days of harvest.</li> <li>• Do not apply more than a total of 29 fl oz of Success (0.45 lb ai of spinosad) per acre per crop.</li> <li>• <b>Minimum Treatment Interval:</b> Do not make applications less than 5 days apart.</li> </ul>

**Specific Use Directions:** Equipment and spray volume should be carefully adjusted to assure thorough uniform coverage of infested parts of the crop.

**Application Timing:** Treat when pests appear, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Resistance Management:** Do not apply Success more than 3 times in any 30 day period. Whenever Success is applied three times in succession, this should be followed by no use of Success for a 30 day period or rotation to another insecticide class.

**Note:** This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

**ACCEPTED**  
SEP 27 2002

*Trademark of Dow AgroSciences LLC T6P / Success / Prop Sec 3 Supp IR-4 Grape / 04-08-02 D06-065-XXX EPA-accepted <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> Initial printing.	Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-292
--	--