

PLEASE NOTE

This image contains more than one label approved for this product on this date.

62719-292

02/27/2008

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

February 27, 2008

Michael D. Culy, PhD,
Regulatory Manager
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Supplemental labeling to add new uses per approved IR-4 Tolerance Petition #6E7148
"Foliar Insect Suppression in Pineapple"

Entrust (EPA Reg. No. 62719-282)
✓ Success (EPA Reg. No. 62719-292)

Your Submissions dated November 17, 2006

Dear Dr. Culy:

The above referenced amendments, submitted in connection with registrations under FIFRA Section 3(c)(7)(B), as amended, are acceptable.

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate the supplemental labeling into each product's main labeling. See enclosed stamped labels for your records.

If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely,

A handwritten signature in black ink, appearing to read "G. LaRocca".

George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure:

2/3

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Success[®] EPA Reg. No. 62719-292

(For Distribution and Use Only in the State of Hawaii)

Foliar Insect Suppression in Pineapple

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

Directions for Use

Refer to product label for Success for General Use Precautions, Mixing and Application instructions.

Pests and Application Rates:

Pests	Success (fl oz/acre)
lepidopteran larvae such as: armyworms banana moth fruit borer caterpillar (<i>Thecia basilides</i>) <i>Gummosos-Batrachedra commosae</i> pineapple caterpillar pink cornworm sugarcane bud moth	4 - 6

ACCEPTED

FEB 27 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-292

Specific Use Directions:

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated to control target pests. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: For resistance management purposes, do not apply Success more than 3 times in any 30-day period. Whenever Success is applied up to 3 times in succession, this should be followed by no use of Success for a 30-day period or rotation to another insecticide class.

3/3

Restrictions:

- Do not apply more than a total of 29 fl oz of Success (0.45 lb ai of spinosad) per acre per year
- Do not make more than 6 applications per calendar year.
- **Minimum Re-treatment Interval:** Do not make applications less than 7 days apart.
- **Preharvest Interval:** Do not apply within 7 days of harvest.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R065-028

Accepted: __/__/__

Initial printing.

NEXT

LABEL

62719-292

02/27/2008

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

February 27, 2008

Michael D. Culy, PhD,
Regulatory Manager
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Supplemental labeling to add new uses per approved IR-4 Tolerance Petition #6E7068
"Foliar Insect Control in Herbs and Hops, Dried Cones"

Entrust (EPA Reg. No. 62719-282)

✓Success (EPA Reg. No. 62719-292)

SpinTor (EPA Reg. No. 62719-294)

Your Submissions dated April 19, 2006

Dear Dr. Culy:

The above referenced amendments, submitted in connection with registrations under FIFRA Section 3(c)(7)(B), as amended, are acceptable subject to the following comments:

- For each of the 3 labels, revise the supplemental labeling title statement "**Foliar Insect Control in Herbs and Hops**" to read "***Foliar Insect Control in Herbs and Hops, Dried Cones***".
- Under "Direction for Use" on each label, delete the phrase "**included but not limited to**" from the first line.
- For the Entrust label, revise the use restriction statement "**Do not apply more than a total of 7.5 oz of Entrust . . .**" to read "***Do not apply more than a total of 9.5 oz of Entrust . . .***" This product is 80% by weight active ingredient.

Supplemental Labeling

FEB 27 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
62719-292



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Success[®]

EPA Reg. No. 62719-292

Foliar Insect Control in Herbs[†] and Hops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

Directions for Use

[†]Herbs, including, but not limited to: Angelica, balm, basil, borage, burnet, camomile, catnip, chervil (dried), chive, chive (Chinese), clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dillweed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram, nasturtium, parsley (dried) pennyroyal, rosemary, rue, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood.

Refer to product label for Success for General Use Precautions, Mixing and Application instructions.

Pests and Application Rates:

Pests	Application Rate		Recommendations and Restrictions
	Active Ingredient (lb/acre)	Success (fl oz/acre)	
armyworms loopers thrips (suppression)	0.062 – 0.094	4 - 6	Rate Range: Use a higher rate in the rate range for larger larvae or high infestations and/or larger plant volume. Use Restrictions: <ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 1 day of harvest. • Do not apply more than a total of 30 fl oz of Success (0.47 lb ai of spinosad) per acre per crop or make more than 5 applications per crop.

Specific Use Directions:

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated to control target pests. Heavy infestations may require repeat applications, not less than 5 days apart, but follow resistance management guidelines.

Resistance Management: For resistance management purposes, do not apply more than 3 times in any 30-day period. Rotate to a different class of insect control products or use no treatment for the next 30 days. Do not make more than 4 applications per crop.

3/3

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R065-027

Accepted: / /

Initial printing.

NEXT

LABEL

62719-292

02/27/2008

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

February 27, 2008

Michael D. Culy, PhD,
Regulatory Manager
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Supplemental labeling to add new uses per approved IR-4 Tolerance Petition #6E7148 "*Control of Lepidopteran Larvae and suppression of Flea Beetles, Dipteran Leafminers and Thrips in Spices (Except Black Pepper) Subgroup 19B*"

- Entrust (EPA Reg. No. 62719-282)
- ✓ Success (EPA Reg. No. 62719-292)
- SpinTor (EPA Reg. No. 62719-294)

Your Submissions dated November 17, 2006

Dear Dr. Culy:


The above referenced amendments, submitted in connection with registrations under FIFRA Section 3(c)(7)(B), as amended, are acceptable subject to the following comment:

- For each of the 3 labels, under "Direction for Use, first line", delete the phrase "*included but not limited to*".

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate the supplemental labeling into each product's main labeling. See enclosed stamped labels for your records.

If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely,


George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure:

Supplemental Labeling



2/3

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Success[®]

EPA Reg. No. 62719-292

Control of Lepidopteran Larvae and Suppression of Flea Beetles, Dipteran Leafminers and Thrips in Spices[†] (Except Black Pepper) (Subgroup 19B)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Success[®] Naturallyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Success according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Success.

Directions for Use

[†]Spices, including, but not limited to: allspice, anise (seed), annatto (seed), black caraway, caper (buds), caraway, cardamom, cassia (buds), celery (seed), cinnamon, clove (buds), common fennel, coriander (seed), culantro (seed), cumin, dill (seed), Florence fennel (seed), fenugreek, grains of paradise, juniper (berry), lovage (seed), mace, mustard (seed), nutmeg, poppy (seed), saffron, star anise, vanilla, white pepper

Refer to product label for Success for General Use Precautions, Mixing and Application instructions.

Pests and Application Rates:

Pests	Success (fl oz/acre)	Restrictions
lepidopteran larvae	4 - 6	• Preharvest Interval: Do not apply within 14 days of harvest. • Re-treatment Interval: 10 days • Do not apply more than a total of 29 fl oz of Success (0.45 lb ai of spinosad) per acre per crop or make more than 5 applications per calendar year.
flea beetles dipteran leafminers [†] thrips [†]	6 - 10	

[†]Suppression of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing.

Specific Use Directions:

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Heavy infestations may require repeat applications, but follow resistance management guidelines.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

FEB 27 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

62719-292

Resistance Management: For resistance management purposes, do not apply Success more than 3 times in any 30-day period or apply sprays less than 10 days apart. Whenever Success is applied up to 3 times in succession, this should be followed by no use of Success for a 30-day period or rotation to another insecticide class.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R065-029

Accepted: __/__/__

Initial printing.