

62719 -294

03/08/2000

1/2

T6P / SpinTor 2SC / Sec 3 Suppl Dried Beans & Peas / 01-04-00

SpinTor* 2SC

EPA Reg. No. 62719-294

Final printed labeling based on EPA accepted copy dated December 20, 1999.

Section 3 supplemental labeling for foliar insect control in dried beans and peas.

ACCEPTED

MAR 8 2000

62719-294

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

SpinTor* 2SC

EPA Reg. No. 62719-294

For Foliar Insect Control in Dried Beans and Peas

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for SpinTor* 2SC Naturalyte* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of SpinTor according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for SpinTor 2SC.

Directions for Use

Dried cultivars of *Lupinus* spp., *Phaseolus* spp. and *Pisum* spp., including but not limited to: lupins, kidney bean, lima bean, navy bean, pinto bean, tepary bean, adzuki bean, blackeyed pea, cowpea, crowder pea, chickpea, mungbean, field pea, pigeon pea and lentil.

Pests and Application Rates:

Pests	Application Rate [†]		
	Active Ingredient (lb/acre)	Product (fl oz/acre)	Acres per Gallon of Product
European corn borer (eggs and larvae)	0.047 - 0.094	3 - 6	43 - 21
Armyworms, corn earworm, loopers	0.062 - 0.094	4 - 6	32 - 21
Thrips, leafminers	0.094	6	21

[†]Use the higher end of the rate range for larger larvae or high infestations.

Specific Use Directions:

Application Timing: To determine when to apply, scout with enough regularity to monitor populations of labeled pests. Treat when a labeled pest infestation first appears, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but make no more than 6 applications per crop. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations for your area.

Restrictions:

- Do not apply more than 12 fl oz ounces (0.186 lb of spinosad) of SpinTor 2SC per acre per season.
- Do not feed treated forage or hay to meat or dairy animals.
- **Pre-harvest Interval:** Do not apply within 28 days of harvest.

*Trademark of Dow AgroSciences LLC

T6P / SpinTor 2SC / Sec 3 Suppl Dried Beans & Peas / 01-04-00
D06-064-009
EPA-accepted 12/20/1999
Initial printing.