

62719-294

11/08/2001

1/0

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## SpinTor\* 2SC

EPA Reg. No. 62719-294

### Foliar Insect Suppression in Cranberry

RECEIVED  
NOV 8 2001  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-294

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for SpinTor\* 2SC Naturalyte\* insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of SpinTor 2SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for SpinTor 2SC.

#### Directions for Use

Refer to product label for SpinTor\* 2SC Naturalyte\* insect control for General Use Precautions, Mixing and Application instructions.

#### Pests and Application Rates:

Pests	Broadcast Application Rate			Dilute Spray
	Active Ingrid. (lb/acre)	Product (fl oz/acre)	Acres/Gallon of Product	Product (fl oz/100 gal)
thrips	0.062 - 0.156	4 - 10	32 - 13	1 - 2.5
armyworms				
leafrollers				
loopers				
fireworms				
sparganothis fruitworm				
currant fruitfly				

#### Specific Use Directions:

**Application Timing:** For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Application rate within the rate range will depend on plant size and volume of foliage present and pest pressure. Use a higher rate in the rate range for larger larvae or moderate to severe infestations and and/or larger plant volume.

**Coverage of Dilute Sprays:** Dilute sprays should be uniformly applied to point of runoff. The rate per 100 gallons of spray is based on a spray volume of 400 gallons per acre. Gallonage of dilute sprays will vary depending on plant size and density.

#### Restrictions:

- Do not apply more than a total of 29 ounces of SpinTor 2SC (0.55 lb ai of spinosad) per acre per crop or make more than 6 applications per calendar year.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- **Pre-harvest Interval:** Do not apply within 3 days of harvest.

\*Trademark of Dow AgroSciences LLC

<p>T6P / SpinTor 2SC / Prop Sec 3 Suppl Cranberry / 12-07-00 D06-064-XXX EPA-accepted _/ _/ _ Initial printing.</p>	
---	--

