

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460	EPA Reg. Number: 62719-306	Date of Issuance: NOV 24 1998
		Term of Issuance: Conditional	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration		Name of Pesticide Product: Starane Plus Salvo	
(under FIFRA, as amended)			
Name and Address of Registrant (include ZIP Code): Dow AgroSciences LCC 9330 Zionsville Road Indianapolis, IN 46268			
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.			
On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.			
<p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following label changes before you release the product for shipment: <ol style="list-style-type: none"> a. Revise the EPA Registration Number to read, "EPA Reg. No. 62719-306". b. Delete the statement "Probable mucosal damage may contraindicate the use of gastric lavage" from the end of the Note to Physician in the Statement of Practical Treatment. c. Within the section for Personal Protective equipment, revise the phrase reading "... (such as probe and pump or spigot) ..." in the statement referring to containers over 1 gallon but less than five gallons by rewriting it to read "... (probe and pump) ...". 			
Signature of Approving Official:		Date: NOV 24 1998	

page 2
EPA Reg. No. 62719-306

- d. Within the section for Engineering Controls Statements, revise the instructions for containers over 5 gallons to read "A closed mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. The mechanical system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)]. The handler PPE requirements may be reduced or modified as specified in the WPS."
 - e. Within the Storage and Disposal block, revise the second sentence under Pesticide Disposal to read "Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater."
 - f. Within the General Use Precautions, revise the maximum amount of this product to be applied per acre per growing season under Maximum Application Rate to 2 2/3 pints.
 - g. Revise the grazing restriction for wheat, barley, and oats to 14 days after application.
3. Submit an acceptable Corrosion Characteristics study (\$830.6320) for this product within one year of the date of this notice.
4. Submit all of the required information to satisfy the data gaps in the data base for fluroxypyr. These data are described in the Notice of Pesticide Registration for Starane F Technical (EPA Registration No. 62719-285) dated September 30, 1998.
5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

DK:305-7546:HB/PM 23

CONCURRENCES

		CONCURRENCES						
SYMBOL	7505C							
SURNAME	D. KENNY							
DATE	11/19/99							

3616

(Base label):

(logo) Dow AgroSciences LLC

Starane* Plus Salvo®

For selective postemergence control of annual and perennial broadleaf weeds and volunteer potatoes in small grains and fallow cropland, and for on-farm non-cropland applications.

Active Ingredient(s):

fluroxypyr: 1-methylheptyl (4-amino-3,5-dichloro-6-fluoro-2-pyridyloxy)acetate ¹	12.1%
2,4-dichlorophenoxyacetic acid, 2-ethylhexyl ester ²	50.8%
Inert Ingredient(s).....	37.1%
Total	100.0%

ACCEPTED
with COMMENTS
In EPA Letter Dated

NOV 24 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 62719-306

¹Acid Equivalent: fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid - 8.4% - 0.75 lb/gal

²Acid Equivalent: 2,4-D: 2,4-dichlorophenoxyacetic acid - 33.6% - 3.00 lb/gal
Isomer specific by AOAC Method 978.05 15th Ed.

Contains xylene range aromatic solvent.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation • Harmful If Swallowed • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton
- Shoes plus socks
- Protective eyewear
- **Note: For containers of over 1 gallon, but less than 5 gallons:** Mixers and loaders who do not use a mechanical system (such as probe and pump or spigot) to transfer the contents of this container must wear coveralls or chemical-resistant apron in addition to other required PPE.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements

For containers of 5 gallons or more: Do not open pour product from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If In Eyes: Flush with plenty of water. Call a physician if irritation persists.

If Swallowed: Call a doctor or get medical attention. Do not induce vomiting. Do not give anything by mouth to an unconscious person. Avoid alcohol.

Note to Physician: May pose an aspiration pneumonia hazard. Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-XXX

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC
®Salvo is a registered trademark of Platte Chemical Co.

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents XXX

7816

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-XXX

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC

®Salvo is a registered trademark of Platte Chemical Co.

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents XXX

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes Eye Irritation • Harmful If Swallowed • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton
- Shoes plus socks
- Protective eyewear
- **Note: For containers of over 1 gallon, but less than 5 gallons:** Mixers and loaders who do not use a mechanical system (such as probe and pump or spigot) to transfer the contents of this container must wear coveralls or chemical-resistant apron in addition to other required PPE.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements

For containers of 5 gallons or more: Do not open pour product from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If In Eyes: Flush with plenty of water. Call a physician if irritation persists.

If Swallowed: Call a doctor or get medical attention. Do not induce vomiting. Do not give anything by mouth to an unconscious person. Avoid alcohol.

Note to Physician: May pose an aspiration pneumonia hazard. Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton
- Shoes plus socks
- Protective eyewear

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms or nurseries: When this product is applied to non-cropland areas, keep unprotected persons out of treated areas until sprays have dried.

Storage and Disposal

Do not contaminate water, food, feed or fertilizer by storage or disposal.

Storage: Store above 10°F or warm and agitate before use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal (Metal): Triple rinse (or equivalent). Puncture and dispose of in sanitary landfill, or by other procedures approved by state and local authorities.

Container Disposal (Plastic): Triple rinse (or equivalent). Puncture and dispose of in sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Starane* Plus Salvo® herbicide is a selective postemergence product for control of annual and perennial broadleaf weeds and volunteer potatoes in wheat (including durum), barley, or oats not under seeded with a legume and fallow cropland, and for on-farm non-cropland uses such as fence rows, building perimeters, around irrigation equipment and on-farm private roadways.

General Use Precautions

- Do not apply this product directly to, or otherwise permit it to come in direct contact with, susceptible crops or broadleaf plants including alfalfa, cotton, lettuce, edible beans, lentils, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, tobacco, grapes, legumes, fruit trees, canola, tame mustard, other vegetables or ornamentals. Vapors from this product may injure susceptible plants in the immediate vicinity.
- Avoid applications where proximity of susceptible crops or other susceptible broadleaf plants is likely to result in exposure to spray or spray drift.
- Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not apply in greenhouses.
- **Maximum Application Rate:** Do not apply more than 2 3/4 pints of Starane Plus Salvo (4.0 oz of fluroxypyr acid equivalent) per acre per growing season.
- A plant-back interval of 120 days must be observed for all rotational crops.
- **Chemigation:** Do not apply this product through any type of irrigation system.

Management of Kochia Biotypes: Research has suggested that many biotypes of kochia can occur within a single field. While kochia biotypes can vary in their susceptibility to Starane Plus Salvo, all will be suppressed or controlled by the 1 1/3 pint labeled rate. Application of Starane Plus Salvo at rates below the 1 1/3 pint rate can result in a shift to more tolerant biotypes within a field.

Precautions for Avoiding Spray Drift

Spray drift, even very small quantities of the spray which may not be visible, may severely injure susceptible crops whether dormant or actively growing. When applying Starane Plus Salvo, use low pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets. Under adverse weather conditions, fine spray droplets that do not settle rapidly onto target vegetation may be carried a considerable distance from the treatment area. A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Applications: To minimize spray drift, apply Starane Plus Salvo in a total spray volume of 8 or more gallons per acre using spray equipment designed to produce large-droplet, low pressure sprays.

11716

Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Spot treatments should be applied only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray. (See Application Directions.)

Aerial Application: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high potential for temperature inversion. Spray drift from aerial application can be minimized by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 3/4 the wing span of the aircraft. Spray pattern and droplet size distribution can be evaluated by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices, such as Automatic Flagman, may also be used. (See Application Directions.)

Do not apply under conditions of a low level air temperature inversion. A temperature inversion is characterized by little or no wind and lower air temperature near the ground than at higher levels. The behavior of smoke generated by an aircraft mounted device or continuous smoke column released at or near site of application will indicate the direction and velocity of air movement. A temperature inversion is indicated by layering of smoke at some level above the ground and little or no lateral movement.

Mixing Instructions

Starane Plus Salvo

Fill the spray tank approximately 1/2 to 3/4 full with water. Add the required amount of Starane Plus Salvo, then finish filling the spray tank. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixing

Starane Plus Salvo may be applied in tank mix combination with labeled rates of other herbicides provided (1) the tank mix product is labeled for the use site (timing and method of application is the same as Starane Plus Salvo); and (2) tank mixing with Starane Plus Salvo is not prohibited by the label of the tank mix product.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates. If products containing the same active ingredient are tank mixed, do not exceed the maximum allowable active ingredient use rates.
- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Starane Plus Salvo and other pesticides, fertilizers, or carriers. Use a clear glass jar with lid and mix the tank mix ingredients in their relative proportions. The tank mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily mix if agitated. An incompatible mixture is indicated by separation

Tank Mixing Instructions:

Fill the spray tank to approximately 1/2 to 3/4 of the total spray volume required. Start agitation. Add different formulation types in the order indicated, allowing time for complete mixing and dispersion after addition of each.

1. Add dry flowables; wettable powders; aqueous suspensions, flowables or liquids.
2. Maintain agitation and fill spray tank to 3/4 of total spray volume and then add Starane Plus Salvo and other emulsifiable concentrates and any solutions.

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Application Directions

Application Timing: Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds which are emerged at the time of application will be affected.** Foliage that is wet at the time of application may decrease control. Starane Plus Salvo herbicide applications are rain-fast within 1 hour after application.

Application Rates: Generally, application rates at the lower end of the recommended rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds) the higher rates within the rate range will be needed. Weeds growing in the absence of crop competition generally require higher rates to obtain satisfactory control or suppression.

Effect of Temperature on Herbicidal Activity: Herbicidal activity of Starane Plus Salvo is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum herbicidal activity is 55°F to 75°F. Reduced activity will occur when temperatures are below 45°F or above 85°F. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance.

Coverage: For best results, apply in 3 or more gallons per acre by air or in 8 or more gallons per acre by ground equipment. Do not exceed 40 gallons per acre total spray volume. Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Inadequate spray volume and coverage may result in decreased weed control. As crop canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Use larger nozzle tips or decrease spraying speed to increase spray volume rather than increasing boom pressure. Refer to manufacturer's recommendations for information on relationships between spray volume, and nozzle size and arrangement.

Spot Treatments: To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to directions provided below.

Hand-Held Sprayers: Hand-held or backpack sprayers may be used for spot applications of Starane Plus Salvo if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq ft. Mix the amount of Starane Plus Salvo (fl oz or ml) corresponding to the desired broadcast rate in one or more gallons of spray. To calculate the amount of product required for larger areas, multiply the table value (fl oz or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3,500 sq ft, multiply the table value by 3.5 (calc. 3,500 ÷ 1,000 = 3.5). An area of 1000 sq ft is approximately 10.5 X 10.5 yards (strides) in size.

Amount of Starane Plus Salvo to Equal Specified Broadcast Rate (Mix with 1 Gallon or More of Water and Apply to 1,000 sq ft)		
1 1/3 pt/acre	2 pt/acre	2 2/3 pt/acre
0.5 fl oz (15 ml)	0.75 fl oz (22 ml)	1.0 fl oz (29 ml)

[†]1 fl oz = 29.6 (30) ml

Weeds Controlled or Suppressed

(Number in parentheses (-) in weeds list refer to footnotes below.)

Weeds Controlled

- | | |
|---------------------|--------------------------------|
| bedstraw (cleavers) | milk vetch |
| bittercress | morningglory, annual |
| bull nettle | mousetail |
| burdock, common | mustards (except blue) (2) |
| burhead | nightshades |
| buttercup | pennycress |
| Carolina geranium | pepperweeds (except perennial) |
| carpetweed | pigweed |
| catnip | plantains |
| chickweed | poorjoe |
| chicory | prickly lettuce |
| cinquefoil | puncture vine |
| cocklebur | purslane, common |
| coffeeweed | quickweed |
| common purslane | ragweed (common, giant) |
| cornflower | rough fleabane |
| creeping Jenny | Russian thistle |
| dock, curly | shepherdspurse |
| evening primrose | sicklepod |
| fanweed | smallseeded falseflax |
| field pennycress | sneezeweed, bitter |
| figwort | sowthistle (annual, spiny) |
| flixweed | Spanishneedles |
| four o'clock | speedwell |
| goatsbeard | stinkweed |
| healall | sunflower |
| hedge bindweed | sweetclover |
| hemp dogbane | tansy mustard |
| horseweed | velvetleaf |
| ironweed | Venice mallow |
| Jacob's ladder | vetches |
| Jerusalem artichoke | Virginia copperleaf |
| jimsonweed | wild hemp |
| klamathweed | wild lettuce |
| kochia (1) | wild parsnip |
| ladysthumb | wild radish |
| lambquarters | yellow rocket |
| marestail | yellow starthistle |
| marshelder | |

1. Includes herbicide tolerant biotypes.
2. Apply prior to bolting.

14/16

Weeds Suppressed (1)

alfalfa	many-flowered aster
beggarticks	musk thistle
bull thistle	nettles
Canada thistle	nightshade
clover, red	peppergrass
dandelion	prickly lettuce
dogbanes	redstem filaree
fiddleneck	smartweed
field bindweed	tansyragwort
goldenrod	volunteer potato
ground ivy	wild buckwheat
hawkweed	wild carrot
henbit	wild garlic
hoarycress	wild onion
knotweed	

1. **Suppression** is expressed as a reduction in weed competition (reduction population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

Approved Uses

Crop Uses**Wheat (Including Durum), Barley, and Oats**

Apply as a broadcast postemergence treatment to actively growing wheat (including durum), barley or oats, from the 4 leaf crop growth stage up to flag leaf emergence (Zadoks scale 36) for control of broadleaf weeds. Apply when weeds are actively growing, but before weeds are 8 inches tall or vining. For control of volunteer potatoes, apply before potato plants are 8 inches tall. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Do not use if cereal crop is underseeded with a legume.**

Spot Application: Spot applications may be made, however, to prevent over-application spot treatments should be applied at rates and spray volumes equivalent to broadcast application. See instructions for "Spot Application" in "Application Directions" section.

Broadcast Application Rates:

(Number in parentheses (-) in table refer to footnotes below.)

Key Weeds Controlled (1)	Application Rate (pint/acre)
kochia bedstraw common ragweed field pennycress hemp dogbane mustard	1 1/3
Weeds Suppressed	
field bindweed nightshade species wild buckwheat volunteer potatoes	1 1/3 - 2 2/3 (2)

1. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.
2. Crop injury may occur at the 2 2/3 pt/acre rate.

Restrictions:

- Do not allow livestock to graze treated areas or harvest treated forage within 7 days of application.
- Do not make more than one application per season.
- **Preharvest Interval:** Do not apply closer than 14 days before cutting of hay or 40 days before harvesting of grain and straw.

Fallow Cropland

For best results, apply as a single broadcast treatment by ground or aerial equipment to control susceptible broadleaf weeds. Apply when weeds are actively growing, but before kochia is 8 inches tall and before wild buckwheat is vining. Starane Plus Salvo may be applied alone or in tank-mix combination with other herbicides (See tank mixing precautions in "Mixing Instructions" section.)

Broadcast Application Rates:

(Number in parentheses (-) in table refer to footnotes below.)

Key Weeds Controlled (1)	Application Rate (pint/acre)
kochia, bedstraw, common ragweed, field pennycress, hemp dogbane, mustard	1 1/3 - 2 2/3
Weeds Suppressed	
field bindweed, nightshade species, wild buckwheat volunteer potatoes	

1. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

On-Farm Non-Cropland

For best results, apply as a single broadcast treatment or spot treatment to control susceptible broadleaf weeds in on-farm non-cropland areas such as fence rows, building perimeters, around irrigation equipment and on-farm private roadways. Apply at the rate of 1 1/3 to 2 2/3 pints per acre when weeds are small and actively growing. Spot treatments should be applied at rates and spray volumes equivalent

16/16

to broadcast application. See instructions for "Spot Application" in "Application Directions" section. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

CRP Acres

Starane Plus Salvo may be applied to Conservation Reserve Program (CRP) acres. For best results, apply as a single broadcast treatment by ground or aerial equipment to control susceptible broadleaf weeds. Apply at the rate of 1 1/3 to 2 2/3 pints per acre when weeds are small and actively growing. Spot treatments should be applied at rates and spray volumes equivalent to broadcast application. See instructions for "Spot Application" in "Application Directions" section. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

Restriction: Grazing or haying of treated CRP acres is prohibited.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC
@Salvo is a registered trademark of Platte Chemical Co.
EPA-Accepted ___/___/___