

62719-322

10-18-2002

1/71

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 05-31-98



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other (X)

OPP Identifier Number Page 1 of 2 281704

Application for Pesticide - Section I

1. Company/Product Number: Dow AgroSciences/62719-322
2. EPA Product Manager: James A. Tompkins
3. Proposed Classification: None [] Restricted []
4. Company/Product (Name): Dow AgroSciences/Glyphomax* Plus
5. Name and Address of Applicant: Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268
6. Expedited Review: NOTIFICATION OCT 18 2002

Section - II

Amendment - Explain below []
Resubmission in response to Agency letter dated []
Notification - Explain below [X]
Final printed labels in response to Agency letter dated []
'Me Too' Application []
Other - Explain below []

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

Propose Changes by Notification:

- 1. Specific Crop Listings Canola, Crambe and Mustard on Page 21: The term "Mustard" has been clarified to read "Mustard Seed (Latest EPA accepted copy for Roundup Ultra is referenced)
2. Grain Sorghum (Hooded sprayers): Added precautionary text from EPA accepted copy for Roundup Ultra relating to use of hooded sprayers in tillering grain sorghum
3. Soybeans with Roundup Ready Gene: Restored specific paragraphs to RR soybean section that provide spray volume information for aerial and ground applications from EPA accepted supplemental labeling for Glyphomax Plus dated August 31, 1999

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging [] Yes* [] No
Unit Packaging [] Yes [] No
Water Soluble Packaging [] Yes [] No
2. Type of Container: Metal [] Plastic [] Glass [] Paper [] Other (Specify) []
3. Location of Net Contents Information: Label [] Container []
4. Size(s) Retail Container: []
5. Location of Label Directions: On Label [] On Labeling accompanying product []
6. Manner in Which Label is Affixed to Product: Lithograph [] Paper glued [] Stenciled [] Other []

Section - IV

1. Contact Point /Complete items directly below for identification of individual to be contacted, if necessary, to process this application)
Name: Steve A. McMaster, Title: Regulatory Manager, Telephone No. (317) 337-4670
Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.
2. Signature: Steve A. McMaster, 3. Title: Regulatory Manager, 4. Typed Name: Steve A. McMaster, 5. Date: September 18, 2002, 8. Date Application Received (Stamped): []

2/7/11



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
Page 2 of 2
281704

Application for Pesticide - Section I

| | | |
|--|---|---|
| 1. Company/Product Number Dow AgroSciences/62719-322 | 2. EPA Product Manager James A. Tompkins | 3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) Dow AgroSciences/Glyphomax* Plus | PM# PM / 23 | |
| 5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in Composition and labeling to: NOTIFICATION EPA Reg. No. OCT 18 2002 Product Name | |

Section - II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other- Explain below. |

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)
Additional Typographical Changes:

- Table on Page 23: Entry for Florida pusley at rates of 40 and 48 (stray hyphens before/after the weed height removed)
- Pokeweed in Table on Page 29: Stray hyphen in front of the range "3-40" removed
- Storage and Disposal box: "in a landfill approved form pesticide disposal ..." corrected to read "for"
- Controlling Droplet Size - Volume: "Nozzles with higher rated flows product larger droplets" (The word "product" corrected to read "produce".)
- Hooded sprayers (Second Paragraph): "extend front and read flaps of the hoods..." (The word "read" corrected read "rear".)
- Grass Seed Production (last sentence): The word "Grow" corrected to "Grower"

Section - IV

1. Contact Point /Complete items directly below for identification of individual to be contacted, if necessary, to process this application)

| | | |
|---------------------------|-----------------------------|---|
| Name Steve A. McMaster | Title Regulatory Manager | Telephone No. (Include Area Code) 317/337-4670 |
|---------------------------|-----------------------------|---|

| | | |
|---|--------------------------------|---|
| <p>Certification</p> <p>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowing false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</p> | | <p>8. Date Application Received (Stamped)</p> |
| 2. Signature | 3. Title Regulatory Manager | |
| 4. Typed Name Steve A. McMaster | 5. Date September 18, 2002 | |
| <p>Trademark of Dow AgroSciences LLC</p> | | |

3/71

NOTIFICATION
OCT 18 2002

Glyphomax* Plus

EPA Reg. No. 62719-322

Registration Notes:

Source label: text based on EPA accepted copy dated March 29, 2001, with conditions of acceptance.

Proposed Changes by Notification 09-16-02:

The following changes are being made at the request of California Department of Pesticide Regulation as a condition of acceptance of this label:

1. **Specific Crop Listings Canola, Crambe and Mustard on Page 21:** The term "Mustard" has been clarified to read "Mustard Seed" (Latest EPA-accepted copy for Roundup Ultra is referenced).
2. **Grain Sorghum (Hooded Sprayers):** Added precautionary text from EPA-accepted copy for Roundup Ultra relating to use of hooded sprayers in tillering grain sorghum.
3. **Soybeans with Roundup Ready Gene:** Restored specific paragraphs to RR soybean section that provide spray volume information for aerial and ground applications from EPA-accepted supplemental labeling for Glyphomax Plus dated August 31, 1999 (See also CDPR-accepted copy for Roundup Original dated July 3, 2002.) Because of the potential for crop injury, Dow AgroSciences has chosen **not** to include the section from the supplemental label for Roundup Original, allowing use of additional non-ionic surfactants, in the label for Glyphomax Plus.

[Editor's Note: Added text is underlined and deleted text is denoted by strike-through.]

The following typo's corrected on June 14, 2001:

1. Table on page 23: entry for Florida pusley at rates of 40 and 48 (stray hyphens before/after the weed height removed).
2. Pokeweed in table on page 29: Stray hyphen in front of the range "3-40" removed.
3. Storage and Disposal box: "in a landfill approved form pesticide disposal...." corrected to read "for."
4. Controlling Droplet Size--Volume: "Nozzles with higher rated flows product larger droplets." (The word "product" corrected to read "produce").
5. Hooded Sprayers (Second Paragraph): "extend front and read flaps of the hoods...(The word "read" corrected to read "rear").
6. Grass Seed Production (last sentence): The word "Grow" corrected to read "Grower"

*Trademark of Dow AgroSciences LLC
Roundup Ready® is a registered trademark of Monsanto Company

4/71

NOTIFICATION
OCT 18 2002

(Base Label):

(logo) Dow AgroSciences

Glyphomax* Plus

For control of annual and perennial weeds and woody plants in various cropping systems, fallow cropland and CRP acres, and farmsteads.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops (except crops with the Roundup Ready® herbicide tolerant gene), desirable plants and trees, because severe injury or destruction may result.

Active Ingredient(s):

| | |
|--|--------|
| glyphosate † N-(phosphonomethyl)glycine, isopropylamine salt..... | 41.0% |
| Inert Ingredients | 59.0% |
| Total Ingredients | 100.0% |

† Contains 4 pounds per gallon glyphosate, isopropylamine salt (3 pounds per gallon glyphosate acid).

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

S/71

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Domestic Animals: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read "Warranty Disclaimer," "Inherent Risks of Use," and "Limitation of Remedies" at end of label booklet. If terms are unacceptable, return at once unopened.**


In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-322

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC



6/71

Roundup Ready® is a registered trademark of Monsanto Company

Dow Agrosciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents __ gal

7/71

(Label Booklet):

(logo) Dow AgroSciences

Glyphomax* Plus

For control of annual and perennial weeds and woody plants in various cropping systems, fallow cropland and CRP acres, and farmsteads.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops (except crops with the Roundup Ready® herbicide tolerant gene), desirable plants and trees, because severe injury or destruction may result.

Active Ingredient(s):

| | |
|---|--------|
| glyphosate †: N-(phosphonomethyl)glycine, isopropylamine salt..... | 41.0% |
| Inert Ingredients..... | 59.0% |
| Total Ingredients..... | 100.0% |

† Contains 4 pounds per gallon glyphosate, isopropylamine salt (3 pounds per gallon glyphosate acid).

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read "Warranty Disclaimer," Inherent Risks of Use," and "Limitation of Remedies" at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.


EPA Reg. No. 62719-322

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC

Roundup Ready® is a registered trademark of Monsanto Company

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.



8/11

Herbicide

Net Contents __ gal

9/7/1

| Contents | Page |
|---|-------------|
| Precautionary Statements | - |
| Hazards To Humans and Domestic Animals | - |
| Environmental Hazards | - |
| Physical Or Chemical Hazards | - |
| Storage and Disposal | - |
| General Information | - |
| (How this product Works) | - |
| Mixing | - |
| Mixing With Water | - |
| Tank Mixing Procedure | - |
| Mixing For Hand-Held Sprayers | - |
| Ammonium Sulfate | - |
| Colorants or Dyes | - |
| Drift Control Additives | - |
| Application Equipment and Techniques | - |
| Aerial Equipment | - |
| Ground Broadcast Equipment | - |
| Hand-Held and High-Volume Equipment | - |
| Selective Equipment (Wipers, etc.) | - |
| Injection Systems | - |
| CDA Equipment | - |
| Cut Stump Application | - |
| Crops (Alphabetical) | - |
| Alfaifa, Clover and other Forage Legumes | - |
| Asparagus | - |
| Canola, Crambe, Mustard (<u>Seed</u>) | - |
| Cereal Crops | - |
| Christmas Trees | - |
| Citrus Crops | - |
| Conservation Reserve Program (CRP) | - |
| Corn | - |
| Cotton | - |
| Dry Peas, Lentils, Chick Peas | - |
| Fallow Systems (Including Postharvest Applications) | - |
| Grain Sorghum (Milo) | - |
| Grass Seed Production | - |
| Herbs | - |
| Pastures | - |
| Peanuts | - |
| Small Fruits and Berries | - |
| Soybeans | - |
| Sugarcane | - |
| Sunflowers | - |
| Tree and Vine Crops (General) | - |
| Tree Fruits | - |
| Tree Nuts | - |
| Tropical Crops | - |
| Vegetable Crops | - |
| Vine Crops | - |
| Roundup Ready® Crops | - |
| Roundup Ready Corn | - |

10/7/11

- Roundup Ready Soybeans -
- Farmsteads -
- General Nonselective Weed Control, Trim-and-Edge and Bare Ground -
- Chemical Mowing -
- Habitat Management -
- Habitat Restoration and Maintenance -
- Wildlife Food Plots -
- Annual Weeds Rate Tables (Alphabetically by Species) -
- Rates For 10 To 40 GPA -
- Tank Mixtures of Glyphomax* Plus with 2,4-D, Dicamba or Tordon* 22K -
- Perennial Weeds Rate Table (Alphabetically by Species) -
- Woody Brush and Trees Rate Table (Alphabetically by Species) -
- Warranty Disclaimer -
- Inherent Risks of Use -
- Limitation of Remedies -

11/7/1

combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

This is an end-use product. Dow AgroSciences does not intend and has not registered it for reformulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves, such as butyl rubber \geq 14 mils, or natural rubber \geq 14 mils, or neoprene rubber \geq 14 mils, or nitrile rubber \geq 14 mils
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container contains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

Refillable Portable Containers: Do not reuse this container except to refill in accordance with a valid Dow AgroSciences Repackaging agreement. If not refilled or returned to the authorized repackaging facility, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Bulk Containers: Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

12/17/11

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Causes Eye Irritation

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Domestic Animals: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly

13/7/11

Plastic 1-Way Container Disposal: Do not reuse this container. Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Drums: Do not reuse container. Return container per any Dow AgroSciences container return program. If not returned, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information **(How this product works)**

Glyphomax* Plus herbicide is a postemergence, systemic herbicide with no soil residual activity and is intended for control of annual and perennial weeds and woody plants in various cropping systems, fallow cropland and CRP acres, and farmsteads. Glyphomax Plus is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Do not add surfactants, additives containing surfactants, buffering agents or pH-adjusting agents to the spray solution when Glyphomax Plus is the only pesticide used. Ammonium sulfate, drift control additives, or dyes and colorants may be used. See the "Mixing" section of this label for instructions.

Time to Symptoms: The active ingredient in Glyphomax Plus moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of Glyphomax Plus and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of above ground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the annual, perennial, woody brush and trees rate tables for recommendations for specific weeds.

Always use the higher rate of Glyphomax Plus per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced herbicidal activity may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash Glyphomax Plus off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action: The active ingredient in Glyphomax Plus inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

14/71

No Soil Activity: Weeds must be emerged at the time of application to be controlled by Glyphomax Plus. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow.

When Glyphomax Plus comes in contact with soil, it is bound to soil particles. Under recommended use situations, once Glyphomax Plus is bound to soil particles, it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treated area or if the soil is transported off-site. The strong affinity of Glyphomax Plus to soil particles prevents Glyphomax Plus from leaching out of the soil profile and entering ground water.

Biological Degradation: Degradation of Glyphomax Plus is primarily a biological process carried out by soil microbes.

Volatility: Glyphomax Plus is non-volatile. Therefore, it cannot move as a vapor after application to affect nearby vegetation.

Toxicology Testing: Exposure to workers and other applicators generally is expected to pose minimal risks based on results of short-term toxicity studies. Glyphosate has been thoroughly tested and determined not to cause cancer or other adverse long-term health effects.

Tank Mixing: Glyphomax Plus does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of Glyphomax Plus with herbicides or other materials that are not expressly recommended in this labeling. Mixing Glyphomax Plus with herbicides or other materials not recommended on this label may result in reduced performance.

Annual Maximum Use Rate: Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of Glyphomax Plus per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated use rate.

For noncrop uses, the combined total of all treatments must not exceed 10.6 quarts of Glyphomax Plus per acre per year.

Attention

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

AVOID DRIFT. Extreme care must be used when applying Glyphomax Plus to prevent injury to desirable plants and crops.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of Glyphomax Plus can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of Glyphomax Plus increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or

157/11

when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **Avoid applying at excessive speed or pressure.**

NOTE: Use of Glyphomax Plus in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information:**

Importance of Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size:

Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

16/71

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, presence of an inversion can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Mixing

Clean sprayer parts immediately after using Glyphomax Plus by thoroughly flushing with water.

NOTE: reduced results may occur if water containing soil is used, such as visibly muddy water or water from ponds and ditches that is not clear.

Mixing with Water

Glyphomax Plus mixes readily with water. Mix spray solutions of Glyphomax Plus as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of Glyphomax Plus near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

1/1/71

Tank Mixing Procedure

Mix labeled tank mixtures of Glyphomax Plus with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it **slowly** through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture **slowly** through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of Glyphomax Plus near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive and water-soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of Glyphomax Plus with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" section under "General Information" for additional precautions.

Mixing for Hand-held Sprayers

Prepare the desired volume of spray solution by mixing the amount of Glyphomax Plus in water as shown in the following table:

Spray Solution

| Spray Concentration (percent) | Amount of Glyphomax Plus for Desired Volume: | | |
|-------------------------------|--|--------|---------|
| | 1 gal | 25 gal | 100 gal |
| ½% | 2/3 fl oz | 1 pt | 2 qt |
| 1% | 1 1/3 fl oz | 1 qt | 1 gal |
| 1 ½% | 2 fl oz | 1 ½ qt | 1 ½ gal |
| 2% | 2 2/3 fl oz | 2 qt | 2 gal |
| 5% | 6 1/2 fl oz | 5 qt | 5 gal |
| 10% | 13 fl oz | 10 qt | 10 gal |

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of Glyphomax Plus be mixed with water in a larger container. Fill sprayer with the mixed solution.

18/11

Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Glyphomax Plus, particularly when tank mixed with certain residual herbicides on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

Note: When using ammonium sulfate, apply Glyphomax Plus at rates recommended in this label. Lower rates will result in reduced performance.

Colorants or Dyes

Agriculturally-approved colorants or marking dyes may be added to Glyphomax Plus. Colorants or dyes used in spray solutions of Glyphomax Plus may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

Drift Control Additives

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and CDA equipment. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Application Equipment and Techniques

Do not apply Glyphomax Plus through any type of irrigation system.

Glyphomax Plus may be applied with the following application equipment:

Aerial: Fixed Wing and Helicopter

Ground Broadcast Spray: Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held and High-Volume Spray Equipment: Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, mistblowers¹, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

¹Glyphomax Plus is not registered in California or Arizona for use in mistblowers.

Selective Equipment: Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems: Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA): Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.

19/7/1

Cut Stump Application: Apply using suitable equipment to ensure coverage of the entire cambium of cut stems.

Aerial Equipment

Do not apply Glyphomax Plus using aerial spray equipment except under conditions as specified within this label.

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 1 quart per acre. Aerial applications of Glyphomax Plus may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems and preharvest applications. Refer to the individual use area sections of this label for recommended volumes and application rates.

For aerial application in California or Arkansas, refer to the federal supplemental label for aerial applications in that state for specific instructions, restrictions and requirements. Tank mixtures of Glyphomax Plus plus Banvel (dicamba) or 2,4-D herbicide may not be applied by air in California.

Avoid direct application to any body of water.

AVOID DRIFT: do not apply during low-level inversion conditions, when winds are gusty or under any other condition that favors drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Ensure uniform application: To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of Glyphomax Plus accumulated during spraying or from spills. **Prolonged exposure of Glyphomax Plus to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

Ground Broadcast Equipment

Use the recommended rates of Glyphomax Plus in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Hand-Held and High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

20/7/1

For control of weeds listed in the annual weeds rate tables, apply a 0.5 percent solution of Glyphomax Plus to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

Selective Equipment

Glyphomax Plus may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

Avoid contact of herbicide with desirable vegetation.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and hooded applicators

Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **Extreme care must be exercised to avoid contact of herbicide with desirable vegetation.**

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Speed of operation must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

21/7/1

Wiper applicators and sponge bars

Wiper applicators are devices that physically wipe appropriate amounts of Glyphomax Plus directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using Glyphomax Plus by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators: Mix 1 gallon of Glyphomax Plus in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this section.

For Porous-Plastic Applicators: Solutions ranging from 33 to 100 percent of Glyphomax Plus in water may be used in porous-plastic wiper applicators.

When applied as recommended, Glyphomax Plus **controls** the following weeds:

| | |
|-----------------|------------------|
| corn, volunteer | sicklepod |
| panicum, Texas | spanishneedles |
| rye, common | starbur, bristly |
| shattercane | |

When applied as recommended, Glyphomax Plus **suppresses** the following weeds:

| | |
|------------------------|------------------|
| beggarweed, Florida | pigweed, redroot |
| bermudagrass | ragweed, common |
| dogbane, hemp | ragweed, giant |
| dogfennel | smutgrass |
| guineagrass | sunflower |
| johnsongrass | thistle, Canada |
| milkweed | thistle, musk |
| nightshade, | vaseygrass |
| silverleaf | velvetleaf |

22/1/1

Injection Systems

Glyphomax Plus may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix Glyphomax Plus with the concentrate of other products when using injection systems.

CDA Equipment

The rate of Glyphomax Plus applied per acre by vehicle-mounted controlled droplet application (CDA) equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units, apply a 20 percent solution of Glyphomax Plus at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of Glyphomax Plus at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

Cut Stump Application

Types of Application: Treating cut stumps in any noncrop site listed on this label

Specific Use Recommendations: Glyphomax Plus will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply Glyphomax Plus using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of Glyphomax Plus to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion

| | |
|-------------|-----------|
| alder | saltcedar |
| eucalyptus | sweetgum |
| madrone | tan oak |
| oak | willow |
| reed, giant | |

Precautions and Restrictions: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Injury resulting from root grafting may occur in adjacent woody brush or trees.

CROPS (Alphabetical)

This section is organized alphabetically by crop category. There may be several labeled crops listed in a crop category.

Unless otherwise specified, applications may be made to control any weeds listed in the annual, perennial and woody brush tables. Also refer to the "Selective Equipment" section.

23/1/11

For any crop not listed in this "Crops" section, applications must be made at least 30 days prior to planting.

For broadcast postemergence treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

When applying Glyphomax Plus prior to transplanting crops into plastic mulch, residues may be removed from the plastic by 0.5 inches of water via sprinkler irrigation or natural rainfall.

Alfalfa, Clover, and Other Forage Legumes

Labeled Crops: Alfalfa, clover, kudzu, lespedeza, lupin, sainfoin, trefoil, velvet bean, vetch, crown vetch, milk vetch

Types of Applications: Preplant, preemergence, at-planting, spot treatment (alfalfa and clover only), wiper applicators (alfalfa and clover only), renovation, preharvest

Preplant, Preemergence and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting crops listed in this section. Applications must be made prior to emergence of the crop.

Precautions and Restrictions: Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Preharvest (Alfalfa only)

Specific Use Recommendations: Glyphomax Plus may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This application will severely injure or destroy the stand of alfalfa. Glyphomax Plus will control annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa. The treated crop and weeds can be harvested and fed to livestock after 36 hours. Allow a minimum of 36 hours between application and harvest. Applications may be made at any time of the year. Make only one application to an existing stand of alfalfa per year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

Precautions and Restrictions: Do not apply more than 1 quart of Glyphomax Plus per acre as a preharvest treatment. Do not use for alfalfa grown for seed, as a reduction in germination or vigor may occur.

Spot treatment or Wiper applications (Alfalfa and Clover only)

Specific Use Recommendations: Glyphomax Plus may be applied as a spot treatment in alfalfa or clover. Glyphomax Plus may be applied with wiper applicators to control or suppress the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label. Applications may be made in the same area at 30-day intervals.

Precautions and Restrictions: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

24/71

Renovation

Specific Use Recommendations: Glyphomax Plus may be applied as a broadcast spray to existing stands of alfalfa, clover, and other labeled forage legumes. Labeled crops may be planted into the treated area.

Precautions and Restrictions: Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Asparagus

Types of Applications: Preplant, preemergence, spot treatment, postharvest

Preplant, Preemergence

Specific Use Recommendations: Glyphomax Plus may be applied prior to emergence of asparagus.

Precautions and Restrictions: Do not apply within a week before the first spears emerge.

Spot treatment

Specific Use Recommendations: Glyphomax Plus may be applied immediately after cutting, but prior to the emergence of new spears.

Precautions and Restrictions: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Postharvest

Specific Use Recommendations: Glyphomax Plus may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears.

Precautions and Restrictions: Direct contact of the spray with the asparagus may result in serious crop injury. Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

Canola, Crambe, Mustard (Seed)

Types of Applications: Preplant, preemergence, at-planting, post-harvest

Preplant, Preemergence and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting of canola, crambe, or mustard. Applications must be made prior to emergence of the crop.

Precautions and Restrictions: Do not apply more than 1.6 quarts of this product per acre by ground.

25/7/11

Postharvest

Specific Use Recommendations: Glyphomax Plus may be applied after harvest of canola, crambe, or mustard. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures of Glyphomax Plus with 2,4-D or dicamba may be used.

Precautions and Restrictions: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Do not harvest or feed treated vegetation for 8 weeks following application.

Cereal Crops

Labeled Crops: Barley, buckwheat, millet (pearl, proso), oats, rice, rye, teosinte, triticale, wheat (all), wild rice

Types of Applications: Preplant, preemergence, at-planting, spot treatment (except rice), post-harvest, preharvest (wheat only), wiper applicators (wheat only)

Do not treat rice fields or levees when the field contains floodwater.

Preplant, Preemergence and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting of cereal crops. Applications must be made prior to emergence of the crop.

Spot treatment (except rice)

Specific Use Recommendations: Glyphomax Plus may be applied as a spot treatment in cereal crops. Apply Glyphomax Plus before heading in small grains.

Precautions and Restrictions: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

Postharvest

Specific Use Recommendations: Glyphomax Plus may be applied after harvest of cereal crops. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures of Glyphomax Plus with 2,4-D or dicamba may be used.

Precautions and Restrictions: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Do not harvest or feed treated vegetation for 8 weeks following application.

Preharvest (wheat only)

Specific Use Recommendations: Glyphomax Plus provides weed control when applied prior to harvest of wheat. Apply after the hard-dough stage of grain (30% or less grain moisture) and at least 7 days prior to harvest. Wheat stubble may be grazed immediately after harvest.

Glyphomax Plus may be applied using either aerial or ground spray equipment. For ground or aerial applications, apply Glyphomax Plus in 3 or more gallons of water per acre.

Precautions and Restrictions: Do not apply more than 1 quart of Glyphomax Plus per acre. Do not apply to wheat grown for seed, as a reduction in germination or vigor may occur.

26/7/1

Wiper applications (wheat only)

Specific Use Recommendations: Wiper applications may be used in wheat. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth, when the rye is at least 6 inches above the wheat crop.

Precautions and Restrictions: Allow at least 35 days between application and harvest. Do not use roller applicators.

Red Rice Control Prior to Planting Rice: Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled. **Avoid spraying during low humidity conditions as reduced control may result. Do not treat fields or levees when the fields contain water. Do not re-flood treated fields for 8 days following application.**

Christmas Trees

Types of Applications: Post-directed, spot treatment, site preparation

Post-directed, Spot treatment

Specific Use Recommendations: Glyphomax Plus may be used as a post-directed spray and spot treatment around established Christmas trees.

Precautions and Restrictions: Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. **Glyphomax Plus is not recommended for use as an over-the-top broadcast spray in Christmas trees.** Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established Christmas trees.

Site preparation

Specific Use Recommendations: Glyphomax Plus may be used prior to planting Christmas trees.

Precautions and Restrictions: Precautions should be taken to protect nontarget plants during site preparation applications.

Citrus Crops

Labeled Crops: Calamondin, chironja, citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), orange (all), pummelo, tangelo, tangor

Types of Applications: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment

NOTE: for general use directions, see the "tree, nut and vine (general)" section. The following directions are specific to citrus crops.

Florida and Texas only: For burndown or control of the weeds listed below, apply the recommended rates of Glyphomax Plus in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

27/71

For goatweed, apply 2 to 3 quarts of Glyphomax Plus per acre. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Karmax II herbicide or Karmex herbicide may improve control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Perennial Weeds:

| Weed Species | Glyphomax Plus Rate Per Acre | | | |
|--|------------------------------|------|------|------|
| | 1 qt | 2 qt | 3 qt | 5 qt |
| bermudagrass | B | -- | PC | C |
| guineagrass (area) (Texas and Florida ridge) (Florida flatwoods) | B | C | C | C |
| | -- | B | C | C |
| paragrass | B | C | C | C |
| torpedograss | S | -- | PC | C |

S = Suppression B = Burndown
PC = Partial control C = Control

Precautions and Restrictions: Allow a minimum of 1 day between last application and harvest.

Conservation Reserve Program (CRP)

Types of Applications: Renovation (rotating out of CRP), site preparation, postemergence, wiper

Rotating out of CRP, Site preparation

Specific Use Recommendations: Glyphomax Plus may be used to prepare CRP land for crop production.

Postemergence, Wiper

Specific Use Recommendations: Glyphomax Plus may be used to suppress competitive growth and seed production of undesirable vegetation in CRP acres. Such applications may be made with wiper application equipment or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 8 to 10 fluid ounces of Glyphomax Plus per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Precautions and Restrictions: Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant.

Corn

Types of Corn: Field corn, seed corn, sweet corn and popcorn

Types of Applications: Preplant, preemergence, at-planting, hooded sprayers, spot treatment, preharvest, post-harvest

Preplant, Preemergence and At-Planting

28/7/1

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting corn. Applications must be made prior to emergence of the crop.

Tank Mixes: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. See the map in the Annual Weeds section of this label for areas included in this recommendation.

Tank mixtures with the following herbicide products may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue:

| | | |
|------------------|-------------------|-----------------|
| atrazine | FulTime | LeadOff |
| Bicep II | Guardsman | Micro-Tech |
| Bicep II Magnum | Harness | Partner |
| Bicep Lite II | Harness Xtra | Pendimax* |
| Magnum | Harness Xtra 5.6L | (pendimethalin) |
| Bladex/Cyanazine | Hornet* | Python* |
| Bullet | Hornet WDG | Simazine |
| dicamba | Lariat | Surpass EC |
| Degree | Lasso/Alachlor | TopNotch |
| Degree Xtra | Linex | |
| Dual II | Lorox | |
| Dual II Magnum | Marksman | |
| Extrazine | | |
| Frontier | | |

For improved burndown, Glyphomax Plus may be tank mixed with 2,4-D or dicamba.

Annual weeds: For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Glyphomax Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of Glyphomax Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall.

Precautions and Restrictions: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn.

The tank mix recommendations in this section are not registered in California.

Hooded Sprayers

Specific Use Recommendations: This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray pattern. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- Spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application
- Corn must be at least 12 inches tall, measured without extending the leaves.

29/7/1

- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph
- Maximum wind speed: 10 mph
- Use low drift nozzles

Crop injury may occur when the foliage of treated weeds comes in direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Precautions and Restrictions: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers. Do not apply more than 3 quarts per acre per year of this product using hooded sprayer application.

Spot treatment

Specific Use Recommendations: For spot treatments, apply Glyphomax Plus prior to silking of corn.

Precautions and Restrictions: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

Preharvest

Specific Use Recommendations: Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts per acre of this product. For aerial applications, apply up to 1 qt per acre of this product.

Precautions and Restrictions: Allow a minimum of 7 days between application and harvest. It is not recommended that corn grown for seed be treated because a reduction in germination or vigor may result.

Post-harvest

Specific Use Recommendations: Glyphomax Plus may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures of Glyphomax Plus with 2,4-D or dicamba may be used.

Precautions and Restrictions: Do not harvest or feed treated vegetation for 8 weeks following application.

Cotton

Types of Applications: Preplant, preemergence, at-planting, hooded sprayer, selective equipment, spot treatment, preharvest

Preplant, Preemergence, and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting cotton. Applications must be made prior to emergence of the crop.

30/7/11

Hooded sprayer, Selective equipment

Specific Use Recommendations: Glyphomax Plus may be applied through hooded sprayers, recirculating sprayers, shielded applicators or wiper applicators in cotton. Allow at least 7 days between application and harvest.

Precautions and Restrictions: See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.

Spot treatment

Specific Use Recommendations: For spot treatments, apply Glyphomax Plus prior to boll opening of cotton.

Precautions and Restrictions: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

Preharvest

Specific Use Recommendations: Glyphomax Plus provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the annual, perennial and woody brush tables. Apply 1 pint to 2 quarts of Glyphomax Plus per acre for cotton regrowth inhibition. Allow a minimum of 7 days between application and harvest of cotton.

Glyphomax Plus may be applied using either aerial or ground spray equipment. For ground applications, apply Glyphomax Plus in 10 to 20 gallons of water per acre. For aerial applications, apply Glyphomax Plus in 3 to 10 gallons of water per acre.

Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

Glyphomax Plus may be tank mixed with DEF 6, Folex, or Prep defoliant to provide additional enhancement of cotton leaf drop.

Precautions and Restrictions: Do not feed or graze treated cotton forage or hay following preharvest applications. **Do not apply more than 1 quart of Glyphomax Plus per acre by air.** Do not apply more than 2 quarts of Glyphomax Plus per acre by ground. Do not apply to cotton grown for seed, as a reduction in germination or vigor may occur.

Dry Peas, Lentils, Chick Peas

(See "Vegetable Crops" section for specific use directions)

Fallow Systems (Including Post Harvest Applications)

Types of Applications: Chemical fallow, preplant fallow beds, aid-to-tillage

Post Harvest Use

Specific Use Recommendations: Glyphomax Plus may be applied to control existing weeds or volunteer crop following harvest of labeled crops. Weeds should be allowed to regrow after damage incurred during harvest and recover from environmental stress before application. Apply prior to heading of grass weeds

3/1/71

and, if possible, before broadleaf weeds exceed a height of 24 inches. Applications may be made during the fallow period up until the planting or emergence of labeled crops, but for any crop not listed on this label, applications must be made at least 30 days prior to planting. Ground or aerial equipment may be used.

Refer to annual or perennial weeds rate tables for application rates and species controlled. If Glyphomax Plus, applied post harvest, may be tank mixed with other herbicides. See "Chemical Fallow" section below for specific recommendations for tank mixing.

Chemical fallow

Specific Use Recommendations: Glyphomax Plus may be applied during the fallow period prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 30 days prior to planting. Glyphomax Plus may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures of Glyphomax Plus with 2,4-D, dicamba, Tordon[®] 22K herbicide, atrazine or cyanazine herbicide may be used.

Precautions and Restrictions: Tank mixtures of Glyphomax Plus with Banvel (dicamba), Tordon 22K or 2,4-D may not be applied by air in California.

Follow planting, cropping, crop rotation and other restrictions and use precautions on the labels of each product used in tank mixtures.

Dicamba: Some crop injury may occur if dicamba is applied within 45 days of planting.

Tordon 22K[†]: The addition of Tordon 22K in a mixture with Glyphomax Plus may provide short-term residual control of selected weed species. Application of Glyphomax Plus in tank mix with Tordon 22K should be made only to land that will be planted the following year to grass, barley, oats, wheat, grain sorghum (milo) or fallowed. Some crop injury may occur if Tordon 22K is applied within 45 days of planting. Do not plant grain sorghum within 8 months after application. Tordon 22K is not intended for use on land planted to sweet sorghum.

[†] Tordon 22K is not registered for use in California.

Preplant fallow beds

Specific Use Recommendations: Glyphomax Plus may be applied to fallow beds prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 30 days prior to planting. Glyphomax Plus will control weeds listed in the annual, perennial and woody brush tables.

In addition, 12 fluid ounces of Glyphomax Plus plus 2 to 3 oz of Goal 2XL per acre will control the following weeds with the maximum height or length indicated: 3" -- common cheeseweed, chickweed, groundsel; 6" -- London rocket, shepherd's-purse.

16 fluid ounces of Glyphomax Plus plus 2 to 3 oz of Goal 2XL per acre will control the following weeds with the maximum height or length indicated: 6" -- common cheeseweed, groundsel, marestail (*Coryza canadensis*), 12" -- chickweed, London rocket, shepherd's-purse.

Aid-to-tillage

Specific Use Recommendations: Glyphomax Plus may be used in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of Glyphomax Plus in 3 to 10 gallons of water per acre. Make

32/1/11

applications before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

Precautions and Restrictions: Tank mixtures Glyphomax Plus with residual herbicides may result in reduced performance.

Flax

Types of Applications: Preplant, preemergence, at-planting, post-harvest

Preplant, Preemergence and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting of flax. Applications must be made prior to emergence of the crop.

Postharvest

Specific Use Recommendations: Glyphomax Plus may be applied after harvest of flax. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures of Glyphomax Plus with 2,4-D or dicamba may be used.

Precautions and Restrictions: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Do not harvest or feed treated vegetation for 8 weeks following application.

Grain Sorghum (Milo)

Types of Applications: Preplant, preemergence, at-planting, spot treatment, wiper applicators, hooded sprayers, preharvest, post-harvest

Preplant, Preemergence, At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting grain sorghum. Applications must be made prior to emergence of the crop.

The following herbicide products may be applied in tank mix combination with Glyphomax Plus in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. Apply before, during or after planting in conventional tillage systems, into a cover crop, established sod or over previous crop residue.

| | |
|----------|------------------|
| atrazine | Lariat |
| Bicep II | Lasso / alachlor |
| Bullet | Micro-Tech |
| Dual II | Partner |

Annual weeds: For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Glyphomax Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of Glyphomax Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall.

33/71

Spot treatment and Wiper applications

Specific Use Recommendations: Glyphomax Plus may be applied as a spot treatment in grain sorghum. Make spot treatments before heading of milo. Glyphomax Plus may be applied with wiper applicators to control or suppress the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label.

Precautions and Restrictions: For spot treatment, do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

For wiper applicators, allow at least 40 days between application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

Hooded Sprayers

Specific Use Recommendations: This product may be used through hooded sprayers for weed control between the rows of grain sorghum. Only hooded sprayers that completely enclose the spray pattern may be used.

When applying to grain sorghum that is grown on raised beds, ensure that the hood is designed to completely enclose the spray pattern. If necessary, extend the front and ~~rear~~rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- Spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application
- Grain sorghum must be at least 12 inches tall, measured without extending the leaves. Treat before milo extends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph
- Maximum wind speed: 10 mph
- Use low drift nozzles

Crop injury may occur when the foliage of treated weeds comes in direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Precautions and Restrictions: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. Do not graze or feed grain sorghum forage or fodder following applications of this product through hooded sprayers. Do not apply more than 3 quarts per acre per year of this product using hooded sprayer application.

Preharvest

Specific Use Recommendations: Glyphomax Plus may be applied prior to harvest of grain sorghum. Make applications at 30% grain moisture or less.

Precautions and Restrictions: Do not apply more than 2 quarts of this product per acre. Allow a minimum of 7 days between application and harvest of sorghum. It is not recommended that sorghum

34/71

grown for seed be treated, as reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California.

Post-harvest

Specific Use Recommendations: Glyphomax Plus may be applied after harvest of grain sorghum. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures of Glyphomax Plus with 2,4-D or dicamba may be used.

Glyphomax Plus may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of Glyphomax Plus per acre for control, or 1.5 pints of Glyphomax Plus per acre for suppression.

Precautions and Restrictions: Do not harvest or feed treated vegetation for 8 weeks following application.

Grass Seed Production

Types of Applications: Preplant, preemergence, renovation, site preparation, shielded sprayers, wiper applicators, spot treatments, creating rows in annual ryegrass.

Specific Use Recommendations: Applications may be made before, during or after planting or renovation of turf or forage grass areas grown for seed production. Applications must be made prior to the emergence of the crop to avoid crop injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control.

Precautions and Restrictions: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

Do not feed or graze treated areas for 8 weeks following application.

Shielded Sprayers

Specific Use Recommendations: Apply 1-3 quarts of this product as a broadcast spray in 10 to 20 gallons of water per acre to control weeds in rows. Uniform planting in straight rows aids in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.

Precautions and Restrictions: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

Wiper Applications

Precautions and Restrictions: Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators should be adjusted so that the wiper contact point is at least two (2) inches above the desirable vegetation. Weeds should be a minimum of six (6) inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when height of weeds varies so that not all weeds are contacted. In these

357/11

instances, repeat treatments may be necessary. Better results may be obtained if 2 applications are made in opposite directions.

Spot Treatments

Specific Use Recommendations: Use a 1 - 1.5% solution.

Precautions and Restrictions: Apply this product prior to heading of grasses. Do not treat more than 10 percent of the total field to be harvested. The crop receiving the spray in the treated area will be killed and, for the same reason, take care to avoid drift or spray outside target areas.

Creating Rows in Annual Ryegrass

Specific Use Recommendations: Use Us 16 - 32 fl oz of this product per acre mixed with water. Use the higher rate when the ryegrass is greater than 6 inches tall. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height.

Precautions and Restrictions: Set nozzle heights to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use of low pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

Grower assumes all responsibility for crop losses from misapplication.

Herbs

Types of Herbs: Peppermint, spearmint

Specific Use Recommendations: Glyphomax Plus may be used as a spot treatment in spearmint and peppermint. Apply spray-to-wet with hand-held equipment, such as backpack and knapsack sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution on to a limited area.

Precautions and Restrictions: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. No more than one-tenth of any acre should be treated at one time. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for this reason.

Pastures

Type of Pastures: Bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa and clover

Types of Applications: Spot treatment, wiper application, preplant, preemergence, pasture renovation

Spot treatment and Wiper application

Specific Use Recommendations: Glyphomax Plus may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals.

Precautions and Restrictions: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated

36/71

at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

Preplant, Preemergence and Pasture renovation

Specific Use Recommendations: Glyphomax Plus may be applied prior to planting or emergence of forage grasses and legumes. In addition, Glyphomax Plus may be used to control perennial pasture species listed on this label prior to re-planting.

Precautions and Restrictions: Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Peanuts

Types of Applications: Preplant, preemergence, at-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting peanuts. Applications must be made prior to the emergence of the crop.

Small Fruits and Berries

Labeled Crops: Blackberry, blueberry, boysenberry, cranberry, currant, dewberry, elderberry, gooseberry, huckleberry, loganberry, olallieberry, raspberry (black, red), youngberry

Types of Applications: Preplant, preemergence, directed spray (except cranberry), wiper application

Specific Use Recommendations: Glyphomax Plus may be applied as a preplant or preemergence broadcast application or as a wiper application for crops listed in this section. Directed sprays may be applied to any crop except cranberries. For wick or wiper applicators, mix 1 gallon of Glyphomax Plus in 4 gallons of water to prepare a 20 percent solution. In severe infestations, reduce equipment ground speed to ensure that adequate amounts of Glyphomax Plus are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Precautions and Restrictions: Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes or foliage. Allow a minimum of 30 days between last application and harvest of cranberries. For other small fruits and berries, allow a minimum of 14 days between last application and harvest.

Soybeans

Types of Applications: Preplant, preemergence, at-planting, spot treatment, preharvest, selective equipment, hooded sprayers (For Roundup Ready soybeans, refer "Roundup Ready® Crops" section of this label.)

Preplant, Preemergence and At-planting

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting soybeans. Applications must be made prior to emergence of the crop.

37/71

Tank mixtures of Glyphomax Plus with the following herbicide products may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue:

| | | |
|----------------|-----------------|---------------|
| Canopy | Lasso/Alachlor | Pursuit |
| Command | Linex | Pursuit Plus |
| Dual | Lorox/Linuron | Python |
| Dual II Magnum | Lorox Plus | Scepter |
| FirstRate* | Micro-Tech | Sencor/Lexone |
| Frontier | Partner | Squadron |
| Frontrow* | Preview | Turbo |
| Fusion | Pendimax | |
| Gemini | (pendimethalin) | |

For improved burndown, Glyphomax Plus may be tank-mixed with 2,4-D or 2,4-DB herbicide. See the 2,4-D label for intervals between application and planting.

Annual weeds: For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Glyphomax Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of Glyphomax Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall.

Precautions and Restrictions: The tank mix recommendations in this section are not registered in California.

Spot treatment

Specific Use Recommendations: For spot treatments, apply Glyphomax Plus prior to initial pod set in soybeans.

Precautions and Restrictions: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

Preharvest

Specific Use Recommendations: Glyphomax Plus provides weed control when applied prior to harvest of soybeans.

Apply at rates given in the annual, perennial and woody brush tables. Glyphomax Plus may be applied using either aerial or ground spray equipment. For ground applications, apply Glyphomax Plus in 10 to 20 gallons of water per acre. For aerial applications, apply Glyphomax Plus in 3 to 10 gallons of water per acre.

Apply after pods have set and lost all green color. Allow a minimum of 7 days between application and harvest of soybeans. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

Precautions and Restrictions: Do not graze or harvest treated crop for livestock feed within 25 days of last preharvest application. **Do not apply more than 6 quarts per acre of Glyphomax Plus for preharvest applications. Do not apply more than 1 quart per acre of Glyphomax Plus by air.** Do not apply to soybeans grown for seed as a reduction in germination or vigor may occur.

38/71

Selective equipment

Specific Use Recommendations: Glyphomax Plus may be applied through recirculating sprayers, shielded applicators, hooded sprayers, wiper applicators or sponge bars in soybeans. Allow at least 7 days between application and harvest.

Precautions and Restrictions: See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.

Sugarcane

Types of Applications: Preplant, preemergence, spot treatment, fallow, hooded sprayers

Preplant, Preemergence

Specific Use Recommendations: Glyphomax Plus may be applied in or around sugarcane fields or in fields prior to the emergence of plant cane.

Precautions and Restrictions: Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Spot treatment

Specific Use Recommendations: Glyphomax Plus may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, make a 1 percent solution of Glyphomax Plus in water and spray to wet the foliage of vegetation to be controlled. Volunteer or diseased sugarcane should have at least 7 new leaves.

Precautions and Restrictions: Avoid spray contact with healthy cane plants since severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

Fallow treatments

Specific Use Recommendations: Glyphomax Plus may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. Glyphomax Plus may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4 to 5 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage.

Hooded sprayers

Specific Use Recommendations: Glyphomax Plus may be used through hooded sprayers for weed control between the rows of sugarcane. A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution.

Minimize the potential for spray particles to escape from under the hood by operating the sprayer at appropriate ground speeds, nozzle pressures and wind speeds. Operation on rough or sloping ground may result in spray particles escaping from the hood.

When applying to sugarcane that is grown on raised beds, ensure that the hood is designed to completely enclose the spray. If necessary, extend the front and rear flaps of the hoods to reach the ground in furrows between the rows.

39/71

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting the crop. Contact of Glyphomax Plus in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

Precautions and Restrictions: Do not allow treated weeds to come into contact with the crop. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction.

Sunflowers

Types of Applications: Preplant, preemergence

Specific Use Recommendations: Glyphomax Plus may be applied before, during or after planting sunflowers. Applications must be made prior to emergence of the crop.

A tank mixture with Pendimax 3.3 or Prowl (pendimethalin) may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod, or in previous crop residue.

Precautions and Restrictions: Do not apply more than 1 quart of Glyphomax Plus per acre for sunflowers. Make only one preplant or preemergence application per year. Do not feed or graze sunflower forage following application of Glyphomax Plus.

Tree and Vine Crops (General)

Types of Applications: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment (except kiwi), perennial grass suppression

NOTE: This section gives general directions that apply to all citrus crops, tree fruits, tree nuts and vine crops. See the individual crop sections for instructions, preharvest intervals, precautions and restrictions for specific crops.

Glyphomax Plus may be applied in middles, strips and for general weed control in established citrus groves, tree fruit and tree nut orchards, and vineyards. Apply at 1 pint to 5 quarts per acre. Repeat applications may be made up to a maximum of 10.6 quarts per acre per year. Glyphomax Plus may also be used for site preparation prior to transplanting these crops. Allow a minimum of 3 days between application and transplanting. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

Middles (between rows)

Specific Use Recommendations: Glyphomax Plus will control or suppress annual and perennial weeds and ground covers growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

A tank mixture of Glyphomax Plus plus Goal 2XL may be used for annual weeds in middles between rows of citrus crops, tree fruits, tree nuts and vine crops. This mixture is recommended when weeds are stressed or growing in dense populations. Application of 16 to 32 oz/A of Glyphomax Plus plus 3 to 12 oz/A of Goal 2XL will control annual weeds with a maximum height or diameter of 6 inches, including crabgrass, hairy fleabane (*Conyza bonariensis*), common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, common

40/71

cheeseweed (malva), filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). Application of 12 to 32 oz/A of Glyphomax Plus plus 3 to 12 oz/A of Goal 2XL will control common cheeseweed (malva) with a maximum height or diameter of 3 inches.

Strips (in rows)

Specific Use Recommendations: Glyphomax Plus may be applied in rows of tree or vine crops and may also be tank mixed with the following herbicide products:

| | |
|-----------------------------|--------------------|
| Devrinol 50 DF | Princep Caliber 90 |
| Direx 4L | Simazine 4L |
| Goal 2XL | Simazine 80w |
| Karmex DF | Sim-Trol 4L |
| Krovar I | Solicam DF |
| Krovar II | Surflan* A.S. |
| Pendimax (pendimethalin) | |

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Apply 1 pint to 5 quarts of Glyphomax Plus per acre in these tank mixtures. Use rates at the higher end of the recommended rate range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

Perennial grass suppression

Glyphomax Plus will suppress perennial grasses such as bahiagrass, bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of Glyphomax Plus in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of Glyphomax Plus per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply Glyphomax Plus 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of Glyphomax Plus in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of Glyphomax Plus per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

For burndown of bermudagrass, apply 1 to 2 quarts of Glyphomax Plus in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

41/71

For suppression of bermudagrass, apply 6 to 16 fluid ounces of Glyphomax Plus per acre east of the Rocky Mountains and 16 fluid ounces of Glyphomax Plus per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

Selective equipment

Shielded and wiper applicators may be used in tree crops and grapes. Refer to the individual crop sections for time interval between application and harvest.

General Precautions/Restrictions: For citron and olive, apply as a post-directed spray only.

Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees and vines. Contact of Glyphomax Plus with other than matured brown bark can result in serious crop damage.

Avoid painting cut stumps with Glyphomax Plus as injury resulting from root grafting may occur in adjacent trees.

Tree Fruits

Labeled Crops: Apple, apricot, cherry (sweet, sour), crabapple, loquat, mayhaw, nectarine, olive, peach, pear, plum/prune (all), quince

Types of Applications: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment

NOTE: For general use directions, see the "Tree, Nut and Vine (General)" section. The following directions are specific to tree fruits.

Restrictions on application equipment

For cherries, any application equipment listed in this section may be used in all states.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use wiper equipment only.

For **peaches** grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of Glyphomax Plus with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. **Extreme care must be taken to ensure no part of the peach tree is contacted.**

Precautions and Restrictions: Allow a minimum of 1 day between last application and harvest for apple, crabapple, loquat, mayhaw, pear, quince.

4/2/71

Allow a minimum of 17 days between last application and harvest for apricot, cherry, nectarine, olive, peach, plum/prune.

Tree Nuts

Labeled Crops: Almond, beechnut, brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia, pecan, pistachio, walnut (black, English)

Types of Applications: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment

NOTE: For general use directions, see the "tree, Nut and Vine (General)" section. The following directions are specific to tree nuts.

Precautions and Restrictions: Allow a minimum of 3 days between last application and harvest of tree nuts.

Tropical Crops

Labeled Crops: Atemoya, avocado, banana, Barbados cherry (acerola), breadfruit, canistel, carambola, cherimoya, cocoa beans, coconuts, coffee, dates, durian, figs, guava, jaboticaba, jackfruit, longan, lychee, mango, mangosteen, marmaladebox (genip), papaya, passion fruit, persimmon, pineapple, plantain, pomegranate, rambutan, sapodilla, sapote (black, mamey, white), soursop, sugar apple, tamarind, tea.

Specific Use Recommendations: Glyphomax Plus may be applied for general weed control or for site preparation prior to transplanting crops listed in this section. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

Precautions and Restrictions: Allow a minimum of 14 days between last application and harvest of acerola, atemoya, avocado, breadfruit, canistel, carambola, cherimoya, cocoa beans, coconuts, dates, figs, genip, jaboticaba, jackfruit, longan, lychee, mango, mayhaw, passion fruit, persimmon, pomegranate, sapodilla, sapote, soursop, sugar apple, tamarind, and tea.

Allow a minimum of 28 days between last application and harvest of coffee.

Allow a minimum of 1 day between last application and harvest of banana, guava papaya, and plantain.

Do not feed or graze treated pineapple forage following application.

Vegetable Crops

Labeled Crops: Amaranth, arrugula, artichoke (Jerusalem), beans (all), beet greens, garden beets, broccoli (all), brussels sprouts, cabbage (all), cabbage (Chinese), cantaloupe, cardoon, cavalo broccolo, carrot, cauliflower, casaba melon, celery, celery (Chinese), celeriac, celtuce, chard (Swiss), chayote, chervil, chick peas, chicory, chrysanthemum, collards, corn salad, crenshaw melon, cress, cucumber, dandelion, dock (sorrel), eggplant, endive, fennel (florence), garlic, gherkin, ginseng, gourds, ground cherry, guar, honeydew melon, honey ball melon, horseradish, kale, kohlrabi, leek, lentils, lettuce, mango melon, melons (all), mizuna, muskmelon, mustard greens, okra, onion, oriental radish, parsley, parsnips, peas (all), pepinos, pepper (all), Persian melon, potato (Irish), pumpkin, purslane, radish, rape greens,

43/11

rhubarb, rutabaga, salsify, shallot, spinach (all), mustard spinach, squash (summer, winter), sugar beets, sweet potato, tomatillo, tomato, turnip, watercress, watermelon, yams.

Specific Use Recommendations: Glyphomax Plus may be applied prior to the emergence of direct seeded vegetables or prior to transplanting vegetables.

Precautions and Restrictions: When applying Glyphomax Plus prior to transplanting crops into plastic mulch, care must be taken to remove residues of Glyphomax Plus, which could cause crop injury, from the plastic prior to transplanting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or sprinkler system. Applications made at emergence will result in injury or death to emerged seedlings.

For the following crops, apply only prior to planting. Allow at least 3 days between application and planting of cantaloupe, casaba melon, crenshaw melon, cucumber, eggplant, gherkin, gourds, ground cherry, honeydew melon, honey ball melon, mango melon, melons (all), muskmelon, pepper (all), Persian melon, pumpkin, squash (summer, winter), tomatillo, watercress, and watermelon.

Wiper applicators may be used in rutabagas. Allow at least 14 days between application and harvest.

Vine Crops

Labeled Crops: Grapes (raisin, table, wine), kiwi fruit

Types of Applications: General weed control, middles (between rows), strips (in row), selective equipment

NOTE: For general use directions, see the "Tree, Nut and Vine (General)" section. The following directions are specific to vine crops.

Applications should not be made when green shoots, canes or foliage are in the spray zone.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury, or make applications with shielded sprayers or wiper equipment.

Precautions and Restrictions: Allow a minimum of 14 days between last application and harvest.

Roundup Ready® Crops

The following instructions include all applications that can be made onto Roundup Ready® crops during the complete cropping season. Do NOT combine these instructions with other recommendations made for crop varieties that do not contain the Roundup Ready gene, in the "CROPS (ALPHABETICAL)" section of this label.

Glyphomax Plus is recommended for postemergence application only on crop varieties designated as containing the Roundup Ready gene.

- Applying Glyphomax Plus to crop varieties which are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain the Roundup Ready gene, since severe injury or destruction will result.

44/7/11

- Roundup Ready crop varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance is not warranted when Glyphomax Plus is used in conjunction with "brown bag" or seed saved from previous year's crop production and replanted.
- The Roundup Ready designation indicates that the crop variety contains a patented gene that provides tolerance to glyphosate herbicides. Information on Roundup Ready crop varieties may be obtained from your seed supplier.

ATTENTION: Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops, which do not contain the Roundup Ready gene.

See "General Information" and "Application Instructions" sections of this label for essential use directions and restrictions for the application of this product.

Thoroughly clean the spray tank and all lines and filters to eliminate potential contamination from other herbicides prior to mixing and applying Glyphomax Plus.

Note: The following recommendations are based on a clean start at planting by using a burn-down application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burn-down treatment of 16-64 fluid ounces per acre of this product is recommended to control existing weeds prior to crop emergence.

There are no rotational crop restrictions following the application of this product.

Corn with the Roundup Ready® Gene

This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of Glyphomax Plus are not to exceed 1 quart per acre. Sequential in-crop applications of Glyphomax Plus from emergence through the V8 stage or 30 inches must not exceed 2 quarts per acre per growing season.

Maximum Yearly Rates Allowed

Preplant: Maximum amount of Glyphomax Plus which can be applied prior to crop emergence is 5 quarts per acre.

In-crop: Maximum combined total of multiple in-crop applications from emergence through the V8 stage or 30 inches is 2 quarts per acre.

Preharvest: Maximum amount of Glyphomax Plus that can be applied after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest is 1 quart per acre.

Cropping Season: Combined total per year for all applications may not exceed 8 quarts per acre.

When applied as directed, Glyphomax Plus controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of Glyphomax Plus. Applications should be made to actively growing weeds before they reach the maximum size listed in the "Weeds Controlled" section of the label booklet for Glyphomax Plus herbicide.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Glyphomax Plus under hard water conditions, drought conditions

45/11

or when tank mixed with Bullet, Micro-Tech or Partner Herbicides. Refer to the "Mixing" section of the label booklet for proper use instructions. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including fertilizers and micronutrients are not recommended with Glyphomax Plus since this may result in increased potential for crop injury.

Allow a minimum of 50 days between application of Glyphomax Plus and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 10 days between in-crop applications of Glyphomax Plus. **In California, do not graze, harvest or feed corn forage or silage following sequential in-crop applications of Glyphomax Plus on Roundup Ready corn.** There are no rotational crop restrictions following applications of Glyphomax Plus.

ATTENTION: Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops that do not contain the Roundup Ready gene.

Thoroughly clean the spray tank and all lines and filters to eliminate potential contamination from other herbicides prior to mixing and applying this product.

For ground applications: Use the recommended rates of Glyphomax Plus in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of Glyphomax Plus in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See the "Annual and Perennial Weeds Rate Tables" in this label. **Avoid drift - do not apply during inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent vegetation, appropriate buffer zones must be maintained.**

Weed Control Recommendations

Apply 24 to 32 fluid ounces of Glyphomax Plus herbicide per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the "Annual Weeds Rate Table" for rate recommendations for specific annual weeds. Glyphomax Plus herbicide applied at up to 1 quart per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpet creeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "Perennial Weeds Rate Table".

Preemergence followed by Postemergence Weed Control Program

This product may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of Glyphomax Plus should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. A single in-crop application of Glyphomax Plus at the recommended rate will provide control of emerged weeds listed on this label. This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first.

Postemergence Only Weed Control Program

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of Glyphomax Plus should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of Glyphomax Plus at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied postemergence to Roundup Ready

4/6/71

corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first.

This product may be applied in tank mixture with a labeled rate of FulTime, Surpass, Surpass 100, TopNotch, Bicep II, Bicep Lite II Magnum, Bicep II Magnum, Dual II, Dual II Magnum, Frontier, Guardsman, LeadOff, Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Hornet WDG, Micro-Tech, Bullet, Partner, Permit or Atrazine herbicides. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines - the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

| Tank Mix Partner | Maximum Height Of Corn For Application |
|---|--|
| Bicep II Bicep II Magnum Bicep Lite II Magnum Bullet † Dual II Dual II Magnum Micro-Tech † Partner † | 5 inches |
| Frontier Guardsman LeadOff | 8 inches |
| FulTime Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6 Surpass EC TopNotch | 11 inches |
| Hornet Hornet WDG | 20 inches |
| Permit | 24 inches |
| Atrazine | 12 inches |

† Bullet, Micro-Tech and Partner are not registered for use as a postemergence application in Texas.

Soybeans with the Roundup Ready® Gene

Specific Use Directions

This product may be applied postemergence to Roundup Ready soybeans from the cracking stage throughout flowering. Allow a minimum of 14 days between application and harvest of soybeans.

Maximum Allowable Application Rates:

- Combined total for all applications 8 quarts per acre
- Preplant, preemergence applications 5 quarts per acre

47/71

- Total in-crop applications from cracking throughout flowering 3 quarts per acre
- Maximum preharvest application rate 1 quart per acre

When applied as directed, Glyphomax Plus will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of Glyphomax Plus.

Precautions and Restrictions: The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in-crop application is 2 quarts per acre. The maximum combined total of this product that can be applied during flowering is 2 quarts per acre. Allow a minimum of 14 days between final application and harvest of soybeans.

For ground applications: Use the recommended rates of Glyphomax Plus in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles that provide a flat fan pattern. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of Glyphomax Plus in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of Glyphomax Plus per acre. Do not apply during low level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. Maintain appropriate buffer zones to prevent injury to adjacent desirable vegetation.

Annual Weed Rate Tables

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the "Annual Weeds Rate Tables" section for rate recommendations for specific annual weeds.

Dow AgroSciences will not warrant crop safety or weed control when Roundup Ready soybeans are treated with herbicides not specified on this supplemental label. Because of the potential for; 1) crop injury, 2) poor weed control from antagonism, and/or 3) rotational crop restrictions, herbicides not specified on this supplemental label should not be used, whether applied preemergence or applied postemergence as a tank mixture with Glyphomax Plus herbicide.

Up to 64 fluid ounces per acre of Glyphomax Plus may be used in any single application for control of annual weeds, where heavy weed densities exist.

Midwest/ Mid-Atlantic Recommendations

Narrow row or drilled soybeans: A single in-crop application of Glyphomax Plus will provide effective control of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/acre), on 4-8" weeds is recommended. Weeds will generally be 4 - 8" tall 3 to 5 weeks after planting. If the initial application is delayed and weeds are 8-18" tall, use 48 fl oz/acre for best results.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of Glyphomax Plus at 24 to 32 fluid ounces per acre may be necessary to control late flushes of weeds. The combined total application in-crop must not exceed 96 fluid ounces per acre.

Wide row soybeans: An in-crop application of Glyphomax Plus will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/acre), on 4-8" weeds is recommended. Weeds will generally be 4-8" tall 3 to 5 weeks after planting. If new flushes of weeds occur, they can be controlled by sequential applications of Glyphomax Plus.

48/71

Initial Treatment and Sequential Applications (if Needed)

| Weed Height (inches) | Rate (fl oz/acre) |
|-------------------------|----------------------|
| 1 - 3 | 24 |
| 4 - 7 | 32 |
| 8 - 12 | 48 |

Giant ragweed: Apply 32 fl oz/acre when the weed is 8-12" tall to avoid the need for sequential application.

Black nightshade, Pennsylvania smartweed, velvetleaf and waterhemp. Apply 1 quart per acre to weeds 3-6 inches tall and 48 fl oz when weeds are up to 12 inches tall. For morningglory species, apply 32 fl oz when weeds are up to 4 inches tall, and 48 fl oz when weeds are up to 6 inches tall.

Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed, with multiple germination times may require a sequential application of Glyphomax Plus. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 24 fluid ounces of Glyphomax Plus per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

Southeast Recommendations

Narrow row, drilled, or wide-row soybeans: An in-crop application of Glyphomax Plus will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/acre), on 3-6" weeds is recommended. Weeds will generally be 3-6" tall 2 to 3 weeks after planting.

Initial Treatment

| Weed Height (inches) | Rate (fl oz/acre) |
|-------------------------|----------------------|
| 3 - 6 | 32 |
| 6 - 12 | 48 |

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of Glyphomax Plus at 16 to 32 fluid ounces per acre may be necessary to control late flushes of weeds.

Sequential Application (if needed)[†]

| Weed Height (inches) | Rate (fl oz/acre) |
|-------------------------|----------------------|
| 2 - 3 | 16 |
| 3 - 5 | 24 |
| 6 - 12 | 32 |

Florida pusley, hemp sesbania and spurred anoda: Apply 32 fl oz/acre to weeds 2-4" for the initial application. Apply 32 oz/acre when these weeds are 3-6" tall if a sequential application is necessary.

4/9/71

Morningglory, black nightshade, groundcherry, and Pennsylvania smartweed: Apply 24 fl oz/acre on 1-3" weeds, 32 fl oz/acre on 3-6" weeds, or 48 fl oz/acre on 6-12" weeds for the initial application.

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of Glyphomax Plus. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of Glyphomax Plus per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

Delta/Mid-South Recommendations

Narrow row, drilled, or wide row soybeans: An in-crop application of Glyphomax Plus will provide effective control of the initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. For best results, an initial application of 32 fluid ounces per acre (fl oz/acre), on 2-4" weeds is recommended. Weeds will generally be 2-4" tall 2 to 3 weeks after planting.

Initial Treatment

| Weed Height (inches) | Rate (fl oz/acre) |
|-------------------------|----------------------|
| 2 - 4 | 32 |
| 5 - 12 | 48 |

Sequential Application (if needed)[†]

| Weed Height (inches) | Rate (fl oz/acre) |
|-------------------------|----------------------|
| 2 - 3 | 16 |
| 3 - 6 | 24 |
| 6 - 12 | 32 |

Hemp sesbania and spurred anoda: Apply a sequential treatment of 32 fl oz/acre on 3-6" weeds if necessary

Some weeds, such as black nightshade, broadleaf signaigrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of Glyphomax Plus. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of Glyphomax Plus per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

Perennial Weeds Rate Recommendations

A rate of 32 to 64 fluid ounces per acre (single or multiple applications) of Glyphomax Plus will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly.

For best results, allow perennial weed species to reach a height of at least 6" before spraying. For additional information on perennial weeds, see the "Perennial Weeds Rate Table" section. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

50/71

Farmsteads

Labeled Use Sites: Glyphomax Plus may be used in farmsteads (including building foundations, along and in fences, dry ditches, dry canals, along ditchbanks, farm roads, shelterbelts, prior to landscape plantings and equipment storage areas).

Types of Applications: General nonselective weed control, trim-and-edge, chemical mowing, cut stumps, and habitat management.

General nonselective weed control, Trim-and-edge

Glyphomax Plus may be tank mixed with the following herbicide products. Refer to these product labels for labeled application sites and application rates. For annual weeds, use 1 quart per acre of Glyphomax Plus when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are greater than 6 inches tall. For perennial weeds, apply 2 to 5 quarts per acre in these tank mixes. For tank mixtures of Glyphomax Plus with these products through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "Hand-Held And High Volume Equipment" section of this label for recommended rates.

| | |
|------------------|----------------|
| Arsenal | Plateau |
| Banvel (dicamba) | Princep DF |
| Barricade 35WG | Princep Liquid |
| diuron | Ronstar 50W |
| Endurance | Sahara |
| Escort | simazine |
| Karmex DF | Surflan* |
| Krovar I DF | Telar |
| Oust | Vanquish |
| Pendulum 3.3 EC | 2,4-D |
| Pendulum WDG | |

Tank mixtures of Glyphomax Plus with Banvel and 2,4-D may not be applied by air in California.

Chemical mowing

Perennials: Glyphomax Plus will suppress perennial grasses listed in this section to serve as a substitute for mowing. Apply Glyphomax Plus at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces of Glyphomax Plus per acre when treating tall fescue, fine fescue, orchardgrass or quackgrass covers. Use 6 fluid ounces of Glyphomax Plus per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

Precautions and Restrictions: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

5/7/11

| Weed Species | Region | Rate of Glyphomax Plus (Fluid Ounces Per Acre) | | | | | |
|---|--------|---|------|------|-----|-----|-----|
| | | 12 | 16 | 24 | 32 | 40 | 48 |
| | | Maximum Height/Length | | | | | |
| annoda, spurred | | - | 1" | 2" | 3" | 5" | 8" |
| barley | | - | 18" | 18"+ | - | - | - |
| barnyardgrass | South | - | 3" | 5" | 7" | 9" | 12" |
| | North | - | - | 6" | 12" | - | - |
| bassia, fivehook | | - | - | - | 6" | - | - |
| bittercress | | - | 12" | 20" | - | - | - |
| bluegrass, annual | | - | 10" | - | - | - | - |
| brome, downy | | 6" | - | - | - | - | - |
| brome, Japanese | | - | 6" | - | 24" | - | - |
| browntop panicum | | - | 6" | 8" | 12" | - | 24" |
| burcucumber | | - | 6" | 12" | - | - | - |
| buttercup | | - | 12" | 20" | - | - | - |
| Carolina foxtail | | - | 20" | - | - | - | - |
| Carolina geranium | | - | - | - | 4" | - | 9" |
| carpetweed | | - | - | 6" | 12" | - | - |
| cheat | | - | 6" | 20" | - | - | - |
| chervil | | - | 20" | - | - | - | - |
| chickweed | | - | 12" | 18" | - | - | - |
| cocklebur | | - | 12" | 18" | 24" | - | - |
| copperleaf, hophornbeam | | - | 1" | 2" | 3" | 4" | 6" |
| copperleaf, Virginia | | - | 1" | 2" | 3" | 4" | 6" |
| corn | | - | 12" | 20" | - | - | - |
| corn speedwell | | - | 12" | - | - | - | - |
| crabgrass | | - | 12" | 18" | - | - | - |
| cutleaf evening primrose | | - | - | - | 3" | 3" | 6" |
| dwarf dandelion | | - | 20" | - | - | - | - |
| eastern mannagrass | | - | 8" | 12" | - | - | - |
| eclipta | | - | 4" | 8" | 12" | - | - |
| fall panicum | South | - | 4" | 6" | 8" | 12" | 24" |
| | north | - | 6" | 12" | 18" | - | - |
| falsedandelion | | - | 20" | - | - | - | - |
| falseflax, smallseed | | - | 12" | - | - | - | - |
| fiddleneck | | - | - | - | 6" | 6" | 12" |
| field pennycress | | - | 6" | 12" | - | - | - |
| filaree | | - | - | - | - | - | 12" |
| fleabane, annual | | - | 6" | 20" | - | - | - |
| fleabane, hairy (<i>conyza bonariensis</i>) | | - | 6" | - | - | - | - |
| fleabane, rough | | - | 3" | 6" | 12" | - | - |
| Florida pusley | | - | - | - | 4" | 4" | 6" |
| foxtail | South | - | 8" | 12" | 20" | - | - |
| | North | 18" | 18"+ | - | - | - | - |
| goatgrass, jointed | | - | 6" | - | - | - | - |
| goosegrass | | - | 3" | 5" | 8" | - | 18" |
| grain sorghum (milo) | | - | 6" | 12" | 20" | - | - |
| groundsel, common | | - | 6" | - | - | - | - |
| hemp sesbania | | - | - | 2" | 4" | 6" | 8" |

53/71

| | | | | | | | |
|---|-------|-----|------------|------|-----|-----|-----|
| henbit | | - | - | - | 6" | - | 20" |
| horseweed/marestail (<i>conyza canadensis</i>) | South | - | - | 12" | 30" | - | - |
| | North | - | 6" | 12" | 18" | - | - |
| itchgrass | | - | 6" | 12" | 18" | - | - |
| jimsonweed | | - | - | - | 6" | 6" | 12" |
| johnsongrass (seedling) | South | - | - | - | 18" | - | - |
| | North | - | 12" | 18" | - | - | - |
| jungerice | | - | 3" | 5" | 7" | 9" | 12" |
| knotweed | | - | 3" | 8" | 12" | - | 20" |
| kochia ¹ | | - | 3 to 6" | 12" | - | - | - |
| lambsquarters | | - | 6" | 8" | 12" | - | 20" |
| little barley | | - | 20" | - | - | - | - |
| London rocket | | - | 6" | - | - | - | - |
| mayweed | | - | - | 2" | 6" | 12" | 18" |
| morningglory (<i>ipomoea spp.</i>) | | - | - | 2" | 4" | - | 6" |
| mustard, blue | | 6" | - | - | - | - | - |
| mustard, tansy | | 6" | 12" | 20" | - | - | - |
| mustard, tumble | | 6" | - | - | - | - | - |
| mustard, wild | | 6" | 12" | 18" | - | - | - |
| nightshade, black | | 6" | 12" | - | - | - | - |
| nightshade, hairy | | - | 6" | 12" | - | - | - |
| oats | | - | - | 6" | 20" | - | - |
| pigweed | | - | 12" | 18" | 24" | - | - |
| prickly lettuce | | - | 6" | 12" | 20" | - | - |
| purslane | | - | - | - | 6" | 6" | 12" |
| ragweed, common | South | - | 4" | 6" | 8" | - | 11" |
| | North | - | 6" | 12" | 18" | - | - |
| ragweed, giant | | - | - | 4" | 6" | - | 11" |
| red rice | | - | - | - | 4" | - | - |
| Russian thistle | | - | 6" | - | - | - | - |
| rye | South | - | 6" | 20" | 60" | - | - |
| | North | - | 18" | 18"+ | - | - | - |
| ryegrass | | - | - | - | 6" | - | 7+" |
| sandbur, field | | 12" | - | - | - | - | - |
| shattercane | | - | 12" | 18" | - | - | - |
| shepherd's-purse | | - | 6" | 12" | - | - | - |
| sicklepod | | - | - | 2" | 4" | - | 8" |
| signalgrass, broadleaf | | - | 3" | 5" | 7" | 9" | 12" |
| smartweed, ladysthumb | | - | 4" | 6" | 8" | - | 12" |
| smartweed, pennsylvania | | - | 4" | 6" | 8" | - | 12" |
| sowthistle, annual | | - | - | - | 6" | - | 12" |
| spanishneedles | | - | - | - | 8" | - | 18" |
| speedwell, purslane | | - | 12 | - | - | - | - |
| sprangletop | | - | 6" | 12" | 20" | - | - |
| spurge, prostrate | | - | 6" | 12" | 20" | - | - |
| spurge, spotted | | - | 6" | 12" | 20" | - | - |
| spurry, umbrella | | 6" | - | - | - | - | - |
| stinkgrass | | 12" | - | - | - | - | - |
| sunflower | | - | 12" | 18" | - | - | - |

54/71

| | | | | | | | |
|-----------------------|-------|----|-----|------|-----|-----|-----|
| teaweed/ prickly sida | | 1" | 2" | 3" | 4" | 6" | |
| Texas panicum | | 6" | 8" | 12" | - | 24" | |
| velvetleaf | South | - | 2" | 3" | 4" | 5" | 8" |
| | North | - | 3" | 6" | 12" | - | - |
| Virginia pepperweed | | - | 18" | - | - | - | - |
| waterhemp | | - | - | 6" | 12" | - | - |
| wheat | South | - | 6" | 30" | - | - | - |
| | North | - | 18" | 18"+ | - | - | - |
| wheat (over-wintered) | | - | 6" | 18" | - | - | - |
| wild oats | | - | 12" | - | - | - | - |
| wild proso millet | | - | - | 6" | 12" | 12" | 18" |
| witchgrass | | - | 12" | - | - | - | - |
| woolly cupgrass | | - | 6" | 12" | - | - | - |
| yellow rocket | | - | - | 12" | 20" | - | - |

¹ Do not treat kochia in the button stage.

Annual Weeds Rate Table, West Region

| Weed Species | Rate of Glyphomax Plus (Fluid Ounces Per Acre) | | | | |
|---|---|-----|----|-----|-----|
| | 12 | 16 | 24 | 32 | 48 |
| | Maximum Height/Length | | | | |
| barley | 12" | - | - | - | - |
| barnyardgrass | 6" | - | - | - | - |
| bluegrass, annual | 6" | - | - | - | - |
| bluegrass, bulbous | - | 6" | - | - | - |
| brome, downy ¹ | 6" | - | - | - | - |
| buttercup | - | 12" | - | - | - |
| cheat | - | 6" | - | - | - |
| chickweed | - | 6" | - | - | - |
| cocklebur | - | 12" | - | - | - |
| corn | - | 6" | - | - | - |
| crabgrass | - | 12" | - | - | - |
| dwarf dandelion | - | 12" | - | - | - |
| fall panicum | - | 12" | - | - | - |
| false flax, smallseed | - | 12" | - | - | - |
| field pennycress | - | 6" | - | - | - |
| filaree | - | - | - | - | 12" |
| fleabane, hairy (<i>conyza bonariensis</i>) | - | 6" | - | - | - |
| Florida pusley | - | - | - | 12" | - |
| foxtail | (8 fl. oz. for up to 12") | | | | |
| goatgrass, jointed | - | 6" | - | - | - |
| groundsel, common | - | 6" | - | - | - |
| henbit | - | 6" | - | - | - |
| horseweed/marestail (<i>conyza canadensis</i>) | - | 6" | - | - | - |
| johnsongrass, seedling | - | 12" | - | - | - |
| lambsquarters | - | 6" | - | - | - |
| London rocket | - | 6" | - | - | - |

55/11

| | | | | | |
|--------------------------------------|-----|-----|---|---|---|
| morningglory (<i>ipomoea spp.</i>) | - | 2" | - | - | - |
| mustard, blue | 6" | - | - | - | - |
| mustard, tansy | 6" | - | - | - | - |
| mustard, tumble | 6" | - | - | - | - |
| mustard, wild | 6" | - | - | - | - |
| pigweed | - | 12" | - | - | - |
| rye | 12" | - | - | - | - |
| ryegrass, Italian | - | 6" | - | - | - |
| sandbur, field | 12" | - | - | - | - |
| shattercane | 12" | - | - | - | - |
| shepherd's-purse | - | 6" | - | - | - |
| sowthistle, annual | - | 6" | - | - | - |
| spurge, annual | - | 6" | - | - | - |
| stinkgrass | 12" | - | - | - | - |
| Texas panicum | - | 12" | - | - | - |
| wheat | 18" | - | - | - | - |
| wild oats | - | 12" | - | - | - |
| witchgrass | - | 12" | - | - | - |

¹For control of downy brome in no-till systems, use 16 fluid ounces per acre.

Annual Weeds--Water Carrier Volumes of 10 to 40 Gallons per Acre

Apply 1 to 1.5 quarts of Glyphomax Plus per acre. Use 1 quart per acre if weeds are less than 6 inches tall and 1.5 quarts per acre if weeds are over 6 inches tall. These rates will provide control of weeds listed in the annual weed control tables when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Annual Weeds -- Tank Mixtures with 2,4-D, Dicamba or Tordon 22K

Application of 12 to 24 fluid ounces of this product plus 0.25 pound a.i. of dicamba or 0.5 pound a.i. of 2,4-D or 1 to 2 ounces of Tordon 22K per acre will control the following weeds with the maximum height or length indicated: 6" -- prickly lettuce, marestail/horseweed (*Conyza canadensis*), morningglory (*Ipomoea spp.*), kochia (dicamba only); wild buckwheat (Tordon 22K only); 12" -- cocklebur, lambsquarters, pigweed, Russian thistle.

Application of 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. of 2,4-D per acre will control the following weeds when they are a maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf.

Application of 12 fluid ounces of Glyphomax Plus plus 0.25 pound a.i. of dicamba or 0.5 pound a.i. of 2,4-D per acre will control foxtail up to 18".

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

Tank mixtures of Glyphomax Plus with Banvel (dicamba), Tordon 22K or 2,4-D may not be applied by air in California.

56/71

Annual Weeds-Tank Mixtures with Atrazine or Bladex for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound atrazine per acre.

Application of 16 ounces of this product plus 1 to 2 pounds of atrazine or 2.4 to 4 pounds of cyanazine per acre will control the following weeds: barnyardgrass (barnyardgrass requires 26 ounces of Glyphomax Plus for control), downy brome, green foxtail, lambsquarters, prickly lettuce (*Lactuca serriola*), tansy mustard, pigweed, field sandbur (*Cenchrus* spp.), stinkgrass, Russian thistle (*Salsola kali*), volunteer wheat, witongrass (*Panicum capillare*) and kochia (for Kochia, add 4 fl oz of Banvel/dicamba for control).

Perennial Weeds Rate Table (Alphabetically By Species)

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage.

Best results are obtained when soil moisture is adequate for active weed growth.

| Weed Species | Rate (qt/acre) | Water Volume (gpa) | Hand-Held (% Solution) |
|---|----------------|--------------------|------------------------|
| Alfalfa | 1 - 2 | 3 - 10 | 2% |
| Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up. | | | |
| Alligatorweed | 4 | 3 -20 | 1.5% |
| Partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain control. | | | |
| Anise (fennel) | - | - | 1 - 2% |
| Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. | | | |
| Bahiagrass | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants have reached the early head stage. | | | |
| Bentgrass | 1.5 | 10 - 20 | 2% |
| For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is | | | |

57/71

| | | | |
|---|------------------|---------------|-----------|
| recommended for best results. | | | |
| Bermudagrass | 3 - 5 | 3 - 20 | 2% |
| For control, apply 5 quarts of Glyphomax Plus per acre. For partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. | | | |
| Bermudagrass, water (knotgrass) | 1 - 1.5 | 5 - 10 | 2% |
| Apply 1.5 quarts of Glyphomax Plus in 5 to 10 gallons of water per acre. Apply when water bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field. Fall applications only: Apply 1 quart of Glyphomax Plus in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water bermudagrass that is 12 to 18 inches in length. Glyphomax Plus is not registered in California for use on water bermudagrass. | | | |
| Bindweed, field | 0.5 - 5.0 | 3 - 20 | 2% |
| Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. For control, apply 4 to 5 quarts of Glyphomax Plus per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Also for control, apply 2 quarts of Glyphomax Plus plus 0.5 pound a.i. of dicamba in 10 to 20 gallons of water per acre. Do not apply by air. For suppression on irrigated agricultural land, apply 1 to 2 quarts of Glyphomax Plus plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth. For suppression, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. of 2,4-D or 0.25 pound a.i. of dicamba in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length. In California only, apply 1 to 5 quarts of Glyphomax Plus per acre. The actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage. | | | |
| Bluegrass, Kentucky | 1 - 2 | 3 - 40 | 2% |
| Apply 2 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. | | | |
| Blueweed, Texas | 3 - 5 | 3 - 40 | 2% |
| Apply 4 to 5 quarts of Glyphomax Plus per acre west of the Mississippi River and 3 to 4 quarts per | | | |

58/11

| | | | |
|--|--------------|---------------|-----------------|
| acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. | | | |
| Brackenfern | 3 - 4 | 3 - 40 | 1 - 1.5% |
| Apply to fully expanded fronds, which are at least 18 inches long. | | | |
| Bromegrass, smooth | 1 - 2 | 3 - 40 | 2% |
| Apply 2 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. | | | |
| Bursage, woolly-leaf | - | 3 - 20 | 2% |
| For control, apply 2 quarts of Glyphomax Plus plus 0.5 lb a.i. of dicamba per acre. For partial control, apply 1 quart of Glyphomax Plus plus 0.5 lb a.i. of dicamba per acre. Apply when plants are producing new active growth, which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering. | | | |
| Canarygrass, reed | 2 - 3 | 3 - 40 | 2% |
| For best results, apply when most plants have reached the boot-to-head stage of growth. | | | |
| Cattail | 3 - 5 | 3 - 40 | 2% |
| Apply when most plants have reached the early head stage. | | | |
| Clover; red, white | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants have reached the early bud stage. | | | |
| Cogongrass | 3 - 5 | 2 - 40 | 2% |
| Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control. | | | |
| Dallisgrass | 3 - 5 | 2 - 20 | 2% |
| Apply when most plants have reached the early head stage. | | | |
| Dandelion | 3 - 5 | 3 - 40 | 2% |
| Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre. | | | |
| Dock, curly | 3 - 5 | 3 - 40 | 2% |
| Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre. | | | |
| Dogbane, hemp | 4 | 3 - 40 | 2% |
| Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. | | | |

59/71

| | | | |
|---|------------------|---------------|-------------------|
| For suppression, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred. | | | |
| Fescue (Except tall) | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants have reached the early head stage. | | | |
| Fescue, tall | 1 - 3 | 3 - 40 | 2% |
| Apply 3 quarts of Glyphomax Plus per acre when most plants have reached boot-to-early seedhead stage of development. | | | |
| Fall applications only: Apply 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of Glyphomax Plus will improve long-term control and control seedlings germinating after fall treatments or the following spring. | | | |
| Guineagrass | 3 | 3 - 40 | 1% |
| Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. | | | |
| Horsenettle | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants have reached the early bud stage. | | | |
| Horseradish | 4 | 3 - 40 | 2% |
| Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall. | | | |
| Iceplant | - | - | 1.5 - 2.0% |
| Iceplant should be at or beyond the early bud stage of growth. Thorough coverage is necessary for best control. | | | |
| Jerusalem artichoke | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early bud stage. | | | |
| Johnsongrass | 0.5 - 3.0 | 3 - 40 | 1% |
| In annual cropping systems apply 1 to 2 quarts of Glyphomax Plus per acre. Apply 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Glyphomax Plus when applying 10 to 40 gallons of water per acre. In noncrop or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre. | | | |
| For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using the 1 quart per acre rate. | | | |
| For burndown of Johnsongrass, apply 1 pint of Glyphomax Plus in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage. | | | |
| Spot treatment (partial control or suppression): Apply a 1 percent solution of Glyphomax Plus when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete. | | | |
| Kikuyugrass | 2 - 3 | 3-40 | 2% |
| Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth). Allow 3 or more days after application before tillage. | | | |

60/71

| | | | |
|--|----------------|---------------|------------------|
| Knapweed | 4 | 3-40 | 2% |
| Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall. | | | |
| Lantana | - | - | 1 - 1.25% |
| Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. | | | |
| Lespedeza | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants have reached the early bud stage. | | | |
| Milkweed, common | 3 | 3 - 40 | 2% |
| Apply when most plants have reached the late bud to flower stage of growth. | | | |
| Muhly, wirestem | 1 - 2 | 3 - 40 | 2% |
| Use 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Glyphomax Plus when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. | | | |
| Mullein, common | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early bud stage. | | | |
| Napiergrass | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early head stage. | | | |
| Nightshade, silverleaf | 2 | 3 - 10 | 2% |
| Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. | | | |
| Nutsedge; purple, yellow | 0.5 - 3 | 3 - 40 | 1 - 2% |
| <p>Apply 3 quarts of Glyphomax Plus per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets, which have not germinated, will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.</p> <p>Sequential applications: 1 to 2 quarts of Glyphomax Plus in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.</p> <p>For partial control of existing plants, apply 1 pint to 2 quarts of Glyphomax Plus in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.</p> | | | |
| Orchardgrass | 1 - 2 | 3 - 40 | 2% |
| Apply 2 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. | | | |

6/1/71

| | | | |
|---|-----------------|----------------|-----------------|
| Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of Glyphomax Plus in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results. | | | |
| Pampasgrass | - | - | 1.5 - 2% |
| Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control. | | | |
| Paragrass | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early head stage. | | | |
| Phragmites | 3 - 5 | 10 - 40 | 1 - 2% |
| For partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop. | | | |
| Poison hemlock | - | - | 1 - 2% |
| Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. | | | |
| Pokeweed, common | - | 3 - 40 | 2% |
| Apply to actively growing plants up to 24 inches tall. | | | |
| Quackgrass | 1 - 3 | 3 - 40 | 2% |
| In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of Glyphomax Plus. Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results. In pastures, sods or noncrop areas where deep tillage does not follow application: Apply 2 to 3 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall. | | | |
| Redvine | 0.75 - 2 | 5 - 10 | 2% |
| For suppression, apply 24 fluid ounces of Glyphomax Plus per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost. | | | |
| Reed, giant | - | - | 2% |
| Best results are obtained when applications are made in late summer to fall. | | | |
| Ryegrass, perennial | 1 - 3 | 3 - 40 | 1% |
| In annual cropping systems apply 1 to 2 quarts of Glyphomax Plus per acre. Apply 1 quart of Glyphomax Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Glyphomax Plus when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not practiced, | | | |

02/71

| | | | |
|---|--------------|----------------|-----------|
| apply 2 to 3 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre. | | | |
| For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using the 1 quart per acre rate. | | | |
| Smartweed, swamp | 3 - 5 | 3 - 40 | 2% |
| Apply when most plants have reached the early bud stage of growth. | | | |
| Also for control, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. | | | |
| Sowthistle, perennial | 2 - 3 | 3 - 40 | 2% |
| Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage. | | | |
| Spurge, leafy | - | 3 - 10 | 2% |
| For suppression, apply 16 fluid ounces of Glyphomax Plus plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. | | | |
| Starthistle, yellow | 2 | 10 - 40 | 2% |
| Best results are obtained when applications are made during the rosette, bolting and early flower stages. | | | |
| Sweet potato, wild | - | - | 2% |
| Partial control. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required. | | | |
| Thistle, artichoke | - | - | 2% |
| Partial control. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required. | | | |
| Thistle, Canada | 2 - 3 | 3 - 40 | 2% |
| Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Glyphomax Plus. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage. | | | |
| For suppression, apply 1 quart of Glyphomax Plus, or 1 pint of Glyphomax Plus plus 0.5 pound a.i. 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage. | | | |
| Timothy | 2 - 3 | 3 - 40 | 2% |
| For best results, apply when most plants have reached the boot-to-head stage of growth. | | | |
| Torpedograss | 4 - 5 | 3 - 40 | 2% |
| For partial control. Apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. | | | |

6/3/71

| | | | |
|---|--------------|---------------|-----------|
| Trumpet creeper | 2 | 5 - 10 | 2% |
| Partial control. Apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost. | | | |
| Vaseygrass | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early head stage. | | | |
| Velvetgrass | 3 - 5 | 3 - 20 | 2% |
| Apply when most plants are in the early head stage. | | | |
| Wheatgrass, western | 2 - 3 | 3 - 40 | 2% |
| For best results, apply when most plants have reached the boot-to-head stage of growth. | | | |

Woody Brush And Trees Rate Table (Alphabetically By Species)

Apply Glyphomax Plus after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

| Weed Species | Rate (qt/acre) | Water Volume (gpa) | Hand-Held (% Solution) |
|--|---------------------------|-------------------------------|-----------------------------------|
| Alder For control | 3 - 4 | 3 - 40 | 1 - 1.5% |
| Ash Partial control | 2 - 5 | 3 - 40 | 1 - 2% |
| Aspen, quaking For control | 2 - 3 | 3 - 40 | 1 - 1.5% |
| Bearmat (Bearclover) For partial control | 2 - 5 | 3 - 40 | 1 - 2% |
| Beech Partial control | 2 - 5 | 3 - 40 | 1 - 2% |

64/71

| | | | |
|---|--------------|----------------|-----------------|
| Birch | 2 | 3 - 40 | 1% |
| For control | | | |
| Blackberry | 3 - 4 | 10 - 40 | 1 - 1.5% |
| For control. Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4 percent solution of Glyphomax Plus. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of Glyphomax Plus in 10 to 40 gallons of water per acre. | | | |
| Blackgum | 2 - 5 | 3 - 40 | 1 - 2% |
| For control | | | |
| Bracken | 2 - 5 | 3 - 40 | 1 - 2% |
| For control | | | |
| Broom; French, Scotch | - | - | 1.5 - 2% |
| For control | | | |
| Buckwheat, California | - | - | 1 - 2% |
| For partial control. Thorough coverage of foliage is necessary for best results. | | | |
| Cascara | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Catsclaw | - | - | 1 - 1.5% |
| Partial control | | | |
| Ceanothus | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Chamise | - | - | 1% |
| For control. Thorough coverage of foliage is necessary for best results. | | | |
| Cherry; bitter, black, pin | 2 - 3 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Coyote brush | - | - | 1 - 1.5% |
| For control. Apply when at least 50 percent of the new leaves are fully developed. | | | |
| Dogwood | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Elderberry | 2 | 3 - 40 | 1% |
| For control | | | |
| Elm | 2 - 5 | 3 - 40 | 1 - 2% |

| | | | |
|---|--------------|---------------|-----------------|
| Peppertree) | | | |
| Partial control | | | |
| Gorse | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Hasardia | - | - | 1 - 2% |
| Partial control. Thorough coverage of foliage is necessary for best results. | | | |
| Hawthorn | 2 - 3 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Hazel | 2 | 3 - 40 | 1% |
| For control | | | |
| Hickory | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Honeysuckle | 3 - 4 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Hornbeam, American | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Kudzu | 4 | 3 - 40 | 2% |
| For control. Repeat applications may be required to maintain control. | | | |
| Locust, black | 2 - 4 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Madrone resprouts | - | - | 2% |
| Partial control. Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments. | | | |
| Manzanita | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Maple, red | 2 - 4 | 3 - 40 | 1 - 1.5% |
| For control, apply a 1 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of Glyphomax Plus per acre. | | | |
| Maple, sugar | - | - | 1 - 1.5% |
| For control. Apply when at least 50 percent of the new leaves are fully developed. | | | |

66/71

| | | | |
|---|-------|--------|----------|
| Monkey flower | - | - | 1 - 2% |
| Partial control. Thorough coverage of foliage is necessary for best results. | | | |
| Oak; black, white | 2 - 4 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Oak, post | 3 - 4 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Oak; northern, pin | - | - | 1 - 1.5% |
| For control. Apply when at least 50 percent of the new leaves are fully developed. | | | |
| Oak; southern red | 2 - 3 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Persimmon | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Pine | 2 - 5 | 3 - 40 | 1 - 2% |
| For control | | | |
| Poison ivy/ Poison oak | 4 - 5 | 3 - 40 | 1 - 2% |
| For control. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color. | | | |
| Poplar, yellow | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Redbud, eastern | 2 - 5 | 3 - 40 | 1 - 2% |
| For control | | | |
| Rose, multiflora | 2 | 3 - 40 | 1% |
| For control. Treatments should be made prior to leaf deterioration by leaf-eating insects. | | | |
| Russian olive | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Sage, black | - | - | 1% |
| For control. Thorough coverage of foliage is necessary for best results. | | | |
| Sage, white | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Sage brush, California | - | - | 1% |
| For control. Thorough coverage of foliage is necessary for best results. | | | |
| Salmonberry | 2 | 3 - 40 | 1% |
| For control | | | |
| Salt-cedar | 2 - 5 | 3 - 40 | 1 - 2% |

6/7/71

| | | | |
|--|--------------|---------------|-----------------|
| For control | | | |
| Sassafras | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Sourwood | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Sumac; poison, smooth, winged | 2 - 4 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Sweetgum | 2 - 3 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Swordfern | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Tallowtree, Chinese | - | - | 1% |
| For control. Thorough coverage of foliage is necessary for best results. | | | |
| Tan oak resprouts | - | - | 2% |
| For partial control. Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications. | | | |
| Thimbleberry | 2 | 3 - 40 | 1% |
| For control | | | |
| Tobacco, tree | - | - | 1 - 2% |
| Partial control | | | |
| Trumpet creeper | 2 - 3 | 3 - 40 | 1 - 1.5% |
| For control | | | |
| Vine maple | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Virginia creeper | 2 - 5 | 3 - 40 | 1 - 2% |
| For control | | | |
| Waxmyrtle, southern | 2 - 5 | 3 - 40 | 1 - 2% |
| Partial control | | | |
| Willow | 3 | 3 - 40 | 1% |
| For control | | | |

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase

price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that Glyphomax Plus conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of Glyphomax Plus. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from Glyphomax Plus (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of Glyphomax Plus unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC
Roundup Ready® is a registered trademark of Monsanto Company

EPA-accepted 03/29/2001

6/1/01

308/2E
September 18, 2002



Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

GLYPHOMAX* PLUS (AI: GLYPHOSATE)
EPA REGISTRATION NUMBER: 62719-322
NOTIFICATION OF LABEL CHANGES REQUESTED BY CDPR PER PR NOTICE 98-10


Dow AgroSciences is respectfully submitting the following changes to Glyphomax Plus herbicide at the request of the California Department of Pesticide Regulation. The source label text is based on the EPA accepted copy dated March 29, 2001.

Propose Changes by Notification:

1. **Specific Crop Listings Canola, Crambe and Mustard on Page 21:** The term "Mustard" has been clarified to read "Mustard Seed" (Latest EPA accepted copy for Roundup® Ultra is referenced)
2. **Grain Sorghum (Hooded sprayers):** Added precautionary text from EPA accepted copy for Roundup® Ultra relating to use of hooded sprayers in tillering grain sorghum
3. **Soybeans with Roundup® Ready Gene:** Restored specific paragraphs to RR soybean section that provide spray volume information for aerial and ground applications from EPA accepted supplemental labeling for Glyphomax Plus dated August 31, 1999. (See also CDPR accepted copy for Roundup® Original dated July 3, 2002) Because of the potential for crop injury, Dow AgroSciences has chosen **not** to include the section from the supplemental label for Roundup® Original, allowing use of additional non-ionic surfactants, in the label for Glyphomax Plus.

Additional Typographical Changes:

1. Table on Page 23: Entry for Florida pusley at rates of 40 and 48 (stray hyphens before/after the weed height removed)
2. Pokeweed in Table on Page 29: Stray hyphen in front of the range "3-40" removed



70/71

GLYPHOMAX* PLUS (AI: GLYPHOSATE)
EPA REGISTRATION NUMBER: 62719-322
NOTIFICATION OF LABEL CHANGES PER PR NOTICE 98-10
September 18, 2002

Page 2

3. Storage and Disposal box: "in a landfill approved form pesticide disposal ..." corrected to read "for"
4. Controlling Droplet Size – Volume: "Nozzles with higher rated flows product larger droplets" (The word "product" corrected to read "produce".)
5. Hooded sprayers (Second Paragraph): "extend front and read flaps of the hoods... (The word "read" corrected read "rear".)
6. Grass Seed Production (last sentence): The word "Grow" corrected to "Grower"


This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Contents of Submission

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1 (OPP No. 281704) (2 Pages)
- Label titled Glyphomax* Plus (E8A / Glyphomax Plus / Notification With Edits / 09-16-02) (65 Pages plus Registration Notes) (5 Copies)
- Complimentary Copy: Letter from the California Department of Pesticide Regulation dated August 6, 2002

Please contact Richard Bjerregaard (rbjerregaard@dow.com), Label/Registration Specialist at 317/337-4676, or Paula McKinnies (pmckinnies@dow.com), Registration Assistant at 317/337-4679 if you require further information for this product.


Sincerely,



Steve A. McMaster
Regulatory Manager
Regulatory Success – Americas
PH: 317/337-4670 / FX: 317/337-4649
samcmaster@dow.com

Enclosures

/pkm





Department of Pesticide Regulation



Gray Davis
Governor

Winston H. Hickox
Secretary, California
Environmental
Protection Agency

Paul E. Helliker
Director

August 6, 2002

RECEIVED
SEP 09 2002
REGISTRATION

ID# 194809

Brian L. Bret, Ph.D.
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268-1054

COMPLIMENTARY
COPY

Dear Dr. Bret:

GLYPHOMAX PLUS, EPA Reg. No. 62719-322-ZA;
RETURN OF PROPOSED LABEL AMENDMENT TO ADD THE CANOLA,
CRAMBE, MUSTARD, DRY PEAS, LENTILS, CHICK PEAS, FLAX, ROUNDUP
READY CORN AND SOYBEANS, DURIAN, MANGOSTEEN, RAMBUTAN
CROPS, TO ADD COMMON POKEWEED AND PERENNIAL SOWTHISTLE
WEEDS, TO REVISE USE DIRECTIONS, TO REVISE PERSONAL PROTECTIVE
EQUIPMENT (PPE) STATEMENTS, AND TO REVISE STORAGE AND DISPOSAL
STATEMENTS

The following required items were not in your submission for label amendment:

1. Specific crop listings for Mustard on page 20.

Mustard implies too broad a term. Referenced Roundup Ultra Supplemental Label, EPA Reg. No. 524-475-ZB, lists Mustard Greens and Mustard (seed).

2. Complete use directions for Hooded Sprayers, page 29, under Grain Sorghum.

Your proposed label lacks use directions for tillering grain sorghum found on the referenced Roundup Ultra label.

3. Complete use directions for Soybeans with the Roundup Ready Gene. (See enclosed Supplemental Labeling for Roundup Original Herbicide, EPA Reg. No. 524-445-ZF)

4. Other requirements which may become evident upon evaluation of your completed submission.

Since your submission is incomplete, we are returning your request for label amendment. You may resubmit it when the necessary items can be provided.

FLEX YOUR POWER! The energy challenge facing California is real. Every Californian needs to take immediate action to reduce energy consumption. For a list of simple ways you can reduce demand and cut your energy costs, see our Web site at <www.cdpr.ca.gov>.

