

(Base Label):

00XXXXXX

[Insert 2-Point Black Line]

(logo) Dow AgroSciences LLC

# Glyphomax\*


For control of annual, perennial weeds and woody plants in various cropping systems, fallow cropland, CRP acres, and farmsteads, and other noncrop areas.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Active Ingredient(s):

Glyphosate † N-(phosphonomethyl)glycine, isopropylamine salt .....	41.0%
Inert Ingredients .....	59.0%
Total Ingredients .....	100.0%

† Contains 4 pounds per gallon glyphosate, isopropylamine salt  
Acid Equivalent: glyphosate acid – 30.4% (3 lb/gallon)



Keep Out of Reach of Children

## CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

Causes Eye Irritation • Harmful If Swallowed

Avoid contact with eyes or clothing.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**Do not mix, store or apply the product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to label booklet for Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-323

XX XX

EPA Est. \_\_\_\_\_

Superscripts correspond to places 7 & 8 of lot number

900-000000 / 00000000

\*Trademark of Dow AgroSciences LLC  
**Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.**  
[Insert 2-Point Black Line]

## Herbicide

Net Contents \_\_ gal

[Insert Bar Code FPO]  
[Insert DOT shipping classification and diamond(s)]

Lot

(Label Booklet):

(logo) Dow AgroSciences LLC

# Glyphomax\*

**For control of annual, perennial weeds and woody plants in various cropping systems, fallow cropland, CRP acres, and farmsteads, and other noncrop areas.**

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

Active Ingredient(s):

glyphosate† N-(phosphonomethyl)glycine, isopropylamine salt .....	41.0%
Inert Ingredients .....	59.0%
Total Ingredients .....	100.0%

† Contains 4 pounds per gallon glyphosate, isopropylamine salt  
Acid Equivalent: glyphosate acid – 30.4% (3 lb/gallon)

**Keep Out of Reach of Children**

## CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-323

XX XX  
EPA Est. \_\_\_\_\_

Superscripts correspond to places 7 & 8 of lot number  
900-000000 / 00000000

\*Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

# Herbicide

Net Contents \_\_ gal

<b>Contents</b>	<b>Page</b>
Precautionary Statements	-
Hazards To Humans and Domestic Animals	-
Environmental Hazards	-
Physical Or Chemical Hazards	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Mixing, Additives and Application Instructions	-
Mixing	-
Tank Mixtures	-
Additives	-
Surfactants	-
Ammonium Sulfate	-
Colorants and Dyes	-
Application Equipment and Techniques	-
Aerial Equipment	-
Broadcast Equipment	-
Controlled Droplet Application (CDA)	-
Hand-Held and High Volume Equipment	-
Selective Equipment	-
Shielded Applicators	-
Wiper Applicators	-
Weeds Controlled	-
Annual Weeds	-
Low-Volume Broadcast Application (Low Rate Technology)	-
Tank Mixtures	-
High-Volume Broadcast Applications	-
Perennial Weeds	-
Woody Brush and Trees	-
Noncrop Uses	-
Control of Emerged Weeds	-
Preemergence Weed Control	-
Farmstead Weed Control	-
Farm Ditches	-
Conservation Reserve Program (CRP Acres)	-
Dormant Rangeland	-
Habitat Management	-
Christmas Trees	-
Cut Stump Treatments	-
Grass Seed Production	-
Cropping Systems	-
Asparagus	-
Berries and Small Fruits	-
Corn	-
Fallow and Reduced Tillage Systems	-
Tank Mixtures	-
Ecofarming Systems	-
Aid to Tillage	-
Posharvest Grain Sorghum, Sorghum Regrowth Control	-
Pastures	-
Sugarcane	-

7/63

Conservation Tillage, Minimum Tillage and No-Till Systems	-
Corn and Soybeans (Tank Mixtures)	-
Preharvest Applications	-
Soybeans	-
Cotton	-
Grain Sorghum (Milo)	-
Wheat	-
Tree and Vine Crops	-
Middles Management (For Annual Weeds in Middles Between Rows of Tree and Vine Crops)	-
Strips (For Annual and Perennial Weeds in Strips of Tree and Vine Crops)	-
Tank Mixtures with Residual Herbicides	-
Perennial Grass Suppression Orchard Floors	-
Low Volume Application (Florida and Texas)	-
Tree Crops	-
Vine Crops	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

# CAUTION

**Causes Eye Irritation • Harmful If Swallowed**

Avoid contact with eyes or clothing.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.



**Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. *This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.*

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

### Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Disposal:** Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. Do not reuse this container. Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

---

**General Information**

---

Glyphomax\* herbicide is a water soluble liquid which mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

**Do not apply Glyphomax using aerial spray equipment except under conditions as specified within this label.**

The active ingredient in Glyphomax moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of Glyphomax and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of Glyphomax per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

Glyphomax does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of Glyphomax with herbicides or other materials that are not expressly recommended in this labeling. Mixing Glyphomax with herbicides or other materials not recommended on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

**Domestic Animals:** Glyphomax is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of Glyphomax or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Attention

**Avoid drift. Extreme care must be used when applying Glyphomax to prevent injury to desirable plants and crops.**

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of Glyphomax can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of Glyphomax is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **Avoid applying at excessive speed or pressure.**

**NOTE:** Use of Glyphomax in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

### Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information:**

**Importance of Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

#### Controlling Droplet Size:

**Volume-**Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure-**Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of nozzles**-Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation**-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

**Nozzle Type**-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

**Boom Length**-For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application**-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

## Mixing, Additives, and Application Instructions

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Do not apply when wind or other conditions favor drift. Hand-gun applications should be properly directed to avoid spraying desirable plants. NOTE: Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

### Mixing

Glyphomax mixes readily with water. Mix spray solutions of Glyphomax as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of Glyphomax (see the "Directions for Use" and "Weeds Controlled" sections of this label) near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

### Tank Mixtures

Always predetermine the compatibility of labeled tank mixtures of Glyphomax with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of Glyphomax with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it **slowly** through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture **slowly** through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of Glyphomax near the end of the filling process.
7. Where nonionic surfactant is recommended, add this to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

**Clean the entire sprayer after application of Glyphomax.**

Failure to clean the sprayer thoroughly may result in injury to desirable crops which are subsequently sprayed.

1. Add clean water to the tank and thoroughly rinse the entire sprayer system.
2. Fill the tank with water and ammonia. Add 1 quart of household ammonia per 25 gallons of water. Pump enough solution through the hoses, boom and nozzles to fill these parts completely.
3. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

Application or use of other agricultural chemicals with the equipment used for Glyphomax may result in injury to desirable vegetation.

---

**Additives**

---

**Surfactants**

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of Glyphomax when adding surfactant. When adding additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

**Ammonium Sulfate**

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Glyphomax, and Glyphomax plus 2,4-D, dicamba or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**NOTE:** The use of ammonium sulfate as an additive does not preclude the need for additional surfactant. Do not use herbicide rates lower than recommended in this label.

**Colorants or Dyes**

Agriculturally-approved colorants or marking dyes may be added to Glyphomax. Colorants or dyes used in spray solutions of Glyphomax may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

---

## Application Equipment and Techniques

---

**Chemigation:** Do not apply Glyphomax through any type of irrigation system.

Glyphomax may be applied with the following application equipment:

**Aerial:** Fixed Wing and Helicopter

### Broadcast Spray

**Controlled Droplet Applicator (CDA):** Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

**Hand-Held and High-Volume Spray Equipment:** Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, mistblowers<sup>1</sup>, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

<sup>1</sup> Glyphomax is not registered in California or Arizona for use in mistblowers.

**Selective equipment:** Recirculating sprayers, shielded sprayers and wiper applicators.

See the appropriate part of this section for specific instructions and rates of application.

## Aerial Equipment

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. Unless otherwise specified, do not exceed 1 quart per acre. **Aerial applications of Glyphomax may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems and preharvest applications. Refer to the individual use area sections of this label for recommended volumes and application rates.**

**AERIAL APPLICATION IN CALIFORNIA:** Refer to the Federal supplemental label for aerial applications in that state for specific instructions, restrictions and requirements.

Avoid direct application to any body of water.

**AVOID DRIFT. Do not apply during low-level inversion conditions, when winds are gusty or under any other condition which favors drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

**Ensure uniform application:** To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of *Glyphomax* accumulated during spraying or from spills. **Prolonged exposure of *Glyphomax* to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

*Glyphomax* plus tank mixtures of Banvel (dicamba) herbicide or 2,4-D tank may not be applied by air in California.

### Broadcast Equipment

**For control of annual or perennial weeds listed on this label using broadcast equipment:** Use the recommended rates of *Glyphomax* in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

### Controlled Droplet Application (CDA)

The rate of *Glyphomax* applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of *Glyphomax* at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of *Glyphomax* at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

### Hand-Held and High-Volume Equipment (Use coarse sprays only.)

Mix *Glyphomax* in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5 percent solution of *Glyphomax* plus nonionic surfactant to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing.

For annual weeds over 6 inches tall, or when not using additional surfactant, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.



17/63

When using application methods that result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

Prepare the desired volume of spray solution by mixing the amount of Glyphomax in water as shown in the following table:

**Spray Solution**

Desired Volume	Amount of Glyphomax					
	1/2%	1%	1 1/2%	2%	5%	10%
1 gal	2/3 fl oz	1 1/3 fl oz	2 fl oz	2 2/3 fl oz	6 1/2 fl oz	13 fl oz
25 gal	1 pt	1 qt	1 1/2 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1 1/2 gal	2 gal	5 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of Glyphomax be mixed with water in a larger container. Fill sprayer with the mixed solution.

**Selective Equipment**

Glyphomax may be applied through a recirculating spray system, a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

**NOTE: Avoid contact with desirable vegetation.**

Contact of the herbicide solution with the desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

**Shielded Applicators**

When applied as directed under conditions described for shielded applicators, Glyphomax will control those weeds listed in the "Weeds Controlled" section of this label.

Use the following equation to convert from a broadcast rate per acre to a band rate per acre.

$$\begin{array}{l} \text{Band width} \\ \text{in inches} \end{array} \times \begin{array}{l} \text{Herbicide} \\ \text{Broadcast} \\ \text{RATE} \\ \text{per acre} \end{array} = \begin{array}{l} \text{Herbicide} \\ \text{Band} \\ \text{RATE} \\ \text{per acre} \end{array}$$

$$\begin{array}{l} \text{Band width} \\ \text{in inches} \end{array} \times \begin{array}{l} \text{Broadcast} \\ \text{VOLUME of} \\ \text{solution} \\ \text{per acre} \end{array} = \begin{array}{l} \text{Band} \\ \text{VOLUME} \\ \text{of solution} \\ \text{per acre} \end{array}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. **Extreme care must be exercised to avoid contact with desirable vegetation.**

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

### Wiper Applicators

Wiper applicators are devices that physically wipe appropriate amounts of Glyphomax directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using Glyphomax by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

**For Rope or Sponge Wick Applicators:** Mix 1 gallon of Glyphomax in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this "Wiper Applicators" section.

**For Porous-Plastic Applicators:** Solutions ranging from 33 to 100 percent of Glyphomax in water may be used in porous-plastic wiper applicators.

19/63

The following weeds are CONTROLLED when Glyphomax is applied as recommended under the conditions described for " Wiper Applicators ":

<u>Common Name</u>	<u>Scientific Name</u>
<b>(Annual Grasses)</b>	
Corn	<i>Zea mays</i>
Panicum, Texas	<i>Panicum texanum</i>
Rye, common	<i>Secale cereale</i>
Shattercane	<i>Sorghum bicolor</i>
<b>(Annual Broadleaves)</b>	
Sicklepod	<i>Cassia obtusifolia</i>
Spanishneedles	<i>Bidens bipinnata</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>

The following weeds are SUPPRESSED when Glyphomax is applied as recommended under the conditions described for " Wiper Applicators ":

<u>Common Name</u>	<u>Scientific Name</u>
<b>(Annual Broadleaves)</b>	
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Dogfennel	<i>Eupatorium capilliflorum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Sunflower	<i>Helianthus annuus</i>
Thistle, musk	<i>Carduus nutans</i>
Velvetleaf	<i>Abutilon theophrasti</i>
<b>(Perennial Grasses)</b>	
Bermudagrass	<i>Cynodon dactylon</i>
Guineagrass	<i>Panicum maximum</i>
Johnsongrass	<i>Sorghum halepense</i>
Smutgrass	<i>Sporobolus poiretii</i>
Vaseygrass	<i>Paspalum urvillei</i>
<b>(Perennial Broadleaves)</b>	
Dogbane, hemp	<i>Apocynum cannabinum</i>
Milkweed	<i>Asclepias syriaca</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Thistle, Canada	<i>Cirsium arvense</i>

---

**Weeds Controlled**

---

This herbicide controls many annual and perennial grasses and broadleaf weeds.

---

**Annual Weeds**

---

Apply to actively growing grass and broadleaf weeds.

Allow at least 3 days after treatment before tillage.

For maximum agronomic benefit, apply when weeds are 6 inches or less in height.

To prevent seed production, applications should be made prior to seedhead formation.

Glyphomax does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

**Low-Volume Broadcast Application  
(Low Rate Technology)**

When applied as directed under the conditions described, Glyphomax will control the weeds listed below when:

1. Water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are recommended. (See the "Aerial Application" section of this label for approved sites.)
2. A nonionic surfactant is added at 0.5 to 1 percent by total spray volume. Use 0.5 percent surfactant concentration when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration for those surfactants containing less than 70 percent active ingredient.

**NOTE:**

The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of Glyphomax on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.

Do not tank-mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the "**Tank Mixtures**" portion of this section for control of additional broadleaf weeds.

21/63

Weed Species	Maximum Height/Length	Rate per Acre <sup>1</sup> (fl oz)
<b>Chickweed, jagged</b> <i>Holosteum umbellatum</i>	12"	8 oz
<b>Foxtail</b> <i>Setaria spp</i>		
<b>Barnyardgrass</b> <sup>2</sup> <i>Echinochloa crus-galli</i>	6" 0 to 4"	12 oz 16 oz <sup>2</sup>
<b>Bluegrass, annual</b> <i>Poa annua</i>	4 to 6"	24 oz <sup>2</sup>
<b>Brome, downy</b> <sup>3</sup> <i>Bromus tectorum</i>		
<b>Mustard, blue</b> <i>Chorispora tenella</i>		
<b>Mustard, tansy</b> <i>Descurainia pinnata</i>		
<b>Mustard, tumble</b> <i>Sisymbrium altissimum</i>		
<b>Mustard, wild</b> <i>Sinapis arvensis</i>		
<b>Spurry, umbrella</b> <i>Holosteum umbellatum</i>		
<b>Barley</b> <i>Hordeum vulgare</i>	12"	
<b>Rye</b> <i>Secale cereale</i>		
<b>Sandbur, field</b> <i>Cenchrus spp.</i>		
<b>Shattercane</b> <i>Sorghum bicolor</i>		
<b>Stinkgrass</b> <i>Eragrostis cilianensis</i>		
<b>Wheat</b> <i>Triticum aestivum</i>	18"	12 oz
<b>Morningglory</b> <i>Ipomoea spp.</i>	2"	16 oz.
<b>Sicklepod</b> <i>Cassia obtusifolia</i>		
<b>Bluegrass, bulbous</b> <i>Poa bulbosa</i>	6"	
<b>Cheat</b> <i>Bromus secalinus</i>		
<b>Chickweed, common</b> <i>Stellaria media</i>		
<b>Chickweed, mouseear</b> <i>Cerastium vulgatum</i>		
<b>Corn</b> <i>Zea mays</i>		
<b>Goatgrass, jointed</b> <i>Aegilops cylindrica</i>		
<b>Groundsel, common</b> <i>Senecio vulgaris</i>		

<b>Henbit</b> <i>Lamium amplexicaule</i> <b>Horseweed/Marestail</b> <i>Conyza canadensis</i> <b>Lambsquarters, common</b> <i>Chenopodium album</i> <b>Pennycress, field (Fanweed)</b> <i>Thlaspi arvense</i> <b>Rocket, London</b> <i>Sisymbrium irio</i> <b>Ryegrass, Italian</b> <i>Lolium multiflorum</i> <b>Shepherdspurse</b> <i>Capsella bursa-pastoris</i> <b>Spurge, annual</b> <i>Euphorbia spp.</i>		
<b>Buttercup</b> <i>Ranunculus spp.</i> <b>Cocklebur</b> <i>Xanthium strumarium</i> <b>Crabgrass</b> <i>Digitaria spp.</i> <b>Dwarfandelion</b> <i>Krigia cespitosa</i> <b>Falseflax, smallseed</b> <i>Camelina microcarpa</i> <b>Foxtail, Carolina</b> <i>Alopecurus carolinianus</i> <b>Johnsongrass, seedling</b> <i>Sorghum halepense</i> <b>Oats, wild</b> <i>Avena fatua</i> <b>Panicum, fall</b> <i>Panicum dichotomiflorum</i> <b>Panicum, Texas</b> <i>Panicum texanum</i> <b>Pigweed, redroot</b> <i>Amaranthus retroflexus</i> <b>Pigweed, smooth</b> <i>Amaranthus hybridus</i> <b>Witchgrass</b> <i>Panicum capillare</i>	12"	
<b>Sicklepod</b> <i>Cassia obtusifolia</i>	3 to 4"	24 oz
<b>Signalgrass, broadleaf</b> <i>Brachiaria platyphylla</i>	4"	
<b>Horseweed/Marestail</b> <i>Conyza canadensis</i> <b>Lambsquarters, common</b> <i>Chenopodium album</i> <b>Spurge, annual</b> <i>Euphorbia spp.</i>	7 to 12"	
<b>Rice, red</b>	4"	32 oz

<i>Oryza sativa</i> <b>Teaweed</b> <i>Sida spinosa</i>		
<b>Sprangletop</b> <i>Leptochloa spp.</i>	6"	
<b>Geranium, Carolina</b> <i>Geranium carolinianum</i> <b>Goosegrass</b> <i>Eleusine indica</i> <b>Primrose, cutleaf evening</b> <i>Oenothera laciniata</i> <b>Pusley, Florida</b> <i>Richardia scabra</i>	12"	
<b>Sicklepod</b> <i>Cassia obtusifolia</i> <b>Spanishneedles</b> <i>Bidens bipinnata</i>	5 to 12"	
<b>Filaree</b> <i>Erodium spp.</i> <b>Sprangletop</b> <i>Leptochloa spp.</i>	12"	48 oz

<sup>1</sup> For those rates less than 32 fluid ounces per acre, Glyphomax at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

<sup>2</sup> Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana and Texas for preplant treatments.

<sup>3</sup> For control in no-till systems, use 16 fluid ounces per acre.

### Tank Mixtures

- Glyphomax plus Banvel (dicamba) plus Nonionic Surfactant
- Glyphomax plus 2,4-D plus Nonionic Surfactant

Do not apply Banvel (dicamba) herbicide or 2,4-D herbicide tank mixtures by air in California.

These tank mixtures are recommended for use in fallow and reduced tillage areas only. Follow use directions as given in the "Low-Volume Broadcast Application" section.

Glyphomax plus dicamba or 2,4-D will control the annual grasses and broadleaf weeds listed for Glyphomax alone at the indicated heights (except 8 fluid ounces per acre applications), plus the following broadleaf weeds. For those weeds previously listed at 8 fluid ounces of Glyphomax alone per acre, use 12 fluid ounces in these tank mixtures.

**NOTE:** Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting. The addition of dicamba in a mixture with Glyphomax may provide short-term residual control of selected weed species.

Apply 12 to 16 fluid ounces of Glyphomax plus 0.25 pound a.i. of dicamba or 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control dense populations of the following annual broadleaf weeds (Treat when weeds are less than the height indicated.):

Common Name	Scientific Name	Height
Cocklebur	<i>Xanthium strumarium</i>	12"
Kochia <sup>1</sup>	<i>Kochia scoparia</i>	6"
Lambsquarters	<i>Chenopodium album</i>	12"
Lettuce, prickly	<i>Lactuca serriola</i>	6"
Marestail/Horseweed	<i>Conyza canadensis</i>	6"
Morningglory	<i>Ipomoea spp.</i>	6"
Pigweed, redroot	<i>Amaranthus retroflexus</i>	12"
Pigweed, smooth	<i>Amaranthus hybridus</i>	12"
Thistle, Russian	<i>Salsola kali</i>	12"

<sup>1</sup> Controlled with dicamba tank mixture only.

Apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control the following annual broadleaf weeds (Treat when weeds are less than 6 inches in height):

Common Name	Scientific Name
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Velvetleaf	<i>Abutilon theophrasti</i>

### High-Volume Broadcast Applications

When applied as directed under the conditions described, Glyphomax will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Apply 1 to 1.5 quarts of Glyphomax per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume. Use 1 quart per acre if weeds are less than 6 inches tall and 1.5 quarts per acre if weeds are over 6 inches tall. If weeds have been mowed, grazed or cut, allow adequate time for new growth to reach recommended stages prior to treatment. These rates will also provide control of weeds listed in the "Low-Volume Broadcast Application" section.

### Annual Weed Species

Common Name	Scientific Name
Balsamapple <sup>1</sup>	<i>Momordica charantia</i>
Bassia, fivehook	<i>Bassia hyssopifolia</i>
Brome	<i>Bromus spp.</i>
Fiddleneck	<i>Amsinckia spp.</i>
Fleabane, hairy	<i>Conyza bonariensis</i>
Fleabane	<i>Erigeron spp.</i>
Kochia	<i>Kochia scoparia</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Panicum	<i>Panicum spp.</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>



Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sowthistle, annual	<i>Sonchus cleraceus</i>
Sunflower	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>

<sup>1</sup> Apply with hand-held equipment only.

### Perennial Weeds

Apply Glyphomax as follows to control or destroy most perennial weeds:

**NOTE:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Glyphomax on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.

**When applied as recommended under the conditions described, Glyphomax WILL CONTROL the following perennial weeds:**

Common Name	Scientific Name
Alfalfa	<i>Medicago sativa</i>
Alligatorweed <sup>1</sup>	<i>Alternanthera philoxeroides</i>
Anise (fennel)	<i>Foeniculum vulgare</i>
Artichoke, Jerusalem	<i>Helianthus tuberosus</i>
Bahiagrass	<i>Paspalum notatum</i>
Bentgrass	<i>Agrostis spp.</i>
Bermudagrass	<i>Cynodon dactylon</i>
Bermudagrass, water (knotgrass)	<i>Paspalum distichum</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Bluegrass, Kentucky	<i>Poa spp.</i>
Blueweed, Texas	<i>Helianthus ciliaris</i>
Brackenfern	<i>Pteridium aquilinum</i>
Bromegrass, smooth	<i>Bromus inermis</i>
Bursage, woollyleaf	<i>Franseria tomentosa</i>
Canarygrass, reed	<i>Phalaris arundinacea</i>
Cattail	<i>Typha spp.</i>
Clover, red	<i>Trifolium pratense</i>
Clover, white	<i>Trifolium repens</i>
Cogongrass	<i>Imperata cylindrica</i>
Dallisgrass	<i>Paspalum dilatatum</i>
Dandelion	<i>Taraxacum officinale</i>
Dock, curly	<i>Rumex crispus</i>
Dogbane, hemp	<i>Apocynum cannabinum</i>
Fescues	<i>Festuca spp.</i>

Fescue, tall	<i>Festuca arundinacea</i>
Guineagrass	<i>Panicum maximum</i>
Horsenettle	<i>Solanum carolinense</i>
Horseradish	<i>Armoracia rusticana</i>
Ice plant	<i>Mesembryanthemum crystallinum</i>
Johnsongrass	<i>Sorghum halepense</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
Knapweed	<i>Centaurea repens</i>
Lantana	<i>Lantana camara</i>
Lespedeza	<i>Lespedeza spp.</i>
Milkweed	<i>Asclepias spp.</i>
Muhly, wirestem	<i>Muhlenbergia frondosa</i>
Mullein, common	<i>Verbascum thapsus</i>
Najagrass	<i>Pennisetum purpureum</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Orchardgrass	<i>Dactylis glomerata</i>
Pampasgrass	<i>Cortaderia spp.</i>
Paragrass	<i>Brachiaria mutica</i>
Phragmites <sup>1</sup>	<i>Phragmites spp.</i>
Poison hemlock	<i>Conium maculatum</i>
Quackgrass	<i>Agropyron repens</i>
Redvine <sup>1</sup>	<i>Brunnichia ovata</i>
Reed, giant	<i>Arundo donax</i>
Ryegrass, perennial	<i>Lolium perenne</i>
Smartweed, swamp	<i>Polygonum coccineum</i>
Spurge, leafy <sup>1</sup>	<i>Euphorbia esula</i>
Starthistle, yellow	<i>Centaurea solstitialis</i>
Sweet potato, wild <sup>1</sup>	<i>Ipomoea pandurata</i>
Thistle, Canada	<i>Cirsium arvense</i>
Thistle, artichoke	<i>Cynara cardunculus</i>
Timothy	<i>Phleum pratense</i>
Torpedograss <sup>1</sup>	<i>Panicum repens</i>
Trumpetcreeper <sup>1</sup>	<i>Campsis radicans</i>
Vaseygrass	<i>Paspalum urvillei</i>
Velvetgrass	<i>Holcus spp.</i>
Wheatgrass, western	<i>Agropyron smithii</i>

<sup>1</sup> Partial control.

Glyphomax is not registered in California for use on water bermudagrass.

See "Directions for Use" and "Mixing, Additives and Application Instructions" sections of this label for labeled uses and specific application instructions.

**Alfalfa:** Apply 1 quart of Glyphomax per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

**Alligatorweed:** Apply 4 quarts of Glyphomax per acre or apply a 1.5 percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

**Anise (fennel)/Poison hemlock:** Apply a 1 to 2 percent solution of Glyphomax as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.

**Bentgrass:** For suppression in grass seed production areas. For ground applications only, apply 1.5 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should be actively growing and have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.

**Bermudagrass:** For control, apply 5 quarts of Glyphomax per acre. For partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. Allow 7 or more days after application before tillage.

**Bermudagrass, water (knotgrass):** Apply 1.5 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Apply when water bermudagrass is actively growing and 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water bermudagrass that is actively growing and 12 to 18 inches in length. Allow 7 or more days before tillage.

**Bindweed, field:** For control, apply 4 to 5 quarts of Glyphomax per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

Also for control, apply 2 quarts of Glyphomax plus 0.5 pound a.i. of dicamba in 10 to 20 gallons of water per acre. At these rates, apply using ground application only.

The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of Glyphomax plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

**In California only,** apply 1 to 5 quarts of Glyphomax per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions.

For suppression on irrigated land where annual tillage is performed, apply 1 quart of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. Allow 3 or more days after application before tillage.

**Bluegrass, Kentucky / Bromegrass, smooth / Orchardgrass:** Apply 2 quarts of Glyphomax in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.

Orchardgrass (sods going to no-till corn): Apply 1 to 1.5 quarts of Glyphomax per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine herbicide will be necessary for optimum results.

**Blueweed, Texas:** Apply 4 to 5 quarts of Glyphomax per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

**Brackenfern:** Apply 3 to 4 quarts of Glyphomax per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

**Bursage, woollyleaf:** For control, apply 2 quarts of Glyphomax plus 0.5 lb a.i. of dicamba per acre. For partial control, apply 1 quart of Glyphomax plus 0.5 lb a.i. of dicamba per acre. Add 0.5 to 1 percent nonionic surfactant by total spray volume and apply in 3 to 20 gallons of water per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

**Canarygrass, reed/Timothy / Wheatgrass, western:** Apply 2 to 3 quarts of Glyphomax per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.

**Cogongrass:** Apply 3 to 5 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant in 10 to 40 gallons of water per acre. Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

**Dandelion/Dock, curly:** Apply 3 to 5 quarts of Glyphomax per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

**Dogbane, hemp:** Apply 4 quarts of Glyphomax per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

29/63

For suppression, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

**Fescue, tall:** Apply 3 quarts of Glyphomax in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.

Fall applications only, apply 1 quart of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of Glyphomax plus nonionic surfactant will improve long-term control and control seedlings germinating after fall treatments or the following spring.

**Guineagrass:** Apply 3 quarts of Glyphomax per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.

**Johnsongrass/Ryegrass, perennial:** Apply 1 to 3 quarts of Glyphomax per acre. In annual cropping systems apply 1 to 2 quarts of Glyphomax per acre. Apply 1 quart of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of Glyphomax when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of Glyphomax in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart per acre rate.

For burndown of Johnsongrass, apply 1 pint per acre plus 0.5 to 1 percent nonionic surfactant in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression): Apply a 1 percent solution of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

**Kikuyugrass:** Apply 2 to 3 quarts of Glyphomax per acre. Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

**Knapweed/Horseradish:** Apply 4 quarts of Glyphomax per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

**Lantana:** Apply Glyphomax as a 1 to 1.25 percent solution using hand-held equipment only. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.

**Milkweed, common:** Apply 3 quarts of Glyphomax per acre. Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.

**Muhly, wirestem:** Apply 1 to 2 quarts of Glyphomax per acre. Use 1 quart of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of Glyphomax when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. Glyphomax will not provide residual control of wirestem muhly from seeds which germinate after application of Glyphomax. Do not tank mix with residual herbicides when using the 1 quart per acre rate.

**Nightshade, silverleaf:** For control, apply 2 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth.

**Nutsedge; purple, yellow:** Apply 3 quarts of Glyphomax per acre as a broadcast spray, or apply a 1 to 2 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.

Sequential applications of 1 to 2 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre will provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For suppression to partial control of existing plants, apply 1 pint to 2 quarts of Glyphomax per acre, plus 0.5 to 1 percent nonionic surfactant in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.

**Pampasgrass/Ice plant:** Apply Glyphomax as a 1.5 to 2 percent solution using hand-held equipment. Apply to plants that are actively growing. Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

**Phragmites:** For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

**Quackgrass In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage:** Apply 1 to 2 quarts of Glyphomax per acre. For the 1 quart rate, apply 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 6 to 8 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results use a moldboard plow.

**Quackgrass In Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned**

**Following Application:** Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.

**Redvine:** For suppression, apply 24 fluid ounces of Glyphomax per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre plus 0.5 to 1 percent nonionic surfactant by total volume. Apply in late September or early October to actively growing plants, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

**Reed, giant:** For control of giant reed, apply a 2 percent solution of Glyphomax when plants are actively growing. Best results are obtained when applications are made in late summer to fall.

**Smartweed, swamp:** Apply 3 to 5 quarts of Glyphomax per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

**Spurge, leafy:** For suppression, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

**Starthistle, yellow:** Best results are obtained when applications are made during periods of active growth, including the rosette, bolting and early flower stages. For spray-to-wet applications, apply Glyphomax as a 2 percent solution. For broadcast applications, apply 2 quarts per acre in 10 to 40 gallons per acre of water carrier.

**Sweet Potato, wild/Thistle, artichoke:** Apply Glyphomax as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications may be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.

**Thistle, Canada:** Apply 2 to 3 quarts of Glyphomax per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Glyphomax. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression of Canada thistle, apply 1 quart per acre of Glyphomax, or 1 pint of Glyphomax plus 0.5 pound a.i. 2,4-D per acre, plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

**Torpedograss:** Apply 4 to 5 quarts of Glyphomax per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.

**Trumpet creeper:** For control, apply 2 quarts of Glyphomax per acre in 5 to 10 gallons of water per acre. Apply to actively growing plants in late September or October, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

**Other perennials listed on this label:** Apply 3 to 5 quarts of Glyphomax per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

### Woody Brush and Trees

When applied as recommended under the conditions described, Glyphomax controls or partially controls the following woody brush, plants and trees:

Scientific Name	Common Name
Alder	<i>Alnus spp.</i>
Ash <sup>1</sup>	<i>Fraxinus spp.</i>
Aspen, quaking	<i>Populus tremuloides</i>
Bearmat (Bearclover)	<i>Chamaebatia foliolosa</i>
Beech	<i>Fagus grandifolia</i>
Birch	<i>Betula spp.</i>
Blackberry	<i>Rubus spp.</i>
Blackgum	<i>Nyssa spp.</i>
Bracken	<i>Peridium spp.</i>
Broom, French	<i>Cytisus monspessulanus</i>
Broom, Scotch	<i>Cytisus scoparius</i>
Buckwheat, California <sup>1</sup>	<i>Eriogonum fasciculatum</i>
Cascara <sup>1</sup>	<i>Rhamnus purshiana</i>
Catsclaw	<i>Acacia greggi</i>
Ceanothus <sup>1</sup>	<i>Ceanothus spp.</i>
Chamise	<i>Adenostoma fasciculatum</i>
Cherry, Bitter	<i>Prunus emarginata</i>
Cherry, black	<i>Prunus serotina</i>
Cherry, pin	<i>Prunus pensylvanica</i>
Coyote brush	<i>Baccharis consanguinea</i>
Creeper, Virginia <sup>1</sup>	<i>Parthenocissus quinquefolia</i>
Dewberry	<i>Rubus trivialis</i>
Dogwood	<i>Cornus spp.</i>
Elderberry	<i>Sambucus spp.</i>
Elm <sup>1</sup>	<i>Ulmus spp.</i>
Eucalyptus	<i>Eucalyptus spp.</i>
Gorse	<i>Ulex europaeus</i>
Hasardia	<i>Haplopappus squamosus</i>
Hawthorn	<i>Crataegus spp.</i>
Hazel	<i>Corylus spp.</i>
Hickory <sup>1</sup>	<i>Carya spp.</i>
Holly, Florida	<i>Schinus terebinthifolius</i>



(Brazilian Peppertree <sup>1</sup>)

Honeysuckle	<i>Lonicera spp.</i>
Hornbeam, American <sup>1</sup>	<i>Carpinus caroliniana</i>
Kudzu	<i>Pueraria lobata</i>
Locust, black <sup>1</sup>	<i>Robinia pseudoacacia</i>
Madrone	<i>Arbutus menziesii</i>
Manzanita	<i>Arctostaphylos spp.</i>
Maple, red <sup>2</sup>	<i>Acer rubrum</i>
Maple, sugar	<i>Acer saccharum</i>
Maple, vine <sup>1</sup>	<i>Acer circinatum</i>
Monkey Flower <sup>1</sup>	<i>Mimulus guttatus</i>
Oak, Black <sup>1</sup>	<i>Quercus velutina</i>
Oak, northern pin	<i>Quercus palustris</i>
Oak, post	<i>Quercus stellata</i>
Oak, red	<i>Quercus rubra</i>
Oak, southern red	<i>Quercus falcata</i>
Oak, white <sup>1</sup>	<i>Quercus alba</i>
Persimmon <sup>1</sup>	<i>Diospyros spp.</i>
Pine	<i>Pinus spp.</i>
Poison-ivy	<i>Rhus radicans</i>
Poison-oak	<i>Rhus toxicodendron</i>
Poplar, yellow <sup>1</sup>	<i>Liriodendron tulipifera</i>
Raspberry	<i>Rubus spp.</i>
Redbud, eastern	<i>Cercis canadensis</i>
Rose, multiflora	<i>Rosa multiflora</i>
Russian-olive <sup>3</sup>	<i>Elaeagnus angustifolia</i>
Sage; black, white	<i>Salvia spp.</i>
Sagebrush, California	<i>Artemisia californica</i>
Salmonberry	<i>Rubus spectabilis</i>
Salt cedar	<i>Tamarix spp.</i>
Sassafras	<i>Sassafras albidum</i>
Sourwood	<i>Oxydendrum arboreum</i>
Sumac, poison <sup>1</sup>	<i>Rhus vernix</i>
Sumac, smooth <sup>1</sup>	<i>Rhus glabra</i>
Sumac, winged <sup>1</sup>	<i>Rhus copallina</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Swordfern <sup>1</sup>	<i>Polystichum munitum</i>
Tallowtree, Chinese	<i>Sapium sebiferum</i>
Tan Oak	<i>Lithocarpus densiflorus</i>
Thimbleberry	<i>Rubus parviflorus</i>
Tobacco, tree	<i>Nicotiana glauca</i>
Trumpetcreeper	<i>Campsis radicans</i>
Waxmyrtle, southern <sup>1</sup>	<i>Myrica cerifera</i>
Willow	<i>Salix spp.</i>

<sup>1</sup> Partial control.

<sup>2</sup> See below for control or partial control instructions.

<sup>3</sup> Glyphomax is not registered in California for use on Russian-olive.

**NOTE:** If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth.

Apply Glyphomax when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See "Directions for Use" and "Mixing, Additives and Application Instructions" sections of this label for labeled uses and specific application instructions.

Apply Glyphomax as follows to control or partially control the following woody brush and trees.

**Alder/Dewberry/Honeysuckle/Post Oak/Raspberry:** For control, apply 3 to 4 quarts per acre of Glyphomax as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

**Aspen, quaking/Cherry, bitter, black, pin/Hawthorn /Oak, southern red/Sweetgum/Trumpet creeper:** For control, apply 2 to 3 quarts of Glyphomax per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

**Birch/ Elderberry/ Hazel/ Salmonberry/ Thimbleberry:** For control, apply 2 quarts per acre of Glyphomax as a broadcast spray or as a 1 percent solution with hand-held equipment.

**Blackberry:** For control, apply 3 to 4 quarts per acre of Glyphomax as a broadcast spray, or 1 to 1.5 percent solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4 percent solution of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of Glyphomax in 10 to 40 gallons of water per acre.

**Broom: French, Scotch:** For control, apply a 1.5 to 2 percent solution with hand-held equipment.

**Buckwheat, California/Hasardia/Monkey Flower/ Tobacco, tree:** For partial control of these species, apply a 1 to 2 percent solution of Glyphomax as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

**Catsclaw:** For partial control, apply as a 1 to 1.5 percent solution with hand-held equipment.

**Coyote Brush:** For control, apply a 1.5 to 2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

**Eucalyptus:** For control of eucalyptus resprouts, apply a 2 percent solution of Glyphomax with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.

**Kudzu:** For control, apply 4 quarts of Glyphomax per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

**Madrone resprouts:** For suppression or partial control, apply a 2 percent solution of Glyphomax to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

**Maple, red:** For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of Glyphomax per acre as a broadcast spray.

**Maple, sugar/Oak, northern pin/Oak, red:** For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

**Poison-ivy/Poison-oak:** For control, apply 4 to 5 quarts of Glyphomax per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

**Rose, multiflora:** For control, apply 2 quarts of Glyphomax per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

**Sage, black/Sagebrush, California/Chamise/Tallowtree, Chinese:** For control of these species, apply a 1 percent solution of Glyphomax as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

**Tan oak resprouts:** For suppression or partial control, apply a 2 percent solution of Glyphomax to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.

**Willow:** For control, apply 3 quarts of Glyphomax per acre as a broadcast spray or as a 1 percent solution with hand-held equipment.

**Other Woody Brush and Trees listed on this label:** For partial control, apply 2 to 5 quarts of Glyphomax per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment.

---

## Noncrop Uses

---

See "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information and the following "Noncrop" sections for specific recommended uses.

**NOTE: Extreme care must be exercised to avoid contact of spray with foliage, green stems, exposed non-woody roots or fruit of crops, desirable turfgrasses, trees, shrubs or other desirable vegetation since severe damage or destruction may result.**

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Where repeat applications are necessary, do not exceed 10.6 quarts of Glyphomax per acre per year.

Glyphomax does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Glyphomax controls annual and perennial weeds listed on this label growing in areas such as ditch banks, dry ditches, dry canals, fencerows, and noncrop areas.

For specific rates of application and instructions for control of various annual and perennial weeds and woody brush and trees, see the "Weeds Controlled" section of this label.

Glyphomax may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any noncrop site specified on this label. See the "Selective Equipment" part of "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.

### Control of Emerged Weeds

**NOTE:** For backpack sprayer and handgun applications, see the "Hand-Held and High-Volume Equipment" section for recommended rates.

**Annual Weeds:** Apply 1 quart per acre of Glyphomax in these tank mixtures when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are more than 6 inches tall.

**Perennial Weeds:** For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of Glyphomax. Follow the recommendations in the "Weeds Controlled" section of this label for stage of growth and rate of application for specific perennial weeds.

### Preemergence Weed Control

For preemergence weed control, refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements.

Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

### Farmstead Weed Control

When applied as directed for "Noncrop Uses", under conditions described, Glyphomax controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts and for general nonselective farmstead weed control.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

**NOTE:** If spraying areas adjacent to desirable plants, use a shield made of cardboard, sheet metal or plywood while spraying to help prevent spray from contacting foliage of desirable plants.

### Farm Ditches

Glyphomax will suppress perennial grasses along farm ditches. Apply Glyphomax at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces per acre without ammonium sulfate when treating Kentucky bluegrass.

Apply treatments in 10 to 20 gallons of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles.

Add a nonionic surfactant at a rate of 0.5 percent of the spray solution.

Where broadleaf weed control or suppression is desired, tank mix Glyphomax with an appropriate, labeled broadleaf weed herbicide.

---

### **Conservation Reserve Program (CRP Acres)**

---

Glyphomax can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres.

For specific rates of application for various annual and perennial weeds, see the "Weeds Controlled" section of this label.

CRP applications may be made with wiper applicators or conventional spray equipment.

For selective applications with broadcast spray equipment, apply 12 to 16 ounces per acre of Glyphomax in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

---

### **Dormant Rangeland**

---

Glyphomax will control or suppress many weeds, including downy brome, cheat grass, cereal rye, medusahead rye and jointed goatgrass in dormant rangeland.

Apply 8 to 16 ounces per acre of Glyphomax in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass are still truly dormant.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of Glyphomax wear off.

Do not use additional surfactant or ammonium sulfate when spraying dormant rangeland grasses with Glyphomax.

---

### **Habitat Management**

---

Glyphomax is recommended for the restoration and/or maintenance of native habitats and in wildlife management areas. Apply as recommended in the "Noncrop Uses" section of this label.

**Habitat Restoration and Maintenance:** When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to *selectively remove unwanted plants for habitat maintenance and enhancement*. For spot treatments, care should be exercised to keep spray off of desirable plants.

**Wildlife Food Plots:** Glyphomax may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying Glyphomax, or native species may be allowed to *repopulate the area*. If tillage is needed to prepare a seedbed, wait 7 days after applying Glyphomax before tilling.

---

### Christmas Trees

---

**NOTE:** Glyphomax is not recommended for use as an over-the-top broadcast spray in Christmas trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

When applied as instructed for the conditions described for "Noncrop Uses", Glyphomax controls undesirable vegetation listed on this label prior to planting, and as a post directed spray around Christmas trees.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

Where repeat applications are necessary, do not exceed 10.6 quarts of Glyphomax per acre per year.

**Site Preparation:** Following preplant applications of Glyphomax, any Christmas tree species may be planted. Precautions should be taken to protect nontarget plants during site preparation applications.

**Post Directed Spray:** Use as a post directed spray around Christmas trees. Care must be exercised to avoid contact of spray, drift or mist with foliage of or green bark of established species.

### Cut Stump Treatments

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with Glyphomax. Apply Glyphomax using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of Glyphomax to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, Glyphomax will control, partially control or suppress many types of woody brush and tree species, some of which are listed below:

Common Name	Scientific Name
Alder	<i>Alnus spp.</i>
Eucalyptus	<i>Eucalyptus spp.</i>
Madrone	<i>Arbutus menziesii</i>
Oak	<i>Quercus spp.</i>
Reed, giant	<i>Arundo donax</i>
Saltcedar	<i>Tamarisk spp.</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Tan Oak	<i>Lithocarpus densiflorus</i>
Willow	<i>Salix spp.</i>

---

### Grass Seed Production

---

#### Preplant and Renovation

When applied as directed for "Noncrop Uses", under conditions described, Glyphomax controls most existing vegetation prior to the planting or renovation of grass seed production areas.

For specific rates of application and instructions for control of various annual and perennial weeds, and woody brush and trees, see the "Weeds Controlled" section of this label.

For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control.

**Do not disturb soil or underground plant parts before treatment.** Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

Apply Glyphomax to actively growing weeds at the stages of growth recommended in the "Weeds Controlled" section of this label prior to planting or renovation of turf or forage grass areas grown for seed production.

Do not feed or graze treated areas within 8 weeks after application.

---

### Cropping Systems

---

When applied as directed for "Cropping Systems", under the conditions described, Glyphomax controls annual and perennial weeds listed on this label, prior to the emergence of direct seeded crops or prior to transplanting of crops listed on this label.

See "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information.

See the following "Cropping Systems" sections for specific recommended uses.

**Extreme care must be exercised to avoid contact of spray with foliage, green stems or fruit of desirable crops, plants, trees or other desirable vegetation since severe damage or destruction may result.**

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts per acre of Glyphomax per year.

For any crop not listed below, applications must be made at least 30 days prior to planting.

Do not harvest or feed treated vegetation for 8 weeks following application. Following spot treatment or selective equipment use, allow 14 days before grazing domestic livestock or harvesting forage grasses and legumes.

### Row Crops

Corn (All) <sup>1</sup>	Sorghum (Milo) <sup>1</sup>
Cotton <sup>1</sup>	Soybeans <sup>1</sup>
Peanuts	Sugarcane <sup>1</sup>

### Cereal Grains

Barley <sup>1</sup>	Rye <sup>1</sup>
Buckwheat <sup>1</sup>	Triticale <sup>1</sup>
Millet (Pearl, Proso) <sup>1</sup>	Wheat (All) <sup>1</sup>
Oats <sup>1</sup>	Wild Rice <sup>1</sup>
Rice <sup>2</sup>	

### Citrus

Calamondin	Mandarin Orange
Chironja	Orange (All)
Citron	Pummelo
Grapefruit	Tangelo
Kumquat	Tangerine
Lemon	Tangors
Lime	

### Tree Nuts

Almond	Filbert (Hazelnut)
Beechnut	Hickory Nut
Brazil Nut	Macadamia
Butternut	Pecan
Cashew	Pistachio
Chestnut	Walnut (Black, English)
Chinquapin	



**Vine Crops**

Grapes

Kiwi Fruit

**Tree Fruits**

Apple

Olive

Apricots

Peach

Cherry (Sweet, Sour)

Pear

Loquat

Plum/Prune (All)

Mayhaw

Quince

Nectarine

**Vegetables**

Artichoke, Jerusalem

Leek

Asparagus<sup>1</sup>

Lentils

Beans (All)

Lettuce

Beet Greens

Mango Melon<sup>3</sup>

Beets (Red, Sugar)

Melons (All)<sup>3</sup>

Broccoli (All)

Muskmelon<sup>3</sup>

Brussels Sprouts

Mustard Greens

Cabbage (All)

Okra

Cabbage, Chinese

Onion

Cantaloupe<sup>3</sup>

Parsley

Carrot

Parsnips

Cauliflower

Peas (All)

Casaba Melon<sup>3</sup>Pepper (All)<sup>3</sup>

Celeriac

Persian Melon<sup>3</sup>

Celery

Potato (Irish, Sweet)

Chard, Swiss

Pumpkin<sup>3</sup>

Chicory

Radish

Collards

Rape Greens

Crenshaw Melon<sup>3</sup>

Rhubarb

Cucumber<sup>3</sup>

Rutabaga

Eggplant<sup>3</sup>

Shallot

Endive

Spinach (All)

Garlic<sup>3</sup>Squash (Summer, Winter)<sup>3</sup>Gourds<sup>3</sup>Tomatillo<sup>3</sup>Ground Cherry<sup>3</sup>Tomato<sup>3,5</sup>Honeydew Melon<sup>3</sup>

Turnip

Honey Ball Melon<sup>3</sup>Watercress<sup>3</sup>

Horseradish

Watermelon<sup>3</sup>

Kale

Yams

Kohlrabi

**Small Fruits and Berries**

Blackberry

Elderberry

Blueberry

Gooseberry

Boysenberry

Huckleberry

Cranberry

Loganberry

Currant

Olallieberry

Dewberry

Raspberry (Black, Red)

## Forage Crops and Legumes

Alfalfa <sup>1</sup>                      Forage Legumes <sup>1</sup>  
Forage Grasses <sup>1</sup>

## Tropical Crops

Acerola	Longan
Atemoya	Lychee
Avocado	Mango
Banana (Plantains)	Papaya
Breadfruit	Passion Fruit
Canistel	Persimmons
Carambola	Pineapple <sup>4</sup>
Cherimoya	Pomegranate
Cocoa Beans	Sapodilla
Coffee	Sapote (Black, Mamey, White)
Dates	Soursop
Figs	Sugar Apple
Genip	Tamarind
Guava	Tea
Jaboticaba	
Jackfruit	

<sup>1</sup> Spot treatments may be applied in these crops.

<sup>2</sup> Do not treat rice fields or levees when the fields contain flood water.

<sup>3</sup> Apply only prior to planting. Allow at least 3 days between application and planting.

<sup>4</sup> Do not feed or graze treated pineapple forage following application.

<sup>5</sup> Use is restricted to direct seeded crops only.

When applying Glyphomax prior to transplanting crops into plastic mulch, care must be taken to remove residues of Glyphomax from the plastic prior to transplanting. Residues can be removed by 0.5 inch natural rainfall or by applying water via a sprinkler irrigation system.

**Spot Treatment** (Only those crops referenced by footnote "1" above can be spot treated.): Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn, or boll opening on cotton.

For forage grasses and forage legumes see "Spot Treatment" in the "Pastures" section of "Cropping Systems" in this label.

For dilution and rates of application using boom or hand-held equipment, see "Mixing, Additives and Application Instructions" and "Weeds Controlled" sections of this label.

**NOTE: For forage grasses and forage legumes, no more than one-tenth of any acre should be treated at one time. For all other crops, do not treat more than 10 percent of the total field area to be harvested.**

**The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.**

**Selective Equipment:** Glyphomax may be applied through recirculating sprayers, shielded applicators or wiper applicators in cotton and soybeans. Shielded and wiper applicators may also be used in tree crops and grapes. Wiper applicators may be used in wheat, rutabagas, forage grasses and forage legumes, including pasture sites and grain sorghum (milo).

See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.

**Allow at least the following time intervals between application and harvest:**

Cotton, Soybeans .....	7 days
Apples, Citrus, Pear .....	1 day
Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherry, Grapes, Dates, Jaboticaba, Jackfruit, Longan, Lychee, Passion Fruit, Persimmons, Rutabagas, Sapodilla, Sapote, Soursop, Sugar Apple, Tamarind .....	14 days
Stone Fruit .....	17 days
Nut Crops, except pistachios .....	3 days
Pistachio nuts.....	21 days
Wheat <sup>1</sup> .....	35 days
Sorghum (milo) <sup>1, 2</sup> .....	40 days

<sup>1</sup> Do not use roller applicators.

<sup>2</sup> Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

## Asparagus

When applied as directed for "Cropping Systems" under the conditions described, Glyphomax controls weeds listed on this label in asparagus.

For specific rates of applications and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

**Prior to Crop Emergence:** Apply Glyphomax prior to crop emergence for the control of emerged labeled annual and perennial weeds. **Do not apply within a week before the first spears emerge.**

**Spot Treatment:** Apply Glyphomax immediately after cutting, but prior to the emergence of new spears. Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

**Postharvest:** Apply Glyphomax after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. Direct contact of the spray with the asparagus may result in serious crop injury.

**NOTE:** Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

---

## Berries and Small Fruits

---

Wiper applicators may be used in cranberries in accordance with instructions in this section.

For other berries, apply as a preplant broadcast application, or as a directed spray or wiper application post-planting.

See "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information.

See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on recommended use and calibration of this equipment.

Allow a minimum of 30 days between last application and harvest of cranberries. For other small fruits and berries, allow a minimum of 14 days between last application and harvest.

**For Wick or other Wiper Applicators:** Mix 1 gallon of Glyphomax in 4 gallons of water to prepare a 20 percent solution.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of Glyphomax are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes or foliage.

---

## Corn

---

**Hooded Sprayers:** Glyphomax may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

**Follow these requirements:**

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of Glyphomax per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply Glyphomax when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of Glyphomax in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weed Control Section" of this label.

Do not graze or feed corn forage or fodder following applications of Glyphomax through hooded sprayers.

Do not apply more than 3 quarts of Glyphomax per acre per year for hooded sprayer applications.

### **Fallow and Reduced Tillage Systems**

For aerial application in California, refer to supplemental labeling.

Use Glyphomax in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the "Weeds Controlled" section of this label for specific rates and instructions. Glyphomax may be applied using ground or aerial spray equipment. See the "Application Equipment and Techniques" section of this label for instructions.

### **Tank Mixtures**

- **Glyphomax plus Banvel (dicamba) herbicide plus Nonionic Surfactant**
- **Glyphomax plus 2,4-D herbicide plus Nonionic Surfactant**
- **Glyphomax plus Tordon\* 22K herbicide plus Nonionic Surfactant**
- **Glyphomax plus Goal herbicide plus Nonionic Surfactant**
- **Glyphomax plus Atrazine or Bladex (cyanazine) plus Nonionic Surfactant**

### **Glyphomax plus Dicamba, 2,4-D or Tordon 22K Tank Mixtures**

**Do not apply Banvel (dicamba), Tordon 22K or 2,4-D tank mixtures by air in California.**

**Dicamba and 2,4-D:** The addition of dicamba in a mixture with Glyphomax may provide short-term residual control of selected weed species. Some crop injury may occur if dicamba is applied within 45 days of planting. Applications of 2,4-D or Banvel (dicamba) herbicide must be made at least 7 days prior to planting corn. Refer to the labels for dicamba and 2,4-D labels for cropping restrictions and other use instructions.

**Tordon 22K:** The addition of Tordon 22K in a mixture with Glyphomax may provide short-term residual control of selected weed species. Application of Glyphomax in tank mix with Tordon 22K should be made only to land that will be planted the following year to grass, barley, oats, wheat, grain sorghum (milo) or fallowed. Some crop injury may occur if Tordon 22K is applied within 45 days of planting. Do not plant grain sorghum within 8 months after application. Tordon 22K is not intended for use on land planted to sweet sorghum. Refer to the label Tordon 22K for cropping restrictions and other use instructions.

Glyphomax plus dicamba, Tordon 22K or 2,4-D will control the annual grasses and broadleaf weeds previously listed for Glyphomax alone at the indicated heights (except 8 fluid ounces per acre applications), plus the broadleaf weeds listed below in this section. For those weeds previously listed at 8 fluid ounces of Glyphomax alone per acre, use 12 fluid ounces in these tank mixtures.

Apply 12 to 24 fluid ounces of Glyphomax plus 0.25 pound a.i. of dicamba, or 1 to 2 ounces of Tordon 22K, or 0.5 pound a.i. of 2,4-D plus 0.5 percent nonionic surfactant by total spray volume per acre to control dense populations of the following annual broadleaf weeds when less than 6 inches in height:

Common Name	Scientific Name
Buckwheat, Wild <sup>1</sup>	<i>Polygonum convulvulus</i>
Kochia <sup>2</sup>	<i>Kochia scoparia</i>
Lambsquarters	<i>Chenopodium album</i>
Lettuce, prickly <sup>3</sup>	<i>Lactuca serriola</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
Thistle, Russian <sup>3</sup>	<i>Salsola kali</i>

<sup>1</sup> Controlled with Tordon 22K tank mixture only.

<sup>2</sup> Controlled with dicamba tank mixture only.

<sup>3</sup> Controlled with dicamba or 2,4-D tank mixtures only.

#### Glyphomax plus Goal Tank Mixtures

Glyphomax alone or in tank mixtures with Goal plus 0.5 to 1 percent nonionic surfactant by total spray volume will provide control of those weeds listed below.

Make applications when weeds are actively growing and at the recommended stages of growth. Avoid spraying when weeds are subject to moisture stress, when dust is on the foliage or when straw canopy covers the weeds.

Glyphomax (12 fl oz/acre)	Maximum Height or Length in Inches	Glyphomax (16 fl oz/acre)	Maximum Height or Length in Inches	
Wheat	18"	<b>Annual grasses at left plus:</b>		
Barley	12"			
Bluegrass, annual	6"		Ryegrass, annual	6"
Barnyardgrass	6"		Chickweed	6"
Rye	6"		Groundsel	6"
			Marestail	6"
			Rocket, London	6"
			Shepherdspurse	6"
			Crabgrass	12"
			Johnsongrass, seedling	12"
			Lambsquarters	12"
			Oats, wild	12"
			Pigweed, redroot	12"
		Mustards	12"	

NOTE: Use 32 fluid ounces of Glyphomax per acre where heavy weed densities exist.

Glyphomax (12 fl oz/acre) plus Goal <sup>1</sup> (2 to 4 fl oz/acre)	Maximum Height or Length in Inches	Glyphomax (16 fl oz/acre) plus Goal <sup>1</sup> (2 to 4 fl oz/acre)	Maximum Height or Length in Inches
<b>Annual grasses above plus:</b>		<b>Annual weeds above plus:</b>	
Cheeseweed, common	3"	Cheeseweed, common	6"
Chickweed	3"	Groundsel	6"
Groundsel	3"	Chickweed	12"
Rocket, London	6"	Rocket, London	12"
Shepherdspurse	6"	Shepherdspurse	12"

NOTE: Use 32 fluid ounces of Glyphomax per acre in mixtures with 2 to 4 fluid ounces of Goal per acre where heavy weed densities exist.

<sup>1</sup> Use the higher rate of Goal when weeds approach maximum recommended height or stands are dense.

These recommended tank mixtures may be applied using ground or aerial spray equipment. Refer to the "Weeds Controlled" section of this label for specific rates and instructions.

#### **Glyphomax plus Atrazine or Bladex (cyanazine) Tank Mixtures**

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound a.i. of atrazine per acre. The addition of 2 percent dry ammonium sulfate by weight (or liquid equivalent) or 17 pounds per 100 gallons of water is recommended to increase the performance of Glyphomax plus atrazine or Bladex plus nonionic surfactant tank mixtures.

Consult the labels for atrazine or Bladex (90 OF, 4L, 80W) for use rates, soil type, planting, cropping and other restrictions, as well as other precautionary statements and use according to the most restrictive label.

Glyphomax plus atrazine or Bladex plus nonionic surfactant tank mixture provides postemergence control of listed annual weeds in fallow and reduced tillage systems. Apply 12 fluid ounces of Glyphomax with 1 pound a.i. or less of atrazine per acre or 16 fluid ounces of Glyphomax with 2 pounds a.i. or less of atrazine per acre, or 20 fluid ounces of Glyphomax with 1 to 3 pounds a.i. of atrazine per acre, plus 0.5 percent nonionic surfactant by total spray volume. Apply 16 fluid ounces of Glyphomax with 2.4 to 4 pounds a.i. of Bladex (cyanazine) per acre plus 0.5 percent nonionic surfactant. These tank mixtures will provide soil residual control of weeds listed on the atrazine and Bladex label. In addition, these tank mixtures will provide postemergence control of the following weeds:

Common Name	Scientific Name
Barnyardgrass <sup>1</sup>	<i>Echinochloa crus-galli</i>
Brome, downy	<i>Bromus tectorum</i>
Foxtail, green	<i>Setaria, viridis</i>
Kochia <sup>2</sup>	<i>Kochia scoparia</i>
Lambsquarters	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mustard, tansy	<i>Descurainia pinnata</i>
Pigweed	<i>Amaranthus spp.</i>
Sandbur, field	<i>Cenchrus spp.</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Thistle, Russian	<i>Salsola iberica</i>
Wheat	<i>Triticum aestivum</i>
Witchgrass	<i>Panicum capillare</i>

<sup>1</sup> Barnyardgrass will be suppressed at 20 fluid ounces per acre and will require up to 26 fluid ounces per acre for control.

<sup>2</sup> For improved control of kochia, add 4 fluid ounces per acre (.125 pound a.i. per acre) of dicamba to the above tank mixture.

### Postharvest Applications

Glyphomax will provide control of existing weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvest operations, and to recover from environmental stress, before application. Weeds should be treated prior to the heading stage of annual grasses, if possible, and before broadleaf weeds exceed 24 inches in height. Nonionic surfactant should be added at 0.5 percent by total spray volume. Use of ammonium sulfate is recommended and may improve performance on annual weeds.

Apply 12 fluid ounces of Glyphomax per acre plus nonionic surfactant to control downy brome, green foxtail, stinkgrass and volunteer wheat.

To control kochia, lambsquarters, mustard, pigweed and Russian thistle, apply 16 fluid ounces of Glyphomax plus 0.25 pounds a.i. of dicamba or 0.5 pounds a.i. of 2,4-D plus nonionic surfactant per acre.

Apply 20 fluid ounces of Glyphomax per acre plus nonionic surfactant to control barnyardgrass, sandbur, sunflower, and witchgrass. The addition of 0.25 pound a.i. of dicamba or 0.5 pound a.i. of 2,4-D plus nonionic surfactant to 20 fluid ounces of Glyphomax will control prickly lettuce.

**Grain Sorghum (Postharvest Regrowth Control):** Glyphomax may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of Glyphomax per acre for control, or 1.5 pints of Glyphomax per acre for suppression. Use 0.5 percent nonionic surfactant in 3 to 10 gallons of spray solution per acre.

### Ecofarming Systems

The recommendations made in this section are not registered for use in California.

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixture for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System:



**Glyphomax at 16 to 20 fluid ounces per acre**  
**plus**  
**2,4-D at 0.375 to 0.5 pound a.i. per acre**  
**plus**  
**Atrazine herbicide at 0.75 to 1 pound a.i. per acre**  
**plus**  
**Lasso herbicide at 2.5 to 3 quarts per acre**

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume.

**Weeds Controlled:** The following weeds, up to a maximum height of 4 inches, will be controlled:

Common Name	Scientific Name
Brome, downy	<i>Bromus tectorum</i>
Cheat	<i>Bromus secalinus</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria lutescens</i>
Kochia <sup>1</sup>	<i>Kochia scoparia</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Thistle, Russian	<i>Salsola kali</i>
Wheat, volunteer	<i>Triticum aestivum</i>

<sup>1</sup> For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of dicamba to the above tank mixture.

Risk of crop injury from 2,4-D or dicamba can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the label booklet for Lasso for preemergence weed control achieved by this tank mixture.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

### Aid To Tillage

Glyphomax, when used in conjunction with preplant tillage practices, will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. For ground application, apply in 3 to 10 gallons of water per acre. For aerial applications, apply in 3 to 5 gallons of water per acre. Do not tank mix with residual herbicides. Make applications when weeds are actively growing and before they are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage. Tank mixtures with residual herbicides may result in reduced performance. Glyphomax does not provide residual weed control. Repeat treatments may be necessary to control subsequent germinating weeds.

Do not treat weeds under poor growing conditions such as drought stress, or disease or insect damage as reduced weed control may result. Reduced results also may occur when treating weeds heavily covered with dust, or when straw canopy covers the weeds.

### Perennial Weeds

Apply Glyphomax as follows to control or destroy most perennial weeds:

**Note:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages of growth. Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this herbicide on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.

When applied as recommended under the conditions described, Glyphomax will control the following perennial weeds:

Common Name	Scientific Name
Alfalfa	<i>Medicago saliva</i>
Johnsongrass	<i>Sorghum halepense</i>
Larkspur, duncecap	<i>Delphinium occidentale</i>
Larkspur, tall	<i>Delphinium barbeyi</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Quackgrass	<i>Agropyron repens</i>
Ryegrass, perennial	<i>Lolium perenne</i>
Spurge, leafy <sup>1</sup>	<i>Euphorbia esula</i>
Thistle, Canada	<i>Cirsium arvense</i>

<sup>1</sup> Partial control

**Johnsongrass / Ryegrass, perennial:** Apply 1 to 3 quarts of Glyphomax per acre. In annual cropping systems, apply 1 to 2 quarts of Glyphomax per acre. Apply 1 quart of Glyphomax plus 0.5 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Use 2 quarts of Glyphomax when applying 10 to 40 gallons of water per acre. In noncrop or areas where annual tillage (no-till), is not performed, apply 2 to 3 quarts of Glyphomax in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart per acre rate.

For burndown of contacted vegetation, apply 1 pint per acre plus 0.5 percent nonionic surfactant in 5 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

**Larkspur, duncecap / Larkspur, tall:** Apply Glyphomax as a 2 percent solution using hand-held equipment. Apply as a spot treatment when the larkspur is actively growing and in the vegetative stage just prior to budding. Coverage should be uniform and complete.

**Quackgrass (In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage):** Apply 1 to 2 quarts of Glyphomax per acre. For the one quart rate, apply 0.5 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank-mix with residual herbicides when using the 1 quart rate. Spray when

quackgrass is 6 to 8 inches tall and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results, use a moldboard plow.

**Quackgrass (Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application):** Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in the fall or spring prior to spring application. Allow 3 or more days after application before tillage.

**Spurge, leafy:** For suppression, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D per acre plus 0.5 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

**Thistle, Canada:** Apply 2 to 3 quarts of Glyphomax per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Glyphomax. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression of Canada thistle, apply 1 quart per acre of Glyphomax plus 0.5 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

**Other perennials listed on this label:** Apply 3 to 5 quarts of Glyphomax per acre. Apply when actively growing and when most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

See "Directions for Use" and "Mixing Additives, and Application Instructions" sections of this booklet for labeled uses and specific application instructions.

#### Tank Mixtures

When applied as directed under the conditions described, Glyphomax plus dicamba or 2,4-D will **suppress** or **control** the following perennial broadleaf weed:

#### **Bindweed, field** (*Convolvulus arvensis*)

For suppression, apply 16 fluid ounces of Glyphomax plus 0.5 pound a.i. of 2,4-D or 0.25 pound a.i. of dicamba, plus 0.5 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length. These tank mixtures may be applied using aerial application equipment in fallow and reduced tillage systems only.

For control, apply 2 quarts of Glyphomax plus 0.5 pound a.i. of dicamba in 3 to 20 gallons of water per acre. At these rates, apply using ground application equipment only.

---

## Pastures

---

Apply Glyphomax prior to planting forage grasses and legumes.

**Pasture or Hay Crop Renovation:** When applied as a broadcast spray, Glyphomax controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

**Spot Treatment:** When applied as a spot treatment as recommended, Glyphomax controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

**Wiper Application:** When applied as directed, Glyphomax controls or suppresses the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label.

For spot treatment and wiper application, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

---

## Sugarcane

---

When applied as directed for "Cropping Systems", under the conditions described, Glyphomax controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields prior to the emergence of plant cane. Glyphomax will also control undesirable sugarcane.

**NOTE:** Where repeat treatments are necessary, do not exceed a total of 10.6 quarts of Glyphomax per acre per year. Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

**Broadcast Treatment:** Apply Glyphomax in 10 to 40 gallons of water per acre on emerged weeds prior to the emergence of plant cane.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

For removal of last stubble or ratoon cane, apply 4 to 5 quarts of Glyphomax in 10 to 40 gallons of water per acre to new growth having at least 7 or more new leaves. Allow 7 or more days after application before tillage.

**Spot Treatment in or Around Sugarcane Fields:** For dilution and rates of application using hand-held equipment, see "Mixing, Additives and Application Instructions" and "Weeds Controlled" sections of this label.

For control of volunteer or diseased sugarcane, make a 1 percent solution of Glyphomax in water and spray to wet the foliage of vegetation to be controlled.

**NOTE:** When spraying volunteer or diseased sugarcane, the plants should have at least 7 new leaves.

Avoid spray contact with healthy cane plants since severe damage or destruction may result.

Do not feed or graze treated sugarcane forage following application.

## **Conservation Tillage, Minimum Tillage and No-Till Systems**

### **Corn and Soybeans (Tank Mixtures)**

The recommendations made in this section are not registered for use in California.

When applied as recommended under the conditions described, the tank mixtures listed in this section control many emerged weeds, and give preemergence control of many annual weeds where corn or soybeans will be planted directly into a cover crop, established sod or in previous crop residues.

Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures. For mixing instructions, see the "Mixing, Additives and Application Instructions" section of this label.

Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre before, during or after planting. Do not apply these mixtures after crop emergence.

When tank mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. The addition of 1 to 2 percent dry ammonium sulfate by weight may increase the performance of Glyphomax.

**NOTE:** When using these tank mixtures, do not exceed 4 quarts of Glyphomax per acre.

#### **Corn**

For residual control, Glyphomax may be tank-mixed with the following herbicides or combination of herbicides:

Lasso/Alachlor	Atrazine
Lariat	Cyanazine
Bullet	Simazine
Dual	Prowl
Bicep	Micro-Tech
Partner <sup>1</sup>	Hornet*

For improved burndown, Glyphomax may be tank-mixed with 2,4-D or dicamba. Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. See the "Weeds Controlled" section for specific rate information.

<sup>1</sup> Partner herbicide is not registered in California.

## Soybeans

**For residual control, Glyphomax may be tank-mixed with the following herbicides or combination of herbicides:**

Canopy	Partner
Command	Preview
Dual	Prowl
FirstRate*	Pursuit
Frontrow*	Pursuit Plus
Gemini	Python*
Lasso/Alachlor	Scepter
Lexone	Sencor
Linuron	Squadron
Lorox Plus	Turbo
Micro-Tech	

**For improved burndown, Glyphomax may be tank-mixed with the following herbicides:**

2,4-DB  
2,4-D<sup>1</sup>

<sup>1</sup> See the label for 2,4-D for intervals between application and planting.

## Corn and Soybeans

**Annual Weeds:** For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Glyphomax at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of Glyphomax per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. For a complete list of annual weeds controlled, see the "Weeds Controlled" section of this label.

**Perennial Weeds:** At normal application times in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "Weeds Controlled" section of this label for the proper stage of growth for perennial weeds.

Use of 2 to 4 quarts of Glyphomax per acre in the tank mixtures mentioned above, under these conditions provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled, see the "Weeds Controlled" section of this label.

To obtain the desired stage of growth, it may be necessary to apply Glyphomax alone in the late summer or fall and then follow with a label-approved, seedling weed-control program at planting.

**Use of these tank mixtures for bermudagrass or johnsongrass control in minimum tillage systems is not recommended.** For bermudagrass control, follow the instructions under "Control of Perennial Weeds" section of this label and then use a label-approved, seedling weed-control program in a minimum tillage or conventional tillage system. For Johnsongrass control, follow instructions under "Control of Perennial Weeds" section of this label, and then use a label-approved, seedling weed-control program with conventional tillage.

## **Preharvest Applications**

When applied as directed under the conditions described, Glyphomax controls annual and perennial weeds listed on this label prior to the harvest of cotton, soybeans and wheat.

For specific rates and application instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

Glyphomax may be applied by both ground and aerial application equipment. **Do not apply more than 1 quart per acre of Glyphomax by air.** See the "Application Equipment and Techniques" section of this label for instructions for ground and aerial applications.

NOTE: Do not apply to crops grown for seed. Reduction in germination or vigor may occur.

**The use of Glyphomax for preharvest grain sorghum (Milo) is not registered in California.**

### **Soybeans**

Apply after pods have set and lost all green color. Allow a minimum of 7 days between application and harvest of soybeans. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

Do not graze or harvest treated crop for livestock feed within 25 days of last preharvest application.

**Do not apply more than 6 quarts per acre of Glyphomax for preharvest applications.**

### **Cotton**

**Broadcast Applications:** Glyphomax may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply Glyphomax in 10 to 20 gallons of water per acre. For aerial applications, apply Glyphomax in 3 to 10 gallons of water per acre.

Glyphomax provides weed control and cotton regrowth inhibition when applied prior to the harvest of cotton. Apply 1 to 2 quarts of Glyphomax in 3 to 10 gallons of water per acre for cotton regrowth inhibition. Do not apply more than 2 quarts of Glyphomax per acre for preharvest applications. **The use of additives for preharvest application to cotton is prohibited.**

Glyphomax may be tank mixed with DEF 6, Folex, or Prep defoliant to provide additional enhancement of cotton leaf drop.

Allow a minimum of 7 days between application and harvest of cotton.

Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

Do not feed or graze treated cotton forage or hay following preharvest applications.

## Grain Sorghum (Milo)

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

Apply up to 2 quarts of Glyphomax per acre.

## Wheat

Apply after the hard-dough stage of grain (30% or less grain moisture) and at least 7 days prior to harvest.

For control of quackgrass or suppression of Canada thistle, apply 1 quart of Glyphomax plus 0.5% nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

For suppression of field bindweed, apply 1 to 2 pints of Glyphomax plus 0.5 to 1.0 pounds a.i. of 2,4-D plus 0.5% nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

**Do not apply more than 1 quart per acre of Glyphomax for preharvest applications to wheat.**

---

## Tree and Vine Crops

---

Glyphomax is recommended for weed control in established groves, vineyards, and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed in this section. See the "Application Equipment and Techniques" section of this label for specific information on use of equipment.

When applying Glyphomax, refer to the "Weeds Controlled" section of this label and to specific recommendations in this section for rates to be used.

### NOTE:

Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. Glyphomax does not provide residual weed control. For subsequent weed control, use repeated applications of Glyphomax. Do not apply more than 10.6 quarts of Glyphomax per acre per year.

**Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees or vines. Contact of Glyphomax with other than matured brown bark can result in serious crop damage.**

**Avoid painting out stumps with Glyphomax as injury resulting from root grafting may occur in adjacent trees.**

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

For specific rates of applications and instructions, see the "Weeds Controlled" section of this label, and to specific recommendations which follow.



**Middles Management**  
**(For Annual Weeds in Middles Between**  
**Rows of Tree and Vine Crops)**

For citrus crops, treat uniformly between trees.

- **Glyphomax**
- **Glyphomax plus Goal**

Glyphomax alone or in mixtures with Goal will control or suppress the annual weeds listed below.

Apply the recommended rates of Glyphomax, either alone or in mixtures with Goal, plus 0.5 to 1 percent nonionic surfactant by spray volume in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and less than 6 inches in height or diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fluid ounces per acre of Glyphomax may be used to control weeds, which have been mowed, are stressed or are growing in dense populations.

Weed Species	Maximum Height or Diameter (inches)	Glyphomax (fl oz/acre)	Goal (fl oz/acre)
<b>Barley</b> <i>Hordeum vulgare</i>	6	8	-
<b>Bluegrass, annual</b> <i>Poa annua</i>			
<b>Barnyardgrass</b> <i>Echinochloa crus-galli</i>	6	12	-
<b>Chickweed, common</b> <i>Stellaria media</i>			
<b>Red Maids</b> <i>Calandrinia ciliata</i>			
<b>Crabgrass</b> <i>Digitaria spp.</i>	6	16	-
<b>Fleabane, hairy</b> <i>Conyza bonariensis</i>		16 to 32	or + 4 to 16 <sup>1</sup>
<b>Groundsel, common</b> <i>Senecio vulgaris</i>			
<b>Junglerice</b> <i>Echinochloa colonum</i>			
<b>Lambsquarters, common</b> <i>Chenopodium album</i>			
<b>Pigweed, redroot</b> <i>Amaranthus retroflexus</i>			
<b>Rocket, London</b> <i>Sisymbrium irio</i>			
<b>Ryegrass, common</b> <i>Lolium multiflorum</i>			
<b>Shepherdspurse</b> <i>Capsella bursa-pastoris</i>			
<b>Sowthistle, annual</b> <i>Sonchus oleraceus</i>			

<b>Cheesweed, common</b> <i>Malva spp.</i>	3	12 to 32	+	4 to 16
<b>Cheesweed, common</b> <i>Malva spp.</i>	6	16 to 32	+	4 to 16
<b>Filaree</b> <sup>2</sup> <i>Erodium spp.</i>				
<b>Horseweed/Marestail</b> <i>Conyza canadensis</i>				
<b>Nettle, stinging</b> <i>Urtica dioica</i>				
<b>purslane, common</b> <sup>2</sup> <i>Purtulaca oleracea</i>				

<sup>1</sup> The mixture of Glyphomax plus Goal is recommended when weeds are stressed or growing in dense populations.

<sup>2</sup> Suppression only.

### Strips

(For Annual and Perennial Weeds in Strips of Tree and Vine Crops)

### Tank Mixtures With Residual Herbicides

When applied as a tank mixture, Glyphomax provides control of the emerged annual weeds and control or suppression of emerged perennial weeds listed in this label. The following residual herbicides will provide preemergence control of those weeds listed in the individual product labels.

- Glyphomax plus Goal 2XL
- Glyphomax plus Karmex DF
- Glyphomax plus Krovar I
- Glyphomax plus Krovar II
- Glyphomax plus Simazine, Princep Caliber 90
- Glyphomax plus Simazine 4L
- Glyphomax plus Simazine 80W
- Glyphomax plus Solicam 80DF
- Glyphomax plus Surflan A.S.
- Glyphomax plus Simazine (80W, Or 4L, or Princep Caliber 90) plus Surflan A.S.
- Glyphomax plus Goal 2XL plus Surflan A.S.
- Glyphomax plus Goal 2XL plus Simazine (80W, Or 4L, or Princep Caliber 90)
- Glyphomax plus Goal 2XL plus Surflan A.S. plus Simazine (80W, 4L, or Princep Caliber 90)

Do not apply these tank mixtures in Puerto Rico.

When tank-mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution.

Refer to the individual product labels for specific crops, rates, geographical restrictions and precautionary statements.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of all products.

## Recommended Rates

**Annual Weeds:** Apply 1 to 5 quarts per acre of Glyphomax in these tank mixtures. Use rates at the higher end of the recommended range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

**Perennial Weeds:** Apply 1 pint to 5 quarts per acre of Glyphomax in these tank mixtures to control or suppress perennial weeds. Follow the recommendations in the "Weeds Controlled" section of this label for stage of growth and application rates for specific perennial weeds.

## Glyphomax plus Goal plus Simazine/Surflan

Glyphomax plus low rates of Goal in 3-way or 4-way mixtures with simazine and/or Surflan will provide postemergence control of the weeds listed below.

Refer to the individual simazine and Surflan labels for preemergence rates, weeds controlled, precautionary statements and other important information.

Apply these tank mixtures in 3 to 40 gallons of water. Add 0.5 to 1 percent nonionic surfactant by total spray volume to the spray solution.

Apply 1 to 5 quarts per acre of Glyphomax plus 4 to 48 fluid ounces per acre of Goal plus labeled rates of simazine and/or Surflan to control the following weeds:

Common Name	Scientific Name
Barley, wild	<i>Hordeum leporinum</i>
Bluegrass, annual	<i>Poa annua</i>
Cheeseweed, common	<i>Malva spp.</i>
Chickweed, common	<i>Stellaria media</i>
Filaree <sup>1</sup>	<i>Erodium spp.</i>
Fleabane, hairy	<i>Conyza bonariensis</i>
Groundsel, common	<i>Senecio vulgaris</i>
Horseweed/Marestail	<i>Conyza canadensis</i>
Nettle, stinging	<i>Urtica dioica</i>
Pineappleweed	<i>Matricaria matricariodes</i>
Rocket, London	<i>Sisymbrium irio</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>

<sup>1</sup> Use a minimum of 1.5 quarts of Glyphomax in these mixtures.

**NOTE:** This recommendation does not preclude the use of Goal in these mixtures at higher, labeled rates for preemergence weed control.

## **Perennial Grass Suppression (Orchard Floors)**

When applied as directed, Glyphomax will suppress vegetative growth as indicated below.

### **Bahiagrass**

*Glyphomax will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with a single application and approximately 120 days with sequential applications. Apply Glyphomax 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.*

Sequential applications of Glyphomax plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of Glyphomax plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces may be made approximately 45 days after the last application.

### **Bermudagrass**

For burndown, apply 1 to 2 quarts of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. Use 1 quart of Glyphomax in 3 to 20 gallons of water per acre east of the Rocky Mountains. Use 1 to 2 quarts of Glyphomax in 3 to 10 gallons of water per acre west of the Rocky Mountains. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

**Suppression only (east of the Rocky Mountains):** Apply 6 to 16 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Rates of 6 to 10 fluid ounces of Glyphomax plus nonionic surfactant should be used in shaded conditions or where a lesser degree of suppression is desired. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

**Suppression only (west of the Rocky Mountains):** Apply 16 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches in height and no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

### **Cool Season Grass Covers**

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. For best suppression, add ammonium sulfate to the spray solution at a rate of 2 percent by weight or 17 pounds per 100 gallons of spray solution.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of Glyphomax plus 0.5 to 1 percent nonionic surfactant. Do not add ammonium sulfate.

For best results, mow cool-season grass covers in the spring to even their height and apply the recommended rate of Glyphomax 3 to 4 days after mowing. Avoid treating cool season grass covers under poor growing conditions, such as drought stress (drip irrigation), disease or insect damage.

### Low Volume Application (Florida and Texas)

For burndown or control of the weeds listed, apply the recommended rates of Glyphomax plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

#### Annual Weeds

**Goatweed:** Apply 2 to 3 quarts per acre of Glyphomax plus 17 pounds of ammonium sulfate per 100 gallons of water plus 0.5 to 1 percent nonionic surfactant by total spray volume. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches. If goatweed is greater than 8 inches tall, the addition of Krovar II or Karmex may improve control. Use labeled rates for these residual products.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the Krovar II and Karmex labels.

#### Perennial Weeds

Apply when weeds are actively growing and at the growth stages listed in the "PERENNIAL Weeds Controlled" section of this label. If perennial weeds are mowed, allow weeds to regrow to the recommended stage of growth.

S = Suppression      B = Burndown  
PC = Partial Control      C = Control

Weed Species	Glyphomax Rate Per Acre			
	1 qt	2 qt	3 qt	5 qt
Bermudagrass	B	•	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	•	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	•	PC	C

### Tree Crops

**Citrus**<sup>5</sup>: calamondin, chironja, citron, grapefruit, kumquat, lemon, lime, mandarin orange, orange, pummelo, tangelo, tangerine, tangors.

**Nuts**<sup>2</sup>: almond, beechnut, Brazil nut, butternut, cashew, chestnuts, chinquapin, filbert, hazel nut, hickory nut, macadamia, pecan, pistachio, walnut.

**Pome Fruit**<sup>5</sup>: apple, loquat, mayhaw, pear, quince.

**Stone Fruit**<sup>3</sup>: apricots, cherries, nectarines, olives, peaches, plums/prunes.

For cherries, any application equipment listed in this section may be used in all states.

For citron and olives, apply as a directed spray only.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use wiper equipment only.

For **peaches** grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of Glyphomax with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. **Extreme care must be taken to ensure no part of the peach tree is contacted.**

**Tropical Fruit:** acerola <sup>1</sup>, atemoya <sup>1</sup>, avocado <sup>1</sup>, banana <sup>5</sup>, (plantains) <sup>4</sup>, breadfruit <sup>1</sup>, canistel <sup>1</sup>, carambola <sup>1</sup>, cherimoya <sup>1</sup>, cocoa beans <sup>1</sup>, coffee <sup>4</sup>, dates <sup>1</sup>, figs <sup>1</sup>, genip <sup>1</sup>, guava <sup>5</sup>, jaboticaba <sup>1</sup>, jackfruit <sup>1</sup>, longan <sup>1</sup>, lychee <sup>1</sup>, mango <sup>1</sup>, mayhaw <sup>1</sup>, papaya <sup>5</sup>, passion fruit <sup>1</sup>, persimmons <sup>1</sup>, pomegranate <sup>1</sup>, sapodilla <sup>1</sup>, sapote <sup>1</sup>, soursop <sup>1</sup>, sugar apple <sup>1</sup>, tamarind <sup>1</sup>, tea <sup>1</sup>. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

**NOTE:**

- <sup>1</sup> Allow a minimum of 14 days between last application and harvest.
- <sup>2</sup> Allow a minimum of 3 days between last application and harvest of these crops, except pistachio nuts. For pistachio nuts allow a minimum of 21 days between last application and harvest.
- <sup>3</sup> Allow a minimum of 17 days between last application and harvest.
- <sup>4</sup> Allow a minimum of 28 days between last application and harvest.
- <sup>5</sup> Allow a minimum of 1 day between last application and harvest.

---

### Vine Crops

---

**Kiwi Fruit**

**Grapes:** Any variety of table, wine or raisin grape may be treated with any equipment listed in this section.

Applications should not be made when green shoots, canes, or foliage are in the spray zone.

Allow a minimum of 14 days between last application and harvest.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

---

### Warranty Disclaimer

---

Dow AgroSciences warrants that Glyphomax conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of Glyphomax. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from Glyphomax (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of Glyphomax unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC  
EPA-accepted \_\_/\_\_/\_\_