

62719-346

08/12/2005

4/1

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Propiconazole EC

EPA Reg. No. 62719-346

Application through Feekes Growth Stage 10.5 for Disease Control in Wheat


ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Propiconazole* EC before applying. Follow all other applicable directions, restrictions, and precautions on the EPA-registered label.
- Use of Propiconazole* EC according to this supplemental labeling is subject to all applicable use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Propiconazole EC at the rate of 4 fl oz per acre controls powdery mildew (Caused by *Erysiphe graminis* f. sp. *tritici*), leaf and glume blotch (*Septoria tritici* and *Stagonospora nodorum*) and leaf rust (*Puccinia recondita*). Propiconazole EC at the rate of 4 fl oz per acre provides suppression (ca. 50% control) of fusarium head blight (*Fusarium* spp.). Application may be made through Feekes Growth Stage 10.5 (full head emergence) using ground or aerial equipment. **Do not** apply after this crop growth stage to avoid possible illegal residues. Propiconazole EC may be applied earlier if disease symptoms appear.

Note: To avoid possible illegal residues: (1) **Do not** apply more than 4 fl oz of Propiconazole EC per acre per season. (2) **Do not** apply within 40 days of harvest. (3) **Do not** graze or feed livestock treated wheat, straw, forage, or hay. (4) **Do not** cut the green crop for hay or silage. (5) After harvest, the straw may be used for bedding.



R-142-025
EPA-accepted: / /
Initial printing.