

62719-346

11/19/2007

5818350 1/39



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Raymond S. Brinkmeyer
Regulatory Leader
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

NOV 19 2007

Subject: Propiconazole EC
EPA Reg. No. 62719-346
Your amendment dated April 26, 2007
EPA Decision Number 379253

Dear Dr. Brinkmeyer:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. In the section "Hazards to Humans and Domestic Animals" (pages 1, 4, and 7) add the signal word "Warning" immediately before "Causes Substantial But Temporary...."
2. Move the First Aid box to above the Precautionary Statements (pages 1, 4, and 7)
3. In the section Inherent Risks of Use (page 38) change "All such risks shall be assumed by buyer." to "To the extent consistent with applicable law, all such risks shall be assumed by buyer."
4. Reformat the Limitation of Remedies section as follows:

"To the extent consistent with applicable law:

- a) The exclusive remedy for losses...one of the following:
 - i. Refund of purchase price....
 - ii. Replacement of amount of product used.
- b) Dow AgroSciences shall not be liable....
- c) Dow AgroSciences shall not be liable for consequential or incidental damages or losses."

One copy of the label stamped "Accepted with comments" is enclosed for your records. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

2/39

(Base label for rigid containers 5 gal or less):

Propiconazole EC

Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereals, corn, cranberry, mint, nonbearing citrus fruits, onions, peanut, pecan, pineapple, rice, sorghum, stone fruits, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and non-bearing fruit and nut trees in nurseries and landscape settings.

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole	41.8%
Other Ingredients	58.2%
Total	100.0%

Contains petroleum distillates.
Contains 3.6 lb active ingredient per gallon.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

NOV 19 2007

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

62719-346

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Physician: There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Refer to product labeling for use restrictions to protect endangered species.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

4/30

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346

EPA Est. _____

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents ____

(Base label for refillable rigid containers larger than 5 gal):

Propiconazole EC

Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereals, corn, cranberry, mint, nonbearing citrus fruits, onions, peanut, pecan, pineapple, rice, sorghum, stone fruits, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and non-bearing fruit and nut trees in nurseries and landscape settings.

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole	41.8%
Other ingredients	58.2%
Total	100.0%

Contains petroleum distillates.
Contains 3.6 lb active ingredient per gallon.

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

6/39

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Physician: There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Refer to product labeling for use restrictions to protect endangered species.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

7/39

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346

EPA Est. _____

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents _____

(Base label for nonrefillable rigid containers larger than 5 gal):

Propiconazole EC

Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereals, corn, cranberry, mint, nonbearing citrus fruits, onions, peanut, pecan, pineapple, rice, sorghum, stone fruits, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and non-bearing fruit and nut trees in nurseries and landscape settings.

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole	41.8%
Other Ingredients	58.2%
Total	100.0%

Contains petroleum distillates.

Contains 3.6 lb active ingredient per gallon.

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Physician: There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Refer to product labeling for use restrictions to protect endangered species.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

10/39

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346

EPA Est. _____

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents _____

11/39

(Label booklet cover):

Propiconazole EC

Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereals, corn, cranberry, mint, nonbearing citrus fruits, onions, peanut, pecan, pineapple, rice, sorghum, stone fruits, strawberry, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and non-bearing fruit and nut trees in nurseries and landscape settings.

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-	
4-propyl-1,3-dioxolan-2-yl]methyl]-	
1H-1,2,4-triazole	41.8%
Other Ingredients	58.2%
Total	100.0%

Contains petroleum distillates.
Contains 3.6 lb active ingredient per gallon.

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346

EPA Est. _____

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents _____

12/39

[Page 1 to end of label booklet]

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
Engineering Controls	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Physical or Chemical Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Non-Agricultural Use Requirements	-
Storage and Disposal	-
Crops	-
General Information	-
Resistance Management	-
Spray Equipment	-
Mixing Instructions	-
Application Instructions	-
Rotational Crops	-
Uses	-
Almond	-
Berries	-
Carrot	-
Celery	-
Cereals	-
Corn (Field Corn, Corn Grown for Seed, Sweet Corn, and Popcorn)	-
Cranberry	-
Mint (Peppermint, Spearmint)	-
Nonbearing Citrus Fruits	-
Onions	-
Peanut	-
Pecan	-
Pineapple	-
Rice	-
Sorghum	-
Stone Fruits	-
Sugarcane	-
Tree Nuts	-
Turfgrass Grown for Seed	-
Turfgrass and Ornamentals	-
General Information	-
General Use Precautions and Restrictions	-
Mixing Instructions	-
Uses	-
Turfgrass and Dichondra	-
Ornamentals and Non-Bearing Fruit and Nut Trees in Nurseries (Field) and Landscape Plantings	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

13/39

(Page 1 through the end)

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING**

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

14/39

Note to Physician: There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Refer to product labeling for use restrictions to protect endangered species.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter into treated areas without protective clothing until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

15/39

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

Nonrefillable containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or larger:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers 5 gallons or larger:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Crops

General Information

Propiconazole EC fungicide is a broad-spectrum fungicide for the control of certain diseases in cereals (wheat, barley, rye, oats, rice, and wild rice), celery, corn, nonbearing citrus, stone fruit (apricots, nectarines, peaches, plums, and prunes), peanuts, pecans, pineapple, sugarcane, and in grasses grown for seed.

Important: Do not use as a tree injection.

Failure to follow directions and precautions on this label may result in crop injury, poor disease control, or illegal residues.

Resistance Management

Propiconazole EC belongs to the sterol demethylation inhibitor (DMI) class of fungicides. Since certain fungi can develop resistance to this class of products, the use of Propiconazole EC should be part of a resistance management strategy which includes alternation and/or mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Spray Equipment

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use.

Use a pump with sufficient capacity to maintain 35 to 40 psi at nozzles and provide sufficient agitation in tank to keep mixture in suspension (this requires recirculation of 10% of tank volume per minute). Use a jet agitator or liquid sparger tube for agitation. Do not use air sparging.

Although Propiconazole EC is an emulsifiable concentrate, it is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and, where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

Mixing Instructions

Application Rate Reference Table

Application Rate of Propiconazole EC (fl oz/acre)	Active Ingredient Equivalent (lb ai/acre)	Acres per Gallon of Propiconazole EC
2	0.056	64
4	0.1125	32
6	0.169	21.3
8	0.225	16
10	0.28	12.8
12	0.34	10.7
16	0.45	8
20	0.56	6.4
24	0.67	5.3
30	0.84	4.3
32	0.9	4

17/39

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Important: For wheat, barley, rye and rice, custom applicators must use a closed system for mixing and loading this product.

Propiconazole EC - Alone: Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add Propiconazole EC to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after Propiconazole EC has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

Propiconazole EC - Tank Mixes: Propiconazole EC is usually compatible with all tank mix partners listed on this label. To determine the physical compatibility of Propiconazole EC with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Propiconazole EC to the spray tank. Allow Propiconazole EC to completely disperse. Spray the mixture with the agitator running.

Do not apply this product in a tank mix with Cyprex fungicide or crop injury may occur.

If using Propiconazole EC in a tank mix, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations required by the label of the tank mix product. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. This product should not be tank mixed with any product that prohibits such mixing. Tank mixes or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Application Instructions

Propiconazole EC is most effective when applied and allowed to dry before a rainfall. For best results, use sufficient water volume to provide thorough coverage.

Aerial Application: Apply Propiconazole EC by aerial equipment in a minimum of 5 gallons of water per acre unless specified otherwise. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

Ground Application: Apply Propiconazole EC by ground equipment in a minimum of 15 gallons of water per acre unless otherwise specified.

Chemigation: This product may be applied through properly equipped chemigation systems for disease control in the labeled crops. Refer to crop specific use directions for application rates, timing and frequency of application. Do not apply Propiconazole EC by chemigation to other labeled crops except as specified in Dow AgroSciences supplemental labeling or product bulletins. When applying this product by chemigation, do not exceed labeled rates or apply more frequently than recommended for conventional application methods. Propiconazole EC, alone or in combination with other pesticides that are registered for application through irrigation systems, may be applied through irrigation systems. For chemigation application to labeled crops, apply 0.1 to 0.25 inches of water unless otherwise specified. Chemigation with excessive water may lead to a decrease in efficacy.

18/39

Precautions:

- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Note: Do not inject Propiconazole EC at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part Propiconazole EC. Propiconazole EC is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but should be replaced once a year. Do not use viton, Buna-N, Neoprene, or PVC seals.

Specific Equipment Requirements

1. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Center Pivot Irrigation Equipment

- Use only with drive systems that provide uniform water distribution.
- Do not use end guns when applying Propiconazole EC through center pivot systems because of non-uniform application.
- Determine size of area to be treated.
- Determine the time required to apply 1/4 to 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80 to 95% of the manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.

19/39

- Determine the amount of Propiconazole EC required to treat area.
- Add the required amount of Propiconazole EC and sufficient water to meet the injection time requirements of the solution tank.
- Make sure the system is fully charged with water before starting injection of Propiconazole EC. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until the solution of Propiconazole EC has cleared the sprinkler head.

Solid-Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinkler.
- Fill the injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval.
- Determine the amount of Propiconazole EC required to treat the area.
- Add the required amount of Propiconazole EC into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate the system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during the calibration.
- Inject Propiconazole EC at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until the solution of Propiconazole EC has cleared the last sprinkler head.

Banded Application: For banded applications, the treated area is the area covered by the band, not total cropland planted. The following formula can be used to calculate the amount of Propiconazole EC needed per acre of crop when banded applications are made.

$$\frac{\text{Band width in inches}}{\text{Row spacing in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Rotational Crops

Soybeans may be planted as a double crop following a cereal crop that has been treated with Propiconazole EC. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.

To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of an application of Propiconazole EC to the preceding crop unless the second crop appears on this label.

Uses

Almond

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 15 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	4 – 8	Apply at 5 to 10% bloom and 50 to 100% bloom. Under severe disease conditions, use the highest rate. The	• Do not apply more than 32 fl oz of Propiconazole EC (0.9 lb active ingredient) per acre per season.

20/39

		minimum re-treatment interval is 7 days.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 60 days of harvest. • Do not graze livestock in treated areas or cut treated cover crop for feed.
anthracnose (<i>Collectotrichum acutatum</i>)	8	Apply beginning at bud break on a 7- to 14-day interval.	

Berries

Including bingleberry, blackberry, black raspberry, black satin berry, blueberry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, currant, darrowberry, dewberry, dirksen thornless berry, elderberry, gooseberry, Himalayaberry, huckleberry, hullberry, juneberry, lavacaberry, lingonberry, loganberry, lowberry, lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, red raspberry, rossberry, salal, Shawnee blackberry, youngberry, and varieties and/or hybrids of these included in the berries crop grouping

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
cottonball (<i>Monilinia oxycocci</i>)	4 - 6	Make the first application at leaf bud break and repeat in 7 to 10 days. Make the third application at early bloom and repeat in 7 to 10 days. Apply in 20 to 50 gallons of water by ground or in 5 gallons of water by air. Under severe pressure, use the higher rate for control.	<ul style="list-style-type: none"> • Do not apply more than 30 fl oz of Propiconazole EC (0.84 lb active ingredient) per acre per season. • Preharvest Interval: Do not apply within 30 days of harvest.
mummyberry disease (<i>Monilinia vaccinicorymbosi</i>)	6	Make the first application beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, an additional application can be made at pink bud and repeated every 7 to 10 days through petal fall.	
leaf spot, stem canker (<i>Septoria albopunctata</i>) rust (<i>Pucciniastrum vaccinii</i>)		Apply when conditions favor disease development. Repeat applications on a 4-week spray interval.	
leaf and cane spot (<i>Septoria rubi</i>)		Apply as a delayed dormant spray after training in the spring. Repeat the application in late spring, at bud break, and once flowering has begun.	

powdery mildew (<i>Microspheera vaccinii</i>)		Apply at 5 to 10% bloom. Repeat the application at full bloom and on a 14-day interval while conditions are favorable for disease development.	
leaf spot (<i>Septoria</i> spp.)		Make first application any time prior to bloom and again after petal fall. If needed, repeat application just after harvest.	

Carrot

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
leaf blight (<i>Cercospora carotae</i>) powdery mildew (<i>Erysiphe polygoni</i>) <i>Alternaria dauci</i> (suppression only)	4	Apply when conditions favor disease development. Continue applications on a 7- to 10-day interval using the shorter interval when disease conditions are severe. If desired, a spreader-sticker may be used.	<ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 14 days of harvest.
	2	Apply with 0.75 lb active ingredient of chlorothalonil per acre. Begin applications when conditions favor disease development. Continue applications on a 7- to 10-day interval.	

Celery

Apply Propiconazole EC by either ground equipment in a minimum of 10 gallons of water per acre or aerial equipment in a minimum of 5 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
early blight (<i>Cercospora apii</i>) late blight (<i>Septoria apicola</i>)	4	Apply on a 7-day schedule. If desired, a spreader-sticker may be used.	<ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 14 days of harvest.

Cereals

22/39

Including barley, buckwheat, pearl millet, proso millet, oats, rice, rye, teosinte, triticale, wheat, wild rice and other commodities included in the cereal grains crop grouping. For rice and wild rice, see specific use directions elsewhere on the label.

Apply Propiconazole EC by either ground, aerial, or chemigation equipment.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
early season suppression of: glume blotch (<i>Stagonospora nordorum</i>) leaf blight (<i>Septoria tritici</i>) powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) tan spot (<i>Pyrenophora tritici-repentis</i>)	2 - 4	Apply in the spring. Follow up with a second application up to Feekes growth stage 10.5 for season long control. The minimum re-treatment interval is 14 days.	<ul style="list-style-type: none"> Do not apply more than 8 fl oz of Propiconazole EC (0.22 lb active ingredient) per acre per season. Do not apply more than 4 fl oz of Propiconazole EC (0.11 lb active ingredient) per acre per season if forage or hay will be harvested. Preharvest Interval: Do not apply within 30 days of harvest for forage, 40 days of harvest for grain and straw, and 45 days of harvest for hay.
control of leaf diseases: barley scald (<i>Rhynchosporium secalis</i>) barley stripe (<i>Pyrenophora graminea</i>) glume blotch (<i>Stagonospora nordorum</i>) helminthosporium leaf blight (<i>Drechslera tritici-repentis</i>) leaf blight (<i>Septoria tritici</i>) net blotch (<i>Pyrenophora teres</i>) powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) rust (<i>Puccinia</i> spp.) spot blotch (<i>Bipolaris sorokiniana</i>) tan spot (<i>Pyrenophora tritici-repentis</i>)	4	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Propiconazole EC is applied when the flag leaf is 50% to fully emerged. The minimum re-treatment interval is 14 days. Using an oil base adjuvant may improve the spray coverage and canopy penetration.	
foot rot (<i>Pseudocercospora</i> spp.)		Apply with half rates of EPA registered fungicides such as thiophanate-methyl. Apply at tillering but before elongation has occurred.	
fusarium head blight suppression		Apply at approximately 50% flowering. Adding a	

		penetrating type of adjuvant may increase fusarium head blight suppression.	
--	--	---	--

Corn (Field Corn, Field Corn Grown for Seed, Sweet Corn, and Popcorn)

Apply Propiconazole EC by either ground, aerial, or chemigation equipment.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
helminthosporium leaf blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>) northern corn leaf blight (<i>Setosphaeria turcica</i>) northern corn leaf spot (<i>Cochliobolus carbonum</i>) southern corn leaf blight (<i>Cochliobolus heterostrophus</i>)	2 – 4	Apply when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Apply the high rate under heavy disease pressure or if conditions are favorable for disease.	<p>Field corn, field corn grown for seed, popcorn</p> <ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Do not apply more than 8 fl oz of Propiconazole EC (0.22 lb active ingredient) per acre per season on field corn harvested for forage. <p>Sweet corn</p> <ul style="list-style-type: none"> Preharvest Interval: Do not apply within 14 days of harvest for ears or forage.
eye spot (<i>Aureobasidium zeae</i>) gray leaf spot (<i>Cercospora zeae-maydis</i>) rusts (<i>Puccinia</i> spp.)	4	Apply when disease first appears. If conditions favorable for disease persist, continue to apply on a 7- to 14-day schedule. For best disease control, early applications at initial disease onset perform better.	

Cranberry
(For use only in Oregon, Washington and Wisconsin)

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 20 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
cottonball (<i>Monilinia oxycocci</i>)	4 – 6	Make the first application at leaf bud break and repeat in 14 days. Make the third application at early bloom and repeat in 14 days. Under severe disease pressure, use the higher rate for control.	<ul style="list-style-type: none"> Do not apply more than 24 fl oz of Propiconazole EC (0.67 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 45 days of harvest.

Mint (Peppermint, Spearmint)

Apply Propiconazole EC by ground equipment in a minimum of 20 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
rust (<i>Puccinia menthae</i>)	4	Begin applications when the plants are 2 to 4 inches high or when conditions become favorable for disease development. Make a second application within 14 days of the first application.	<ul style="list-style-type: none"> Do not apply more than 8 fl oz of Propiconazole EC (0.22 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 30 days of harvest.

Nonbearing Citrus Fruits

Including calamondin, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin, sour orange, sweet orange, pummelo, satsume mandarin, tangerine, and cultivars and/or hybrids of these included in the citrus fruits crop grouping

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 15 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
greasy spot (<i>Mycosphaerella citri</i>)	6 – 8	Begin applications in June. Apply on a 30-day interval through August.	<ul style="list-style-type: none"> Do not apply more than 24 fl oz of Propiconazole EC (0.67 lb active ingredient) per acre per season. Do not apply to citrus that will bear harvestable fruit within 12 months.

Onions

Including dry bulb onions, dry bulb shallots, garlic, green eschalots, green onions, green shallots, Japanese bunching onions, leeks, scallions, spring onions, and cultivars and/or hybrids of these included in the bulb vegetables crop grouping

Apply Propiconazole EC by either ground equipment in a minimum of 15 gallons of water per acre or aerial equipment in a minimum of 5 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
botrytis leaf blight (<i>Botrytis squamosa</i>) (suppression only) purple blotch (<i>Alternaria pom</i>)	4 (Propiconazole EC only)	Begin applications when conditions favor disease development and continue on a 7- to 10-day interval. Use the higher rate and shorter interval when disease conditions are severe.	<ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 14 days of harvest on dry bulb onions. There is no preharvest interval for green onion types.
	2 – 4 (plus tank mix partner)	Apply with another fungicide registered for control of botrytis leaf blight or purple blotch. Begin applications when	

		<p>conditions favor disease development and continue on a 7-day interval or according to the tank mix partner label. Use the higher rate when disease conditions are severe. To achieve optimum disease control, use a wetting agent or spreader-sticker</p>	
--	--	--	--

Peanut

Apply Propiconazole EC by either ground, aerial, or chemigation equipment.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
early leaf spot (<i>Cercospora arachidicola</i>)	2.5 - 4	<p>Apply beginning 35 to 40 days after planting or at the first appearance of disease. Continue applications on a 10- to 14-day schedule. Under heavy disease pressure, use the higher rate. Propiconazole EC also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based upon environmental factors favorable for disease development.</p>	<ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 14 days of harvest with the low use rate of 4 fl oz per acre. Do not apply within 21 days of harvest with the high use rate of 8 fl oz per acre. Do not feed hay from treated fields to livestock if the high rate of 8 fl oz per acre was used.
late leaf spot (<i>Cercosporidium personatum</i>) rust (<i>Puccinia arachidis</i>) web blotch (<i>Phoma arachidicola</i>)	4		
southern stem rot (<i>Sclerotium rolfsii</i>)		<p>Apply to the crown and pegging zones of the plant using chemigation or directed ground application using one of the following schedules:</p> <p>A. Apply 4 fl oz per acre. Begin applications 45 days after planting or at the first appearance of disease. Repeat on a 14-day interval.</p> <p>B. Apply 8 fl oz per acre. Make the first application at pegging (approximately 60 days after planting) or at the first appearance of disease. Make the</p>	

20/39

		<p>second application 3 to 4 weeks later.</p> <p>When applying in irrigation water for southern stem rot control, use a minimum of 0.25 to 0.5 inches of water per acre. Use enough water so the fungicide penetrates the peanut canopy and reaches the crown of the plant where the disease is most active.</p> <p>When applying via irrigation or as a directed ground application, additional methods should be employed for leaf spot control.</p>	
--	--	--	--

Pecan

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 20 gallons of water per acre.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult <http://www.epa.gov/espp/usa-map.htm>. Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
pecan scab <i>(Cladosporium caryigenum)</i>	4 - 8	Apply 4 to 8 fl oz per acre on a 14-day schedule during bud break and pre-pollination sprays. Apply 6 to 8 fl oz per acre during nut formation and cover sprays. Use higher rates when disease pressure is heavier.	<ul style="list-style-type: none"> Do not apply more than 32 fl oz of Propiconazole EC (0.9 lb active ingredient) per acre per season. Do not apply after shuck split. Do not graze livestock in treated areas or cut treated cover crop for feed.
downy spot <i>(Mycosphaerella caryigena)</i> liver spot (<i>Gnomonia caryae pv pecanae</i>) powdery mildew <i>(Microsphaera penicillata)</i> vein spot (<i>Gnomornia nerviseda</i>) zonate leaf spot <i>(Cristulariella moricola)</i>		Apply 4 fl oz per acre with other products registered for pecans labeled for these mid- to late-season foliar diseases. Observe all directions, precautions, and limitations for the other products.	

Pineapple

(For use only in Hawaii)

Target Disease	Application Rate (fl oz/100 gal water)	Application Timing	Use Restrictions
butt rot (<i>Ceratocystis paradoxa</i>)	0.75 (1:17,000)	Treatments can be made in either a cold or hot water dip. Cold water dip: Immerse crowns to give thorough wetting, remove, and allow to drain. Hot water dip: Maintain water temperature at 125°F (52°C). Soak crown for 20 to 30 minutes, remove, and allow to drain.	<ul style="list-style-type: none"> Do not use treated crowns for food or feed. Do not graze while plant is growing. Do not graze tops until fruit is harvested. Dispose of used dip solution according to local, state, and federal regulations.

Rice

Apply Propiconazole EC by aerial equipment in 5 to 10 gallons of water per acre.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult <http://www.epa.gov/espp/usa-map.htm>. Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
aggregate sheath spot (<i>Rhizoctonia oryzae-sativa</i>) black sheath rot (<i>Gaeumannomyces graminis</i>) brown leaf spot (<i>Helminthosporium oryzae</i>) false smut (<i>Ustilaginoidea virens</i>) (suppression only) kernel smut (<i>Tilletia barclayana</i>) leaf smut (<i>Entyloma oryzae</i>) narrow brown leaf spot (<i>Cercospora oryzae</i>) sheath blight (<i>Rhizoctoria solani</i>) sheath spot (<i>Rhizoctonia oryzae</i>) stem rot (<i>Sclerotium oryzae</i>) (suppression)	6 - 10	Timing of application depends upon disease severity, disease complex and rice variety/growth stage. Consult local extension experts for local economic thresholds established for various rice varieties and diseases. Apply at first internode elongation (up to 2-inch panicle) using one of the following schedules: A. Apply 6 fl oz per acre and repeat at swollen boot. Make the second application 14 days later but before the boot splits and head emerges. Sheath blight is best controlled when the first application is applied at	<ul style="list-style-type: none"> Do not apply more than 12 fl oz of Propiconazole EC (0.34 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 35 days of harvest. Do not apply to stubble or ratoon crop rice. Do not use in rice fields where commercial farming of crayfish will be practiced. Do not drain water from treated rice fields into ponds used for commercial fish farming. Do not use water drained from treated fields to irrigate other crops.

only)		<p>disease appearance in the field. The first application is recommended when 5% or fewer of the tillers are infected.</p> <p>B. Apply 10 fl oz per acre. This rate is recommended if greater than 10% of the tillers are infected with sheath blight. If disease reappears, use another registered fungicide for the second application.</p> <p>Tank mix option: Apply 6 fl oz of Propiconazole EC in a tank mix with registered fungicides for control of rice diseases.</p>	
-------	--	---	--

**Wild Rice
(For use only in Minnesota)**

Apply Propiconazole EC by aerial equipment.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult <http://www.epa.gov/espp/usa-map.htm>. Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
brown spot (<i>Bipolaris</i> spp.) helminthosporium leaf blight (<i>Helminthosporium</i> spp.)	6 – 8	Apply 6 fl oz per acre at both booting and heading, or make a single application of 8 fl oz per acre at booting.	<ul style="list-style-type: none"> • Do not apply more than 12 fl oz of Propiconazole EC (0.34 lb active ingredient) per acre per season. • Preharvest Interval: Do not apply within 35 days of harvest. • Do not apply to stubble or ratoon crop rice. • Do not use in rice fields where commercial farming of crayfish will be practiced. • Do not drain water from treated rice fields into ponds used for commercial fish farming. • Do not use water drained from treated fields to irrigate other crops.

Sorghum

Apply Propiconazole EC by either ground equipment in a minimum of 15 gallons of water per acre or aerial equipment in a minimum of 10 gallons of water per acre.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
ergot (<i>Claviceps sorghi</i>)	3 – 4	Make the first application at or just prior to flowering. Repeat on a 5- to 7-day interval. Make up to 4 applications.	<ul style="list-style-type: none"> Do not apply more than 16 fl oz of Propiconazole EC (0.45 lb active ingredient) per acre per season. Do not apply more than 8 fl oz of Propiconazole EC (0.22 lb active ingredient) per acre per season on sorghum harvested for forage. Preharvest Interval: Do not apply within 30 days of harvest for forage and 21 days of harvest for grain and stover. Do not graze livestock or cut for green chop or silage within 30 days of application.

Stone Fruits

Including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune, sweet cherry, tart cherry, and cultivars and/or hybrids of these included in the stone fruits crop grouping

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 15 gallons of water per acre. Diseases on stone fruits are most effectively controlled by ground applications.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
brown rot blossom blight (<i>Monilinia</i> spp.)	4	Apply at early bloom stage. If disease pressure is low, a second application may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, make a second application from 75 to 100% bloom and a third application at petal fall.	<ul style="list-style-type: none"> Do not apply more than 20 fl oz of Propiconazole EC (0.56 lb active ingredient) per acre per season. Preharvest Interval: 0 days Applications of Propiconazole EC during bloom to stanley plums have occasionally caused fruit to be less oval in

30/39

fruit brown rot (<i>Monilinia</i> spp.)		Apply a maximum of 2 sprays as needed during the preharvest period up to the day of harvest (0 day PHI). If high inoculum and severe disease conditions persist, apply another registered fungicide after applying Propiconazole EC twice.	shape and smaller in size at harvest. To avoid this, do not apply to stanley plums earlier than 21 days before harvest.
cherry leaf spot (<i>Blumeriella jaapii</i>) powdery mildew (<i>Podosphaera</i> spp.) rust (<i>Tranzschelia discolor</i>)		See brown rot blossom blight. Make up to 2 additional applications on a 10- to 14-day interval from the end of petal fall to harvest.	

Sugarcane

Target Disease	Application Rate (fl oz/100 gal water)	Application Timing	Use Restrictions
pineapple disease (<i>Ceratocystic paradoxa</i>)	0.75 (1:17,000)	Apply to cut seed pieces using one of the following methods: Cold water dip: Immerse seed pieces to give thorough wetting, remove, and allow to drain. Hot water dip: Maintain water temperature at 125°F (52°C). Soak seed pieces for 20 to 30 minutes, remove, and allow to drain. Conveyor belt treatment: Treat seed pieces with a solution of Propiconazole EC and water using in-line directed spray sufficient to wet cut ends.	<ul style="list-style-type: none"> Do not use treated seed pieces for food or feed. Dispose of used dip solution according to local, state, and federal regulations.

Tree Nuts

Including almond, beechnut, Brazil nut, butternut, cashew, chestnut, chinaquapin, hickory, macadamia, pecan, walnut and other commodities included in the tree nuts crop grouping. For almond and pecan, see specific use directions elsewhere on the label.

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 15 gallons of water per acre. Diseases on tree nuts are most effectively controlled by ground applications.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
foliar diseases	4 – 8	Apply at first sign of disease. Repeat on a 7- to 14-day interval.	<ul style="list-style-type: none"> Do not apply more than 32 fl oz of Propiconazole EC (0.9 lb active ingredient) per acre per season. Preharvest Interval: Do not apply within 60 days of harvest. Do not graze livestock in treated areas or cut treated cover crop for feed.

Turfgrass Grown for Seed

(For use only in Idaho, Minnesota, Nebraska, Oregon, and Washington)

Apply Propiconazole EC by either ground or aerial equipment in a minimum of 10 gallons of water per acre. Propiconazole EC may also be applied by chemigation.

Target Disease	Application Rate (fl oz/acre)	Application Timing	Use Restrictions
ergot stem disease powdery mildew (<i>Erysiphe graminis</i>) rusts (<i>Puccinia</i> spp.) selenophoma stem eyespot (<i>Selenophoma</i> spp.)	4 – 8	Apply when powdery mildew infections, <i>Seleophoma</i> infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. Apply on a 14- to 21-day interval. To maximize control of severe rust pressure, apply 8 fl oz per acre and make applications at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin applications early in the growing season.	<ul style="list-style-type: none"> Do not apply more than 32 fl oz of Propiconazole EC (0.90 lb active ingredient) per acre per season. Do not feed hay cut within 20 days of the last application. Do not graze treated areas within 140 days of the last application.

Turfgrass and Ornamentals

General Information

Propiconazole EC fungicide is a systemic fungicide for use on turfgrass for the control of dollar spot (*Sclerotinia homeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Laetisaria fuciformis*), pink patch (*Limonomyces roseipellis*), rust (*Puccinia graminis*), powdery

32/39

mildew (*Erysiphe graminis*), stripe smut (*Ustilago striiformis* and *Urocystis agropyri*), summer patch (*Magnaporthe poae*), necrotic ring spot (*Leptosphaeria korrae*), spring dead spot (*Leptosphaeria korrae*, *Leptosphaeria narmari*, *Ophiosphaerella herpotricha*, *Gaeumannomyces graminis*), take-all patch (*Gaeumannomyces graminis*), leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), pink snowmold (*Microdochium nivale*), fusarium patch (*Fusarium nivale*), gray snowmold (*Typhula* spp.), yellow patch (*Rhizoctonia cerealis*), and zoysia patch (*Rhizoctonia solani*).

Propiconazole EC also controls numerous diseases on ornamentals and other landscape and nursery plantings. It controls powdery mildews, rusts, leaf spots, scabs, and blights. Refer to the appropriate section for specified diseases and plants.

General Use Precautions and Restrictions

Failure to follow the directions for use and precautions on this label may result in plant injury or poor disease control.

Chemigation: Do not apply this product through any type of irrigation system.

Do not use this product as a tree injection treatment.

Do not use Propiconazole EC in greenhouses.

Do not apply more than 5.8 fl oz per 1000 sq ft per year of Propiconazole EC.

Mixing Instructions

Fill the tank 1/2 to 3/4 full of water. Add the proper amount of Propiconazole EC and then add the rest of the water. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

If this product is mixed with other products, use the following mixing sequence:

- Always check the compatibility of the tank mix using a jar test with proportionate amounts of Propiconazole EC, other chemicals to be used, and the water before mixing in the spray tank.
- Provide sufficient jet or mechanical agitation during filling and application to keep the tank mix uniformly suspended.
- Fill the tank at least 1/2 full of clean water.
- Add wettable powders to the tank first, allowing them to completely suspend in the water before proceeding. This process can be hastened by premixing the product in water before adding to the tank.
- Add flowables or suspensions next.
- Add Propiconazole EC next.
- Add other emulsifiable concentrates last.
- Do not leave tank mixtures in the spray tank for prolonged periods without agitation. Mix and apply them the same day.

Tank Mixes: For broader spectrum disease control, Propiconazole EC may be mixed with other fungicides. Propiconazole EC is also compatible with numerous herbicides and insecticides. Always check compatibility using a jar test before tank mixing. Add Unite compatibility agent (3 pt per 100 gallons) to tank mixes that are incompatible. Follow the directions under Mixing Instructions for tank mixes. Observe all directions and most stringent precautions and limitations on the labels of all products used in the tank mix. Tank mixes or other applications of products referenced on this label are permitted only in those states where referenced products are registered.

Uses

Turfgrass and Dichondra

- Use Propiconazole EC in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to dry completely before irrigation.
- For control of foliar diseases, allow sprayed area to dry completely before irrigation.
- For control of soilborne diseases, Propiconazole EC can be watered in immediately after application.
- Under conditions that are optimum for high disease pressure, use the higher rate and shorter interval.
- For optimum turfgrass quality and disease control, use Propiconazole EC in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based upon data obtained with no additives.
- Before using any fungicide, proper diagnosis of the organism causing the disease is important. Using diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.
- Do not apply more than 5.8 fl oz per 1000 sq ft per year.
- Do not apply more than 1.79 lb active ingredient per acre per application.

Important: Bermudagrass can be sensitive to Propiconazole EC. Do not exceed 1.44 fl oz per 1000 sq ft every 30 days on any variety of bermudagrass. In Florida, do not apply Propiconazole EC to bermudagrass golf course greens when temperatures exceed 90°F.

Note: Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.

Rate conversion: When treating 1000 sq ft or less, values in rate tables may be difficult to measure accurately in fl oz. In such cases, it is suggested that users measure small quantities in a cylinder graduated in milliliters (1 fl oz = 30 milliliters).

Fluid Ounces to Milliliters Conversion Table				
0.18 fl oz	0.37 fl oz	0.72 fl oz	1.1 fl oz	1.44 fl oz
5.4 mL	10.8 mL	21.6 mL	33 mL	43.2 mL

Specific Diseases, Rates and Application Timing

Disease	Propiconazole EC		Application Interval	Specific Use Instructions
	(fl oz/1000 sq ft)	(fl oz/acre)		
Turfgrass				
anthracnose (<i>Colletotricum graminicola</i>)	0.37 - 0.73	16 - 32	14 - 28 days	Apply when conditions are favorable for disease development. When disease pressure is high, use a higher rate and shorter interval. For broad spectrum control, tank mix with a registered contact fungicide at the label rate. If disease is present, mix 0.73 fl oz of Propiconazole EC per 1000 sq ft with the label rate of the above-mentioned contact fungicides.
brown patch (<i>Rhizoctonia solani</i>)			14 - 21 days	Begin application in May or June before disease is present. Tank mix with the label rate of a contact fungicide registered for control of brown patch. Under conditions of high temperature and humidity, use a

34/39

				higher rate and shorter interval.
dollar spot (<i>Sclerotinia homeocarpa</i>)	0.18	8	7 days	Apply when conditions are favorable for disease development.
			14 days	Tank mix with low label rate of one of the following contact fungicides containing chlorothalonil.
	0.37	16	21 - 28 days	Tank mix with the low label rate of a contact fungicide containing chlorothalonil or Chipco 26019.
	0.37 - 0.73	16 - 32	14 - 28 days	If using the 0.37 - 0.73 fl oz/1000 sq ft rate without tank mixing, make no more than 3 consecutive applications for control of dollar spot before rotating to an alternate EPA-registered fungicide having a different mode of action.
fusarium patch (<i>Fusarium nivale</i>)	0.73 - 1.45		fall - early spring	Apply when conditions are favorable for disease development.
gray leafspot (<i>Pyricularia grisea</i>)	0.37 - 0.73	16 - 32	14 days	Apply when conditions are favorable for disease development. If using the 0.37 fl oz/1000 sq ft rate, tank mix with a registered contact fungicide at the label rate.
melting out, leaf spot (<i>Bipolaris</i> spp.), (<i>Drechslera</i> spp.)				Under light to moderate pressure, apply Propiconazole EC to reduce the severity of leaf spot and melting out caused by <i>Helminthosporium</i> -type pathogens. For broad spectrum disease control, tank mix the 0.37 fl oz/1000 sq ft rate with a registered contact fungicide at the label rate.
necrotic ring spot (<i>Leptosphaeria korrae</i>)	1.45	63	fall or spring	Apply in fall and/or the early spring depending upon local recommendations.
powdery mildew (<i>Erysiphe graminis</i>) rust (<i>Puccinia graminis</i>)	0.37 - 0.73	16 - 32	14 - 28 days	Apply when conditions are favorable for disease development. If disease is present, use 0.73 fl oz of Propiconazole EC per 1000 sq ft.
red thread (<i>Laetisaria fuciformis</i>) pink patch (<i>Limonomyces roseipellis</i>)	0.37	32	14 - 21 days	Apply when conditions are favorable for disease development.
snow mold gray (<i>Typhula</i> spp.)	0.73 - 1.45	32 - 63	late fall	Make 1 application in the late fall before snow cover. Do not apply on top of snow. For optimum

pink (<i>Microdochium nivale</i>)				disease control, 0.73 – 1.45 fl oz/1000 sq ft should be tank mixed with either PCNB or chlorothalonil at label rates.
spring dead spot (<i>Leptosphaeria korrae</i>), <i>Leptosphaeria narmari</i> (<i>Ophiosphaerella herpotricha</i>), (<i>Gaeumannomyces graminis</i>)	1.45	63	30 days	Make 1 to 3 applications. If a single application is made, apply in September or October. For multiple applications, begin sprays in August.
stripe smut (<i>Ustilago striiformis</i>), (<i>Urocystis agropyri</i>)	0.37 - 0.73	16 - 32	fall or spring	Apply once in the fall after turfgrass becomes dormant or in the early spring before turfgrass starts to grow.
summer patch poa patch (<i>Magnaporthe poae</i>)	0.73 1.45	32 63	14 days 28 days	Apply Propiconazole EC beginning in April. Use the 0.73 fl oz/1000 sq ft rate on a 14-day schedule or the 1.45 fl oz/1000 sq ft rate on a 28-day schedule.
take-all patch (<i>Gaeumannomyces graminis</i>)	0.73 – 1.45	32 - 63	spring and fall	Apply Propiconazole EC to reduce the severity of take-all patch. Make 1 or 2 fall applications in September and October or when night temperatures drop below 55°F, and 1 or 2 applications in April and May, depending upon local recommendations.
yellow patch (<i>Rhizoctonia cerealis</i>)	1.1 – 1.45	48 – 63	late fall	Make 1 application in the late fall before snow cover. Do not apply on top of snow. If using the 1.1 fl oz/1000 sq ft rate, tank mix with a registered contact fungicide at the label rate.
zoysia patch, large patch of zoysia (<i>Rhizoctonia solani</i>)	1.1 – 1.45	48 – 63	early fall	Make 1 application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing for your area.
Dichondra				
dichondra rust (<i>Puccinia dichondrae</i>)	0.73	32	14 - 21 days	Apply when conditions are favorable for disease development.

Establishment of Cool Season Turfgrass

Propiconazole EC provides control of many turfgrass diseases; its primary use is as a fungicide for use against the diseases listed on this label. As an additional benefit, Propiconazole EC improves the rate of establishment when it is applied to cool season turfgrass seedlings or sod.

New Seedlings: Apply 0.37 fl oz per 1000 sq ft at the 2 to 3 leaf stage of growth for faster root development and top growth.

Sod: Apply 0.37 fl oz per 1000 sq ft 2 to 6 weeks before cutting for increased sod knitting and faster establishment after laying.

Ornamentals and Non-Bearing Fruit and Nut Trees in Nurseries (Field) and Landscape Plantings

- Use Propiconazole EC in a preventative disease control program. To determine the use directions for controlling a disease on an ornamental plant species, select the plant species in Table 1. The number(s) in parentheses following the listed plant species refers to the disease(s) controlled in Table 2. Find the disease in Table 2. The letter in brackets following the disease refers to the application regime in Table 3.
- Allow spray to dry before applying overhead irrigation.
- Optimum benefit of Propiconazole EC is obtained when used in conjunction with sound disease management practices.

Propiconazole EC may be used at rates of 0.73 to 8.7 fl oz per 100 gallons of water for disease control in ornamentals (see Tables 1, 2 and 3). For general disease control in landscape plantings, apply 2.2 to 3 fl oz per 100 gallons of water every 21 days. For best control, begin applications before disease development.

Note: For outdoor uses, up to 2 gallons of Propiconazole EC may be applied per acre per crop per year.

Plant tolerances to Propiconazole EC have been found acceptable for the specific genus and species of plants listed under the Directions for Use. Other plant species may be sensitive to Propiconazole EC and diseases other than those listed may not be controlled. Before using Propiconazole EC on plants or for diseases that are not listed in the Directions for Use, first test Propiconazole EC on a small scale basis. Do not apply Propiconazole EC to African violets, begonias, Boston fern, or geraniums. Apply the recommended rates for a particular disease, i.e., rust, powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use.

Table 1. Ornamental Plant Species

Number in parentheses (-) refer to diseases controlled in Table 2.

Herbaceous Ornamentals	Woody Ornamentals	Non-Bearing Fruits and Nut Trees (Nursery and Landscape Plantings)
calendula (4a)	amelanchier (4d)	apple (3q, 4d, 5a)
camation (5f)	ash (4c)	bartlett pear (3q, 4c, 5a)
chrysanthemum (2a)	azalea (2c, 4b)	cherry (2b, 3d)
delphinium (4a)	bayberry (3n)	citrus (3m)
English ivy (3e)	camellia (3e)	nectarine (2b)
gompfrena (3a)	cotoneaster (3i)	peach (2b)
impatiens (3a, 3b, 4a)	crabapple (3c, 3g, 4c, 5a)	pecan (3b, 3c, 3f, 3l, 3n, 4e)
iris (5d)	crape myrtle (4a)	plum (2b)
marigold (3a)	dogwood (3h, 4c)	walnut (3j)
monarda (4c)	douglas fir (5b)	
phlox (4c)	elm (4c)	
snapdragon (5d)	euonymus (3e, 4c)	
sweet william (3k) (<i>Dianthus barbatus</i>)	hawthorn (5a)	
zinnia (4c)	holly (3r)	
	juniper (1a)	
	lilac (4c)	

	linden (3e, 3b, 4b) magnolia (3e, 4b) maple (3e, 4f) oaks (3p) pines (1b, 1c) poplars (5b) pyracantha (3o) red tip photinia (3i) raphiolepis (3e, 3i) rhododendron (2c, 3n) roses (3g, 4e, 5c) (outdoor use only) shasta fir (5e) sweetgum (3b, 3c, 3n) sycamore (3e) tulip tree (3e, 4a) wax myrtle (3n)	
--	---	--

Table 2. Plant Diseases

Letters in brackets [-] refer to application regimes in Table 3.

1. Conifer Blights
 - a. *Phomopsis juniperovora* (phomopsis blight) [B]
 - b. *Sirococcus strobolinus* (tip blight) [D]
 - c. *Sphaeropsis sapinea* (diplodia tip blight) [B]
2. Flower Blight
 - a. *Ascochyta chrysanthemi* (ray blight) [C]
 - b. *Monilinia* spp. [A]
 - c. *Ovulinia* spp. [B]
3. Leaf Blights/Spots
 - a. *Alternaria* spp. [B]
 - b. *Cercospora* spp. (brown leaf spot) [C]
 - c. *Cladosporium* spp. (scab) [C]
 - d. *Coccomyces hiemalis* [A]
 - e. *Collectrochum* spp. [B]
 - f. *Cristulariella* spp. (zonate leaf spot) [C]
 - g. *Diplocarpon rosae* (blackspot) [B]
 - h. *Discula* spp. (anthracnose) [A]
 - i. *Fabraea maculata* (syn. *Entomosporium maculata*) [B]
 - j. *Gnomonia leptostylia* (anthracnose) [C]
 - k. *Heterosporium echinulatum* [B]
 - l. *Mycosphaerella caryigena* (downy spot) [C]
 - m. *Mycosphaerella fructicola* (greasy spot) [E]
 - n. *Septoria* spp. (leaf scorch) [C]
 - o. *Spilocaea pyracanthae* [B]
 - p. *Tubakia dryina* [D]
 - q. *Venturia inaequalis* (scab) [A]
 - r. *Rhizoctonia web blight* [B]
4. Powdery Mildew
 - a. *Erysiphe* spp. [B]

- b. *Microsphaera* spp. [C]
- c. *Oidium* spp. [B]
- d. *Podosphaera* spp. [B]
- e. *Sphaerotheca pannosa* [B]
- f. *Phyllactinia* spp. [B]

5. Rust

- a. *Gymnosporangium juniperi-virginianae* [A]
- b. *Melampsora occidentalis* [D]
- c. *Phragmidium* spp. [B]
- d. *Puccinia* spp. [B]
- e. *Pucciniastrum goeppertianum* [D]
- f. *Uromyces dianthi* [B]

Table 3. Application Regimes

- [A] Mix 0.73 to 1.45 fl oz of Propiconazole EC in 100 gallons of water and apply as a full coverage spray to the point of drip. Make repeat applications every 14 to 21 days during the period of primary infection. If disease is present, tank mix with an EPA-registered contact fungicide. For **flower blight**, apply Propiconazole EC when there is 5 to 10% bloom and again at 70 to 100% bloom. For **dogwoods**, apply the 0.73 to 1.45 fl oz rate every 14 days, or apply 3 fl oz of Propiconazole EC every 28 days.
- [B] Mix 1.8 to 3 fl oz of Propiconazole EC in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply as needed, beginning when conditions are favorable for disease development. For **blackspot**, apply in tank mix with a registered contact fungicide labeled for blackspot. For **calendula**, apply every 30 days. For **diplodia tip blight**, make a total of 3 applications every 14 days prior to the major period of infection. For **juniper phomopsis blight**, make an initial application as soon as junipers start to grow and repeat the application every 14 to 21 days during the period of active growth.
- [C] Mix 3 to 4.5 fl oz of Propiconazole EC in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 30 days, beginning when conditions are favorable for disease development. For **pecans**, apply the 4.5 fl oz rate. Beginning at bud break, make a total of 3 applications 14 days apart. For **walnuts**, apply 3 fl oz every 14 to 21 days. For **ray blight**, apply 4.5 fl oz every 7 days or 7.5 fl oz every 14 days. For **impatiens**, **bayberry**, **linden**, **magnolia**, **sweetgum** and **wax myrtle**, the maximum use rate is 8 fl oz.
- [D] Mix 6 fl oz of Propiconazole EC in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 14 to 28 days beginning when conditions are favorable for disease development. For **douglas fir needle rust**, apply once in May. For **tip blight**, make an initial application in mid- to late winter, and 3 additional applications at 2-month intervals.
- [E] Mix 7.3 to 8.7 fl oz of Propiconazole EC in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply within the June to August time period.

Note: To avoid possible illegal residues, do not apply to apple, bartlett pear, cherry, citrus, nectarine, peach, pecan, plum, or walnut trees that will bear harvestable fruit within 12 months.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election; one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

EPA accepted ___/___/___