

62719-346

9/30/2009

1 of 11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Joseph W. Burley
Regulatory Manager
Sipcam Agro USA, Inc.
2520 Meridian Pkwy., Suite 525
Durham, NC 27713

SEP 30 2009

SUBJECT: Applications for Pesticide Notification – Request to Add CPDA Language Appropriate to Approved Adjuvants per PR Notices 98-10, and Revise Storage & Disposal Language per PR Notice 2007-4
Echo® 720 with Eminent 125SL (Co-Pack)
EPA Reg. Nos. 60063-7 and 60063-12
Echo® 720 with Muscle® 3.6F (Co-Pack)
EPA Reg. Nos. 60063-7 and 60063-29
EchoPropiMax Fungicide (Echo® 720 / PropiMax™ EC) (Co-Pack)
EPA Reg. Nos. 60063-7 and 62719-346
Applications Dated August 5, 2009

Dear Mr. Burley:

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notices (PRNs) 98-10 and 2007-4 for the above products. The Registration Division (RD) has conducted a review of these requests for their applicability under PRNs 98-10 and 2007-4, and finds that that the actions requested fall within the scope of these PRNs. The labels submitted with the applications have been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249, or Terri Stowe of my staff at 703-305-6117.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20911



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 60063-7; 62719-346	2. EPA Product Manager Tony Kish; Mary Waller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product Name EchoPropiMax Fungicide Co-Pack	PM# 22; 21	
5. Name and Address of Applicant (Include ZIP Code) Sipcam Agro USA, Inc. 2520 Meridian Parkway, Suite 525 Durham, NC 27713	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Check if this is a new address

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION SEP 30 2009
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146 and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product except as certified in the paragraph below. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. [See attached page for further certification statement.]

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Unit Packaging wgt. No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container

5. Location of Label Directions
 On Label
 On Labeling accompanying product

6. Manner in Which Label is Affixed to Product
 Lithograph Other
 Paper glued
 Stenciled

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Joseph W. Burley	Title Regulatory Manager	Telephone No. (Include Area Code) (919) 226 1297
---------------------------------	------------------------------------	--

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Regulatory Manager	6. Date Application Received (Stamped)
4. Typed Name Joseph W. Burley	5. Date 08/05/2009	

Section II Explanation (continued):

Notification of addition of EPA approved language regarding certified adjuvants per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula except as certified in the paragraph above. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

EchoPropiMax Fungicide Co-Pack
For control of certain diseases in peanuts

50911

NOTIFICATION

ECHO® 720
EPA Reg. No. 60063-7

Active Ingredient: Chlorothalonil
(tetrachloroisophthalonitrile) 54.0%
Other Ingredients: 46.0%
Total: 100.0%

Contains 6.0 Pounds Chlorothalonil Per Gallon

Net Contents: 160 fl.oz. U.S. Standard Measure

ECHO is a registered trademark of Sipcam Agro USA, Inc.

PropiMax* EC SEP 30 2009
EPA Reg. No. 62719-346

Active Ingredient: Propiconazole
(1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole) 41.8%
Inert Ingredients 58.2%
Total 100.0%

Contains petroleum distillates.

Net Contents 20 fl.oz. U.S. Standard Measure

*Propimax is a trademark of Dow AgroSciences


Keep Out of Reach of Children
WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have affected person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center (human health) (800) 345-4735 ASPCA (animal health)
<p>NOTES TO PHYSICIAN: For the Echo 720 (chlorothalonil) portion of this product- Probable mucosal damage may contraindicate the use of gastric lavage. Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids. For the PropiMax EC (propiconazole) portion of this product- There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy. This product may pose an aspiration pneumonia hazard. Contains petroleum distillate.</p>	

EPA Est. No. _____

Manufactured for
Sipcam Agro USA, Inc.
2520 Meridian Parkway, Suite 525
Durham, NC 27713



Front Panel

60011

EchoPropiMax Fungicide Co-Pack
For control of certain diseases in peanuts

Echo 720 Precautionary Statements

Hazards to Humans and Domestic Animals

May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Do not breathe spray mist.

PropiMax EC Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury. Harmful If Swallowed, Inhaled, Or Absorbed Through Skin. Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.

EchoPropiMax Fungicide Co-Pack Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to these products are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear
- A NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

70911

EchoPropiMax Fungicide Co-Pack
For control of certain diseases in peanuts

EchoPropiMax Fungicide Co-Pack Environmental Hazards

These products are toxic to aquatic invertebrates, fish and wildlife. DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwater or rinsate.

Chlorothalonil can contaminate surface water through spray drift. DO NOT apply when weather conditions favor drift from treated areas. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Chlorothalonil degradates are known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

EchoPropiMax Fungicide Co-Pack Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

EchoPropiMax Fungicide Co-Pack Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

DO NOT apply this product in a way that will contact workers or other persons, or pets, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

EchoPropiMax Fungicide Co-pack Agricultural Use Requirements

Use these products only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

This co-pack includes two separate products which are designed to be used and applied together to the intended site. Agricultural use requirements subject to the Worker Protection Standard are hereby stated in accordance with the most protective of the two product labels:

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Special Eye Irritation Provisions: The Echo 720 component of this co-pack product is a severe eye irritant. Although the REI for the co-pack product expires after 24 hours, for the next 6 days entry is permitted only when the following safety measures are provided:

80811

EchoPropiMax Fungicide Co-Pack For control of certain diseases in peanuts

At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.

Workers must be informed, in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes;
- that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes;
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water; and
- how to operate the eyeflush container.

EchoPropiMax Fungicide Co-Pack General Information

Use EchoPropiMax Fungicide Co-Pack only for the control of certain diseases in peanuts as specified on this label.

Failure to follow directions and precautions on this label may result in crop injury, poor disease control, or illegal residues.

This product must not be applied within 150 feet (for aerial and air-blast applications), or 25 feet (for ground applications) from marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

Application Information for Drift Reduction

DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable conditions (see Wind, Temperature).

- Volume- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles- Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation- Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle type- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the

9 of 11

EchoPropiMax Fungicide Co-Pack

For control of certain diseases in peanuts

aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, small drops, etc.).

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Aerial Application:


BOOM LENGTH

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

For those crops where aerial applications are indicated, apply in a **minimum of 5 gallons of water per acre**, unless specified otherwise. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

Ground Application: Apply by ground equipment in a **minimum of 15 gals of water per acre**.

Chemigation: Do not apply this product through any type of irrigation system.



10 of 11

EchoPropiMax Fungicide Co-Pack
For control of certain diseases in peanuts

EchoPropiMax Fungicide Co-Pack Mixing Instructions

Note: Slowly invert container several time to assure uniform mixture.

Thoroughly clean spray equipment before using this product.

The contents of this container are designed to be applied to ten (10) acres of growing peanut crop.

Prepare no more spray mixture than is required for the immediate operation. Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank, and begin agitation. Pour the entire contents both chambers of this container into the tank at the same time, maintaining constant agitation, to form a suspension in water. Slowly add the remaining required amount of water to the tank. Continue agitating the spray suspension before and during application.

Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to a previously treated area of the same crop.

- If using in a tank mixture of this product with other materials or pesticides registered for peanuts, observe all directions for use, use rates, dilution ratios, precautions, and limitations required by the label of the tank mix product. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be tank mixed with any product that prohibits such mixing. When an adjuvant is to be used with this product, Sipcarn Agro USA recommends the use of a Chemical Producers and Distributors Association (CDPA) certified adjuvant.

EchoPropiMax Fungicide Co-Pack Directions For Use on Peanuts

Apply the entire contents of this co-pack container together for the control of the following diseases of peanuts caused by fungi:

- Early leafspot (*Cercospora arachidicola*)
- Late leafspot (*Cercosporidium personatum*)
- Web blotch (*Phoma arachidis*)
- Rust (*Puccinia arachidicola*)

Apply when conditions favor disease, generally when leaf wetness first occurs, or 30 to 40 days after planting. Repeat applications of this product at 14 day intervals if conditions remain favorable for disease. Do not make more than eight (8) applications of this product per growing season. Do not apply within 14 days before harvest (digging) of peanuts. Do not allow livestock to graze in treated areas. Do not feed hay or threshings from treated fields to livestock.

Rotational Crops

To avoid possible illegal residues, do not plant any other crop within 105 days after an application of this product to peanuts, unless the rotational crop to be planted is one of the following: cereals (wheat, barley, rye, oats, rice, and wild rice), celery, corn, nonbearing citrus, stone fruit (apricots, nectarines, peaches, plums, and prunes), peanuts, pecans, or grasses grown for seed.

