

62719-371

05/24/2007

ACCEPTED

MAY 24 2007

1/2

Supplemental Labeling



Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 62719-371

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

Due to Oncogenicity

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FulTime®

EPA Reg. No. 62719-371

Weed Control in Sweet Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for FulTime® herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of FulTime according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

FulTime Applied Alone

Used alone, FulTime® herbicide may be applied to sweet corn using rates recommended in the product label for FulTime and application methods and timing listed below.

Carriers and Spray Volume: Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. Refer to product label for FulTime for details.

Application Timing and Methods: For the optimum period of effective weed control during the time most critical to corn production, preplant applications of FulTime should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Refer to product label for FulTime for details.

Note: Do not apply FulTime to sweet corn as an early postemergence application.

- **Early Preplant:** On medium and fine textured soils (see Table 1 in main label), FulTime may be applied up to 30 days prior to planting.
- **Preplant Incorporation:** FulTime and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting.
- **Preemergence Surface:** FulTime and certain tank mixes may be applied to the soil surface as a broadcast or banded application.

- 2/10
- **Postplant-Preemergence:** FulTime may be applied immediately after planting but prior to corn emergence.
 - **Banding Preemergence:** FulTime may be applied in a 10 to 14 inch band after corn planting but prior to corn emergence.

Sprinkler Irrigation: Do not apply FulTime by sprinkler irrigation. Use a sprinkler system only to incorporate FulTime after application. Refer to product label for FulTime for details.

Cultivation: Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. Refer to product label for FulTime for details.

Soil Texture and Organic Matter: The use rate of FulTime is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Refer to Table 1 in main product label for soil texture classifications.

Use Rates for Conventional Tillage Systems: Refer to table 2 in main label for application rates and detailed information for conventional systems.

Use Rates for Reduced Tillage Systems: Refer to table 3 in main label for application rates and detailed information for reduced tillage systems.

Weeds Controlled: Refer to table 4 in main label for details.

FulTime Applied in Tank Mix Combination

Additional weeds may be controlled with tank mixes of FulTime and other products labeled for use on sweet corn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied using application methods and timing listed above unless otherwise specified in the tank mix product label. Refer to the product label for FulTime for details on carriers, mixing instructions and testing for compatibility of tank mixtures.

®Trademark of Dow AgroSciences LLC

R140-001

EPA Accepted: / /

Initial printing.

62719-371

05/24/2007

111

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE
Due to Oncogenicity
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FulTime®

EPA Reg. No. 62719-371

Rotational Crop Restrictions After Final Application

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for FulTime® herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of FulTime according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Rotational Crop Restrictions

When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

Crop	Interval
Corn (1)	Anytime - 0 Months After Application
Sorghum, Soybean (2)	Spring Following Application
Alfalfa, Barley, Dry beans (3), Lupin (4), Millet, pearl or proso, Oats, Pea (6), Potatoes, Rye, Sugar beets Sunflower, Tobacco (7), Triticale, Wheat, Wild rice	15 Months After Application (5)

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements or crop descriptions below.

- (1) If crop treated with FulTime is lost, corn may be replanted immediately. Do not make a second application of FulTime. Do not apply FulTime after June 10, unless only corn will be planted the following year.
- (2) Due to the risk of atrazine carryover, injury to soybeans may occur the year following corn when planted in north central and northwest Iowa, south central and southwest Minnesota, northern Nebraska and southeast South Dakota on soils having a calcareous surface layer and relatively high pH.
- (3) Dry beans includes: adzuki, kidney, lima, navy, pinto
- (4) Lupin includes: grain, white, white sweet
- (5) Approved rotation crops list does not include any species of succulent beans and peas.
- (6) Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern
- (7) Because of atrazine carryover, injury may occur to tobacco.

®Trademark of Dow AgroSciences LLC
R140-002
EPA Accepted: ___/___/___
Initial printing.

ACCEPTED
MAY 24 2007
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 62719-371