

62719-375

2/29/2008

1 of 28



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Mr. Ray S. Brinkmeyer  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

FEB 29 2008

Dear Mr. Brinkmeyer:

Subject: Quintec  
EPA Reg. No. 62719-375  
Your Submissions of January 15, 2008

The amendments to the supplemental labels for use on strawberries, melons and head and leaf lettuce, referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, are acceptable. A stamped copy of each supplemental label is enclosed for your records.

Submit one copy of the final printed labeling for each crop before you release the product for shipment.

Sincerely yours,

A handwritten signature in black ink, appearing to read "Mary L. Waller".

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505P)

Enclosure

7505P:CGrable:cg:2/27/08

(Base label for rigid containers 5 gal or less):

# Quintec®


## Fungicide

**A protectant fungicide for the control of powdery mildew in pepper, melons, strawberry, lettuce, grapes, hops, and cherry.**

<b>Group</b>	<b>13</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

**Active Ingredient:**

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy) quinoline.....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%



Contains 2.08 lb of active ingredient per gallon.

**Keep Out of Reach of Children**

## CAUTION

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**Harmful If Swallowed • Harmful If Absorbed Through Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.**

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center

or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

40928

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**Net Contents \_\_\_\_\_**

50f28

(Base label for refillable rigid containers larger than 5 gal):

# Quintec®

## Fungicide

**A protectant fungicide for the control of powdery mildew in pepper, melons, strawberry, lettuce, grapes, hops, and cherry.**

<b>Group</b>	<b>13</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy) quinoline .....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%

Contains 2.08 lb of active ingredient per gallon.

### Keep Out of Reach of Children

## CAUTION

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Harmful If Swallowed • Harmful If Absorbed Through Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.**

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center

or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC  
**Produced for**  
**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**Net Contents** \_\_\_\_\_

80f28

(Base label for nonrefillable rigid containers larger than 5 gal):

# Quintec®

## Fungicide

**A protectant fungicide for the control of powdery mildew in pepper, melons, strawberry, lettuce, grapes, hops, and cherry.**

<b>Group</b>	<b>13</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy) quinoline .....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%

Contains 2.08 lb of active ingredient per gallon.

**Keep Out of Reach of Children**

## CAUTION

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Harmful If Swallowed • Harmful If Absorbed Through Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.**

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center



90912

or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

100f28

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.  
If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

**Produced for**  
**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**Net Contents** \_\_\_\_\_

(Label booklet cover)

# Quintec®

## Fungicide

**A protectant fungicide for the control of powdery mildew in pepper, melons, strawberry, lettuce, grapes, hops, and cherry.**

<b>Group</b>	<b>13</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

**Active Ingredient:**

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)	
quinoline .....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%

Contains 2.08 lb of active ingredient per gallon.

### Keep Out of Reach of Children

## CAUTION

**Agricultural Use Requirements**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering life or property involving this product, call collect (517) 636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**Net Contents \_\_\_\_\_**

12 of 19

(Page 1 through end)

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION****Harmful If Swallowed • Harmful If Absorbed Through Skin****Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.****Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

13 of 20

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable containers 5 gallons or less:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### Refillable containers 5 gallons or larger:

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the

14 of 28

container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable containers 5 gallons or larger:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### General Use Directions

Quintec® fungicide is a protectant fungicide for the control of powdery mildew diseases. Quintec does not control existing or latent powdery mildew infections and must be applied on a protectant schedule (latency is the period between infection of plant tissues and the development of visible symptoms). Quintec is compatible with most commonly recommended insecticides, miticides and fungicides.

### General Use Precautions and Restrictions

- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- Do not aerially apply this product.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Rotational Crop Restrictions:** Crops grown for food or feed without registered uses for quinoxyfen may be replanted 12 months after the last application of quinoxyfen.
- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

### Disease Resistance Management

The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is in mode of action Group 13. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a mode of action different from Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow

15 of 28

AgroSciences representative or by calling 800-253-3033 or over the internet at [www.dowagro.com](http://www.dowagro.com).

### **Mixing Instructions**

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and Quintec or other flowable (aqueous suspension) products to the spray tank. Then add the recommended rate of emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

### **Oil Sprays**

Oil sprays may be integrated as part of a powdery mildew management program. Consult local university extension for specific recommendations.

### **Use of Surfactants**

Quintec contains sufficient surfactant to ensure mixability and plant coverage. The use of additional surfactant is not required.

### **Tank Mixing**

Quintec may be applied in tank mix combination with labeled rates of other pesticides, including fungicides, insecticides, or other chemicals used in the culture of the labeled crop provided: (1) the tank mix product is labeled for the timing and method of application for the intended use; and (2) tank mixing with Quintec or other pesticides is not prohibited by the label of the tank mix product.

### **Tank Mixing Precautions:**

- Read carefully and follow all applicable use directions including application rates, use precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates or maximum use rates if products with the same active ingredient are applied together
- For tank mix partners packaged in water soluble packaging, do not use with products containing boron or mix in equipment previously used to apply boron-containing mixtures unless the tank and spray equipment have been adequately cleaned.
- Always perform a (jar) test to ensure compatibility of products used in tank mixtures.

**Tank Mix Compatibility Testing:** A jar test is recommended prior to tank mixing to ensure compatibility of Quintec and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

### **Concentrate and Dilute Sprays**

Concentrate sprays are generally applied in a spray volume of 100 gallons or less per acre to provide uniform coverage, but not to the point of runoff. Dilute sprays are applied in sufficient spray volume to completely wet the target plant to the point of runoff. Gallonage of dilute sprays will vary depending on the volume, density and spacing of the crop canopy and will generally increase in proportion to crop development during the growing season.

**Rainfast Requirement:** When applied alone, Quintec is rainfast 1 hour after application.

### **Application Guidelines**

#### **Application Timing**

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if

16 of 28

multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. Maximum label rates and short application intervals are recommended for severe or threatening disease conditions.

### Cherry, Grape, Hops

Crop	Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre) <sup>1</sup>
cherry	powdery mildew ( <i>Podosphaera clandestina</i> )	10 - 14	7
grape	powdery mildew ( <i>Uncinula necator</i> )	7	1 - 2
		14	3 - 4
		21	5 - 6.6
hops	powdery mildew ( <i>Sphaerotheca macularis</i> )	14	4 - 8.2

<sup>1</sup>When applied as a dilute spray, maintain the per acre use rate.

### Specific Use Restrictions:

#### Cherry

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not make more than five applications of Quintec per calendar year.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or more than 35 fl oz of Quintec per acre per calendar year.
- Do not make more than five applications of Quintec per calendar year.
- **Minimum Spray Interval:** 7 days.

#### Grape

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- Do not make more than five applications of Quintec per calendar year.
- **Maximum Application Rate:** Do not apply more than 6.6 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- Do not make more than five applications of Quintec per calendar year.
- **Minimum Spray Interval:** 5 days for 1 to 2 fl oz per acre rate; 7 days for 3 to 4 fl oz per acre rate; 14 days for 5 to 6.6 fl oz per acre rate.

#### Hops

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- Do not make more than four applications of Quintec per calendar year.
- **Maximum Application Rate:** Do not apply more than 8.2 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per calendar year.
- **Minimum Spray Interval:** 7 days.

### Head and Leaf Lettuce

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Erysiphe cichoracearum</i> )	10 - 14	4 - 6



17 of 28

**Specific Use Precautions and Restrictions:**

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop.
- Do not make sequential applications of Quintec on lettuce. After each application of Quintec, alternate with a product that is effective on powdery mildew and has a different mode of action than Quintec. Apply the alternate product within 14 days of the application of Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

**Melons [Including Citron Melon, Muskmelon (Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon, True Cantaloupe), Watermelon]**

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Sphaerotheca fuliginea</i> )	10 - 14	4 - 6

**Specific Use Precautions and Restrictions:**

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec.
- Do not make more than two consecutive applications of Quintec before rotating to a product that is effective on powdery mildew and has a different mode of action than Quintec. In either case, apply the alternate product within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

**Pepper (Bell Pepper, Chili Pepper, Cooking Pepper, Non-Bell Pepper, Pimento, Sweet Pepper)**

Apply uniformly in a minimum spray volume of 30 gallons per acre that assures thorough coverage of the crop. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Leveillula taurica</i> , <i>Oidiopsis taurica</i> )	10 - 14	4 - 6

**Specific Use Precautions and Restrictions:**

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec.

18 of 28

- It is recommended to alternate Quintec with a product with a different mode of action after each application of Quintec. Apply the alternate product within 14 days of applying Quintec.
- When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different mode of action than Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

### Strawberry

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Sphaerotheca macularis</i> )	10 - 14	4 - 6

### Specific Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec.
- Do not make more than two consecutive applications of Quintec before rotating to a product that is effective on powdery mildew and has a different mode of action than Quintec. In either case, apply the alternate product within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.

---

### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

190928

---

**Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC  
EPA accepted \_\_\_/\_\_\_/\_\_\_

# Supplemental Labeling



Dow AgroSciences LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA

## Quintec® EPA Reg. No. 62719-375

### Control of Powdery Mildew on Strawberry

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

### Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Sphaerotheca macularis</i> )	10 - 14	4 - 6

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

### Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative fungicide. If other diseases are present, use fungicides that are effective on those diseases. If reliable predictive modeling (risk index) systems are available and indicate low disease pressure conditions, minimum label rates and longer application intervals may be used. Maximum label rates and shorter application intervals are recommended for severe or threatening disease conditions.

### Resistance Management

The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is in mode of action Group 13. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a mode of action different from Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.

### Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.

- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec. Apply the alternate product within 14 days of applying Quintec.
- When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different mode of action than Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.
- Do not aerially apply this product.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

**Rotational Crop Restrictions:** Crops grown for food or feed without registered uses for quinoxifen may be replanted 12 months after the last application of quinoxifen.

**Note:** This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC  
R136-025  
Approved: \_\_/\_\_/\_\_  
Replaces R136-009.

# Quintec®

EPA Reg. No. 62719-375

## Registration Notes:

Source Section 3 supplemental labeling based on EPA-accepted text dated September 11, 2006.

### Proposed changes by amendment:

1. Resistance Management: Added section.
2. Use Precautions and Restrictions: (1) Revised third bullet point "Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec."; (2) revised fourth bullet point to add "Apply the alternate product within 14 days of applying Quintec."; (3) revised fifth bullet point.

®Trademark of Dow AgroSciences LLC

RECEIVED  
 FEB 29 2008  
 U.S. DEPARTMENT OF AGRICULTURE  
 OFFICE OF PESTICIDE REGISTRATION  
 1490 L STREET, NW  
 WASHINGTON, DC 20250  
 Tel: 202-720-6275  
 Fax: 202-720-6275  
 E-mail: [Registration@aphis.usda.gov](mailto:Registration@aphis.usda.gov)

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Quintec®

EPA Reg. No. 62719-375

### Control of Powdery Mildew on Head and Leaf Lettuce

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

#### Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Erysiphe cichoracearum</i> )	10 - 14	4 - 6

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

#### Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative fungicide. If other diseases are present, use fungicides that are effective on those diseases. If reliable predictive modeling (risk index) systems are available and indicate low disease pressure conditions, minimum label rates and longer application intervals may be used. Maximum label rates and short application intervals are recommended for severe or threatening disease conditions.

#### Resistance Management

The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is in mode of action Group 13. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a mode of action different from Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.


#### Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.

- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec. Apply the alternate product within 14 days of applying Quintec.
- When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different mode of action than Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not aerially apply this product.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

**Rotational Crop Restrictions:** Crops grown for food or feed without registered uses for quinoxifen may be replanted 12 months after the last application of quinoxifen.

**Note:** This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.



®Trademark of Dow AgroSciences LLC

R136-026

Approved: \_\_\_/\_\_\_/\_\_\_

Replaces R136-010.



# Quintec®

EPA Reg. No. 62719-375

## Registration Notes:


Source Section 3 supplemental labeling based on EPA-accepted text dated September 11, 2006.

### Proposed changes by amendment:

1. Resistance Management: Added section.
2. Use Precautions and Restrictions: (1) Revised third bullet point "Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec."; (2) revised fourth bullet point to add "Apply the alternate product within 14 days of applying Quintec."; (3) revised fifth bullet point.

[Editor's note: Added text is underlined and deleted text is denoted by ~~strikethrough~~.]

®Trademark of Dow AgroSciences LLC



# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Quintec<sup>®</sup>

EPA Reg. No. 62719-375

### Control of Powdery Mildew in Melons<sup>1</sup>

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec<sup>®</sup> fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

#### Directions for Use

<sup>1</sup>Melons including, but not limited to: cantaloupe, casaba, citron melon, crenshaw melon, golden pershaw melon, honey balls, honeydew melon, mango melon, muskmelon, Persian melon, pineapple melon, santa claus melon, snake melon, true cantaloupe, watermelon

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew ( <i>Sphaerotheca fuliginea</i> )	10 - 14	4 - 6

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

#### Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative fungicide. If other diseases are present, use fungicides that are effective on those diseases. If reliable predictive modeling (risk index) systems are available and indicate low disease pressure conditions, minimum label rates and longer application intervals may be used. Maximum label rates and short application intervals are recommended for severe or threatening disease conditions.

#### Resistance Management

The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is in mode of action Group 13. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a mode of action different from Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.

#### Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.

- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec. Apply the alternate product within 14 days of applying Quintec.
- ~~Do not make more than 2 consecutive applications of Quintec before rotating to a product that is effective on powdery mildew and has a different mode of action than Quintec. In either case, apply the alternate product within 14 days of applying Quintec.~~ When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different mode of action than Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not aerially apply this product.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

**Rotational Crop Restrictions:** Crops grown for food or feed without registered uses for quinoxyfen may be replanted 12 months after the last application of quinoxyfen.

**Note:** This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-024

Approved:     /    /    

Replaces R136-008.

# Quintec®

EPA Reg. No. 62719-375

## Registration Notes:

Source Section 3 supplemental labeling based on EPA-accepted text dated September 11, 2006.

### Proposed changes by amendment:

1. Resistance Management: Added section.
2. Use Precautions and Restrictions: (1) Revised third bullet point "Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec."; (2) revised fourth bullet point to add "Apply the alternate product within 14 days of applying Quintec."; (3) revised fifth bullet point.

[Editor's note: Added text is underlined and deleted text is denoted by strikethrough.]

®Trademark of Dow AgroSciences LLC

RECEIVED  
 FEB 29 2008  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 this pesticide is registered.  
 62719-375