

62719-375

3/6/2008

1 of 3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 06 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ray Brinkmeyer
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

Subject: Quintec
EPA Registration No. 62719-375
Your e-mail dated 2/28/2008 regarding supplemental pepper labeling

Dear Mr. Brinkmeyer:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act is acceptable. A stamped copy of the label is enclosed for your records. Submit one copy of the final printed label before you release the product for shipment.

If you have any questions, please contact Tracy Keigwin of my team by phone at (703) 305-6605 or via email at keigwin.tracy@epa.gov.

Sincerely,

Mary L. Waller

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

2083

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Pepper (Bell Pepper, Chili Pepper, Cooking Pepper, Non-Bell Pepper, Pimento, Sweet Pepper)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

| Disease | Recommended Spray Interval (days) | Rate of Quintec (fl oz/acre) |
|---|-----------------------------------|------------------------------|
| powdery mildew (<i>Leveillula taurica</i> , <i>Oidiopsis tuarica</i>) | 10 - 14 | 4 - 6 |

Apply uniformly in a minimum spray volume of 30 gallons per acre that assures thorough coverage of the crop. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. Maximum label rates and short application intervals are recommended for severe or threatening disease conditions.

Resistance Management

The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is in mode of action Group 13. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a mode of action different from Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.

ACCEPTED
 March 06, 2008
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 Reg. No. 62719-375

Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec.
- It is recommended to alternate Quintec with a product with a different mode of action after each application of Quintec. Apply the alternate product within 14 days of applying Quintec.
- When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different mode of action than Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not aerially apply this product.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.

Rotational Crop Restrictions: Crops grown for food or feed without registered uses for quinoxifen may be replanted 12 months after the last application of quinoxifen.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC
R136-020
EPA accepted ___/___/___
Replaces R136-005