

62719-375

10-15-2010

Jacket 1/42



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Dr. Raymond S. Brinkmeyer
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268

OCT 15 2010

Subject: Quintec
EPA Registration No. 62719-375
Your Amendment dated January 20, 2010 Adding aerial application
PRIA Decision No. 427560

Dear Dr. Brinkmeyer:

The master labeling referred to above, submitted in connection with registration under section under 3(c)(7)(b) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided you make the following changes:

Main Labeling

1. Page 10, within the "Application Guidelines" section revise the first sentence for the application method for "Ground" to read, "Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre except as noted in the supplemental labeling..."
2. Page 11, within the directions for use for "Aerial" application revise the first sentence to read, "Apply in a minimum spray volume of 5 gallons of water per acre..."


Supplemental Labeling

3. Pages 19, 23, 26, 29, 32, 35, and 38 within the directions for use for "Aerial" application, revise the sentence "Apply in a minimum spray volume of 5 gallons per acre" to "Apply in a minimum spray volume of 5 gallons of water per acre."

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). One copy of the label stamped "Accepted with comments" is enclosed for your records. If you have any questions, please contact Tamue L. Gibson by phone at (703) 305-9096 or via email at Gibson.tamue@epa.gov.

EPA Reg. No.: 62719-373
Product Name: Quintec
Page 2 of 2

Sincerely,

A handwritten signature in black ink, appearing to read "Shaja B. Joyner". The signature is fluid and cursive, with the first name "Shaja" being more prominent.

Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7504P)

Enclosure

3
42

ACCEPTED
with COMMENTS
In EPA Letter Dated:
OCT 15 2000

(Base label):

Quintec®

Fungicide

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. 62719-375

**A protectant fungicide for the control of powdery mildew in artichoke, gourds,
lettuce, melons, pepper, pumpkin, stone fruits, strawberry, winter squash, cherry,
grapes, and hops.**

Group	13	FUNGICIDE
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)

quinoline..... 22.58%

Other Ingredients 77.42%

Total 100.00%

Contains 2.08 lb of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed • Harmful If Absorbed Through Skin

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling
and before eating, drinking, chewing gum, and using the toilet.**

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables
exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

4
42

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then

5
42

offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read **Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies** at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. _____

6
42

®Trademark of Dow AgroSciences LLC
Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents _____

7
42

(Label booklet cover)

Quintec[®]

Fungicide

A protectant fungicide for the control of powdery mildew in artichoke, gourds, lettuce, melons, pepper, pumpkin, stone fruits, strawberry, winter squash, cherry, grapes, and hops.

Group	13	FUNGICIDE
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)

quinoline..... 22.58%

Other Ingredients 77.42%

Total 100.00%

Contains 2.08 lb of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read **Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies** at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA Est. _____

[®]Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268

Net Contents _____

(Page 1 through end)

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Harmful If Swallowed • Harmful If Absorbed Through Skin**

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, and using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and

springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

In counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur, do not apply this product one-half hour before dusk to one-half hour after dawn.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable containers 5 gallons or larger:

10
42

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Use Directions

Quintec® fungicide is a protectant fungicide for the control of powdery mildew diseases. Quintec does not control existing or latent powdery mildew infections and must be applied on a protectant schedule (latency is the period between infection of plant tissues and the development of visible symptoms). Quintec is compatible with most commonly used insecticides, miticides and fungicides.

General Use Precautions and Restrictions

- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- Do not aerially apply this product on stone fruits, cherry, grapes, and hops. Aerial application is permitted on other crops for which this product is registered for use.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
Brassica (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days

11
42

all other crops grown for food or feed	12 months
--	-----------

¹ **Brassica (cole) leafy vegetables** including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lon), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² **Cucurbit vegetables** including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ **Fruiting vegetables** including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ **Leafy vegetables** including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ **Root and tuber vegetables** including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniel, true, yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Disease Resistance Management

Artichokes, Cherry, Grapes, Hops, Stone Fruits: Quintec is in mode of action Group 13. The number of applications of Quintec per season must not exceed three applications or 50% of the total number of powdery mildew sprays per season, whichever is greater (see crop specific sections for the total number of applications allowed). Apply no more than two consecutive sprays of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

Gourds, Lettuce, Melons, Pepper, Pumpkin, Strawberry, Winter Squash: Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

All Crops:

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow

12
42

AgroSciences representative or by calling 800-258-3033.

Mixing Instructions

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and Quintec or other flowable (aqueous suspension) products to the spray tank. Then add the specified rate of emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

Oil Sprays

Oil sprays may be integrated as part of a powdery mildew management program. Consult local university extension for specific information.

Use of Surfactants

Quintec contains sufficient surfactant to ensure mixability and plant coverage. The use of additional surfactant is not required.

Tank Mixing

Quintec may be applied in tank mix combination with labeled rates of other pesticides, including fungicides, insecticides, or other chemicals used in the culture of the labeled crop provided: (1) the tank mix product is labeled for the timing and method of application for the intended use; and (2) tank mixing with Quintec or other pesticides is not prohibited by the label of the tank mix product.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions including application rates, use precautions, and limitations on the respective product labels.
- Do not exceed specified application rates or maximum use rates if products with the same active ingredient are applied together
- For tank mix partners packaged in water soluble packaging, do not use with products containing boron or mix in equipment previously used to apply boron-containing mixtures unless the tank and spray equipment have been adequately cleaned.
- Always perform a (jar) test to ensure compatibility of products used in tank mixtures.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Quintec and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Rainfast Requirement: When applied alone, Quintec is rainfast 1 hour after application.

Application Guidelines

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Cherry, Grape, Hops, Stone Fruits:

Ground: Apply uniformly by ground in a minimum spray volume of 30 gallons per acre except as noted in

13
42

supplemental labeling. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur.

Aerial: Do not aerially apply this product on stone fruits, cherry, grape, and hops.

Artichoke, Head and Leaf Lettuce, Melons, Pepper, Strawberry, Gourds and Pumpkin, Winter Squash:

Ground: For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur.

Artichoke, Cherry, Grape, Head and Leaf Lettuce, Hops, Melons, Pepper, Stone Fruits, Strawberry, Winter Squash, Gourds and Pumpkin

Crop	Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) ¹
artichoke	powdery mildew (<i>Erysiphe cichoracearum</i>)	6 - 7	8
		10 - 14	4 - 6
cherry	powdery mildew (<i>Podosphaera clandestina</i>)	10 - 14	7
grape	powdery mildew (<i>Uncinula necator</i>)	7	1 - 2
		14	3 - 4
		21	5 - 6.6
head and leaf lettuce	powdery mildew (<i>Erysiphe cichoracearum</i>)	10 - 14	4 - 6
hops	powdery mildew (<i>Sphaerotheca macularis</i>)	14	4 - 8.2
melons ²	powdery mildew (<i>Sphaerotheca fuliginea</i>)	10 - 14	4 - 6
pepper ³	powdery mildew (<i>Leveillula taurica</i> , <i>Oidiopsis taurica</i>)	10 - 14	4 - 6
stone fruits ⁴	powdery mildew (<i>Podosphaera clandestina</i>)	6 - 8	8
		10 - 14	7
strawberry	powdery mildew (<i>Sphaerotheca macularis</i>)	10 - 14	4 - 6
winter squash, gourds, pumpkin	powdery mildew (<i>Sphaerotheca fuliginea</i>)	6 - 8	8
		10 - 14	4 - 6

¹When applied as a dilute spray, maintain the per acre use rate.

²Melons including citron melon, muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, true cantaloupe), watermelon

³Pepper including bell pepper, chili pepper, cooking pepper, non-bell pepper, pimento, sweet pepper

⁴Stone fruits including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh)

Specific Use Restrictions:

Artichoke

- **Preharvest Interval:** None
- **Maximum Application Rate:** Do not apply more than 32 fl oz of Quintec per acre per calendar year.

14
42

- Do not make more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of applications of Quintec per season must not exceed three applications or 50% of the total number of powdery mildew sprays per season, whichever is greater (see crop specific sections for total number of applications allowed). Apply no more than two consecutive sprays of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Cherry

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or more than 35 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 7 days.
- Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

Grape

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6.6 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 7 days for 1 to 2 fl oz per acre rate; 14 days for 3 to 4 fl oz per acre rate; 21 days for 5 to 6.6 fl oz per acre rate.
- Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

Head and Leaf Lettuce

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

15
42

- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Hops

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8.2 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 7 days.
- Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

Melons

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides

16
42

that have a similar target site of action on the same pathogens.

- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Pepper

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Stone Fruits

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or 28 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of

17
42

Quintec per season must not exceed three applications or 50% of the total number of powdery mildew sprays per season, whichever is greater. Apply no more than four applications of Quintec per season. Apply no more than two consecutive sprays of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Strawberry

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Winter Squash, Gourds and Pumpkin

- **Preharvest Interval:** Do not apply within 3 days of harvest.

- **Maximum Application Rate:** Do not apply more than 8 fl oz of Quintec per acre per application or 32 fl oz of Quintec per calendar year.
- Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- **Disease Resistance Management:** Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

19
42

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC
EPA accepted __/__/__

20
42**[Editor's note: Master supplemental labels.]**

List of Supplemental Labels	
Supplemental Title	Previous EPA Accepted Date
Aerial Application Directions on Labeled Crops	New
Control of Powdery Mildew on Stone Fruits (Excluding Cherries)	June 12, 2009
Control of Powdery Mildew on Winter Squash, Gourds, and Pumpkin	June 12, 2009
Control of Powdery Mildew on Artichoke	June 12, 2009
Control of Powdery Mildew on Pepper	May 28, 2009
Control of Powdery Mildew in Melons	May 28, 2009
Control of Powdery Mildew on Strawberry	May 28, 2009
Control of Powdery Mildew on Head and Leaf Lettuce	May 28, 2009

21
42

Supplemental Labeling

**Dow AgroSciences**

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec[®]

EPA Reg. No. 62719-375

Aerial Application Directions on Labeled Crops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec[®] fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing Instructions.

Aerial application of Quintec is permitted on all non-tree and vine crops labeled for use. Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aeriaily.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

[®]Trademark of Dow AgroSciences LLC

R136-035

EPA accepted / /

Initial printing.

22
42

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec[®]

EPA Reg. No. 62719-375

Control of Powdery Mildew on Stone Fruits¹ (Excluding Cherries)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec[®] fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

¹Stone fruits including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh)

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) ¹
powdery mildew (<i>Podosphaera clandestina</i>)	6 - 8	8
	10 - 14	7

¹When applied as a dilute spray, maintain the per acre use rate.

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Apply uniformly by ground in a minimum spray volume of 30 gallons per acre except as noted in supplemental labeling. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of applications of Quintec per season must not exceed three applications or 50% of the total number of powdery mildew sprays per season, whichever is greater (see crop specific sections for the total number of applications allowed). Apply no more than two

23
42

consecutive sprays of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or 28 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Do not aerially apply this product.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
Brassica (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹Brassica (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lon), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

²Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon,

scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, tanager, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-036

EPA accepted / /

Replaces R136-021.

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Winter Squash, Gourds, and Pumpkin

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) ¹
powdery mildew (<i>Sphaerotheca fuliginea</i>)	6 - 8	8
	10 - 14	4 - 6

¹When applied as a dilute spray, maintain the per acre use rate.

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aerially.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is

recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8 fl oz of Quintec per acre per application or 32 fl oz of Quintec per calendar year.
- Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
<i>Brassica</i> (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ *Brassica* (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible

gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniel, true, yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-037

EPA accepted / /

Replaces R136-022.

28
42

Supplemental Labeling


Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Artichoke

ATTENTION:

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) ¹
powdery mildew (<i>Erysiphe cichoracearum</i>)	6 - 7	8
	10 - 14	4 - 6

¹When applied as a dilute spray, maintain the per acre use rate.

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aerially.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of applications of Quintec per season must not exceed three applications or 50% of the total number of powdery mildew sprays per season, whichever is

29
42

greater (see crop specific sections for the total number of applications allowed). Apply no more than two consecutive sprays of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Restrictions:

- **Preharvest Interval:** None
- **Maximum Application Rate:** Do not apply more than 32 fl oz of Quintec per acre per calendar year.
- Do not make more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
Brassica (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ Brassica (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon,

30
42

scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniel, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-038

EPA accepted / /

Replaces R136-023.

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Pepper¹

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

¹Pepper including bell pepper, chili pepper, cooking pepper, non-bell pepper, pimento, sweet pepper

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre)
powdery mildew (<i>Leveillula taurica</i> , <i>Oidiopsis taurica</i>)	10 - 14	4 - 6

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aerially.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
<i>Brassica</i> (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ *Brassica* (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root-and tuber vegetables including arracacha, arrowroot, bitter-cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniar, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-039

EPA accepted / /

Replaces R136-031

34
42

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew in Melons¹

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

¹Melons including cantaloupe, casaba, citron melon, crenshaw melon, golden pershaw melon, honey balls, honeydew melon, mango melon, muskmelon, Persian melon, pineapple melon, santa claus melon, snake melon, true cantaloupe, watermelon

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre)
powdery mildew (<i>Sphaerotheca fuliginea</i>)	10 - 14	4 - 6

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aurally.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50%

35
42

of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
<i>Brassica</i> (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ *Brassica* (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon,

36
42

Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechimā, honey balls, honeydew melon, hubbard squash, hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, lerén, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniel, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-040

EPA accepted ___/___/___

Replaces R136-032.

Supplemental Labeling


Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Strawberry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre)
powdery mildew (<i>Sphaerotheca macularis</i>)	10 - 14	4 - 6

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray, or if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aerially.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per

 37
 42

38
42

crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
Brassica (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ Brassica (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash,

39
42

hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, tanier, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-041

EPA accepted / /

Replaces R136-033.

42
42

Supplemental Labeling


Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

EPA Reg. No. 62719-375

Control of Powdery Mildew on Head and Leaf Lettuce

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre)
powdery mildew (<i>Erysiphe cichoracearum</i>)	10 - 14	4 - 6

Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

Application Method

Ground: For thorough spray coverage and optimum control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

Aerial: Apply in a minimum spray volume of 5 gallons per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Avoid application under conditions when uniform spray coverage cannot be obtained or when excessive spray drift may occur. **Note:** These aerial application directions supersede the previous restriction on the product label against applying this product aurally.

Disease Resistance Management

Quintec is in mode of action Group 13. The number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Alternation with other effective modes of action is recommended after each application of Quintec. When more than one application of Quintec is made per

41
42

crop, at least one of the applications of Quintec must be a tank mixture with a product that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec before alternating to another effective mode of action.

To delay development of fungicide/bactericide resistance, the following practices are recommended:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

Specific Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not graze or harvest forage from cover-crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
<i>Brassica</i> (cole) leafy vegetables (crop group 5) ¹ cucurbit vegetables (crop group 9) ² fruiting vegetables (crop group 8) ³ and okra leafy vegetables (crop group 4) ⁴ root and tuber vegetables (crop group 1) ⁵ sunflower	30 days
all other crops grown for food or feed	12 months

¹ *Brassica* (cole) leafy vegetables including broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai lan), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, white flowering broccoli.

² Cucurbit vegetables including acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese okra, Chinese preserving melon, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, cucuzza, edible gourd, gherkin, golden pershaw melon, hechima, honey balls, honeydew melon, hubbard squash,

42
42

hyotan, mango melon, muskmelon, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini.

³ Fruiting vegetables including bell pepper, chili pepper, cooking pepper, eggplant, groundcherry, hot pepper, pepino, pepper, pimento, sweet pepper, tomatillo, tomato.

⁴ Leafy vegetables including amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock (sorrel), edible-leaved chrysanthemum, endive (escarole), Florence fennel (finocchio), garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, fennel, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket.

⁵ Root and tuber vegetables including arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac (celery root), chayote (root), chicory, Chinese artichoke, chufa, dasheen (taro), edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish (daikon), parsnip, potato, radish, rutabaga, salsify (oyster plant), skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, taniel, true yam, turmeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-042

EPA accepted / /

Replaces R136-034.