



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

November 24, 2015

David Ouimette  
Regulatory Manager  
Dow AgroSciences, LLC.  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: PRIA Label Amendment – Revision to remove restriction on aerial application to tomato  
Product Name: QUINTEC  
EPA Registration Number: 62719-375  
Application Date: 06/25/2015  
Decision Number: 506325

Dear Mr. Ouimette:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 62719-375  
Decision No. 506325

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Maryam Muhammad by phone at 703-347-0301, or via email at Muhammad.maryam@epa.gov.

Sincerely,



Hope Johnson, Product Manager 21  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure      Stamped "Quintec" Approved Label  
Quinoxifen. Review of Label Amendment for Quintec Fungicide (EPA Reg  
No. 62719-375) to Add Aerial Applications to Tomatoes

(Base label):

**Quintec<sup>®</sup>****FUNGICIDE**

**A protectant fungicide for the control of powdery mildew in artichoke, gourds, lettuce, melons, pepper, pumpkin, stone fruits, strawberry, tomato, winter squash, cherry, grapes, and hops.**

Group	13	FUNGICIDE
-------	----	-----------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)	
quinoline .....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%

Contains 2.08 lb of active ingredient per gallon.

**Keep Out of Reach of Children****CAUTION****ACCEPTED**

Nov 24, 2015

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 62719-375

**Precautionary Statements****Hazards to Humans and Domestic Animals****Harmful If Swallowed • Harmful If Absorbed Through Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.**

**Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

**User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

#### (Storage and Disposal for rigid containers 5 gal or less)

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### (Storage and Disposal for refillable rigid containers larger than 5 gal)

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)****Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. \_\_\_\_\_

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced for  
Dow AgroSciences LLC**

**9330 Zionsville Road  
Indianapolis, IN 46268**

**NET CONTENTS \_\_\_\_\_**

(Cover, shipping container)

# Quintec®

## FUNGICIDE

**A protectant fungicide for the control of powdery mildew in artichoke, gourds, lettuce, melons, pepper, pumpkin, stone fruits, strawberry, tomato, winter squash, cherry, grapes, and hops.**

<b>Group</b>	<b>13</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)	
quinoline .....	22.58%
Other Ingredients .....	77.42%
Total .....	100.00%

Contains 2.08 lb of active ingredient per gallon.

**Keep Out of Reach of Children**

## CAUTION

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-375

EPA. Est. \_\_\_\_\_

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET CONTENTS \_\_\_\_\_**

(Page 1 through end)

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

# CAUTION

**Harmful If Swallowed • Harmful If Absorbed Through Skin**

**Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are



more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product one-half hour before dusk to one-half hour after dawn in counties east of the Rocky Mountains where the gray bat, Indiana bat, and the Ozark big-eared bat occur.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Avoid freezing. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable containers 5 gallons or larger:**

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable containers 5 gallons or larger:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Product Information**

Quintec<sup>®</sup> fungicide is a protectant fungicide for the control of powdery mildew diseases. Quintec does not control existing or latent powdery mildew infections and must be applied on a protectant schedule (latency is the period between infection of plant tissues and the development of visible symptoms). Quintec is compatible with most commonly used insecticides, miticides and fungicides.

**Use Restrictions**

- Do not graze or harvest forage from cover crops within areas treated with Quintec.
- Do not aeri ally apply this product on stone fruits, cherry, grapes, or hops. Aerial application is permitted on other crops for which this product is registered for use.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.
- **Rainfast Requirement:** When applied alone, Quintec is rainfast 1 hour after application.
- Not for use in greenhouses or nurseries

**Rotational Crop Restrictions**

The following rotational crops may be planted at intervals defined below following the final application of Quintec.

Crops	Re-Planting Interval
registered crops	no restrictions
all other crops grown for food or feed	30 days

### **Disease Resistance Management**

#### **Cherry, Grapes, Hops, Stone Fruits:**

Quintec is in mode of action Group 13. Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec

**Gourds, Lettuce, Melons, Pepper, Pumpkin, Strawberry, Tomato, Winter Squash:** Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.

#### **All Crops:**

To delay development of fungicide/bactericide resistance:

- Avoid the consecutive use of Quintec or other target site of action Group 13 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local Dow AgroSciences representative or by calling 800-258-3033.

### **Mixing Directions**

Quintec may be applied in tank mix combination with labeled rates of other pesticides, including fungicides, insecticides, or other chemicals used in the culture of the labeled crop provided: (1) the tank mix product is labeled for the timing and method of application for the intended use; and (2) tank mixing with Quintec or other pesticides is not prohibited by the label of the tank mix product.

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and Quintec or other flowable (aqueous suspension) products to the spray tank. Then add the specified rate of emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

#### **Oil Sprays**

Oil sprays may be integrated as part of a powdery mildew management program. Consult local university extension for specific information.

#### **Use of Surfactants**

A moderate rate of a registered nonionic or organosilicone surfactant may improve crop coverage. Penetrants and stickers are not recommended.

#### **Tank Mixing Directions**

- Read carefully and follow all applicable use directions including application rates, use precautions, and limitations on the respective product labels.
- Do not exceed specified application rates or maximum use rates if products with the same active ingredient are applied together
- For tank mix partners packaged in water soluble packaging, do not use with products containing boron

or mix in equipment previously used to apply boron-containing mixtures unless the tank and spray equipment have been adequately cleaned.

- Always perform a (jar) test to ensure compatibility of products used in tank mixtures.

### Tank Mix Compatibility Testing

To determine the physical compatibility of Quintec with other pesticides, perform a jar test. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination must not be used.

## Application Directions

### Application Timing

Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, apply Quintec in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. For optimum results, use maximum label rates and short application intervals for severe or threatening disease conditions.

### Application Method

#### Cherry, Grape, Hops, Stone Fruits:

**Ground:** Apply uniformly by ground application in a minimum spray volume of 30 gallons per acre. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. Do not apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

**Aerial:** Do not aerially apply this product on cherry, grape, hops, and stone fruits.

#### Artichoke, Head and Leaf Lettuce, Melons, Pepper, Strawberry, Gourds and Pumpkin, Tomato, Winter Squash:

**Ground:** For thorough spray coverage and optimum disease control, apply Quintec by ground application. Apply in a minimum spray volume of 30 gallons per acre.

**Aerial:** Apply in a minimum spray volume of 5 gallons of water per acre. Higher spray volumes per acre and conditions that allow uniform spray coverage will provide the greatest level of disease control with Quintec. Do not apply under conditions when uniform spray coverage cannot be obtained or when spray drift may occur.

#### Artichoke, Cherry, Grape, Head and Leaf Lettuce, Hops, Melons (Subgroup 9A), Pepper, Stone Fruits, Strawberry, Tomato, Winter Squash, Gourds and Pumpkin

Crop	Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) <sup>1</sup>	Quinoxifen rate (lb a.i./acre)
artichoke	powdery mildew ( <i>Leveillula taurica</i> )	10 - 14	4 - 6	0.065 - 0.098
cherry	powdery mildew ( <i>Podosphaera clandestina</i> )	10 - 14	7	0.114
grape	powdery mildew ( <i>Erysiphe necator</i> )	14	4	0.065
		21	6.6	0.108
head and leaf lettuce	powdery mildew ( <i>Erysiphe cichoracearum</i> )	10 - 14	4 - 6	0.065 - 0.098

Crop	Disease	Re-Treatment Interval (Days)	Rate of Quintec (fl oz/acre) <sup>1</sup>	Quinoxifen rate (lb a.i./acre)
hops	powdery mildew ( <i>Podosphaera macularis</i> )	14	4 - 8.2	0.065 - 0.134
melons (subgroup 9A) <sup>2</sup>	powdery mildew ( <i>Erysiphe cichoracearum</i> <i>Sphaerotheca fuliginea</i> )	10 - 14	4 - 6	0.065 - 0.098
pepper <sup>3</sup>	powdery mildew ( <i>Leveillula taurica</i> , <i>Oidiopsis taurica</i> )	10 - 14	4 - 6	0.065 - 0.098
stone fruits <sup>4</sup>	powdery mildew ( <i>Sphaerotheca pannosa</i> , <i>Podosphaera leucotricha</i> )	10 - 14	7	0.114
strawberry	powdery mildew ( <i>Sphaerotheca macularis</i> )	10 - 14	4 - 6	0.065 - 0.098
tomato	powdery mildew ( <i>Leveillula taurica</i> , <i>Oidiopsis taurica</i> )	10 - 14	4 - 6	0.065 - 0.098
winter squash, gourds, pumpkin	powdery mildew ( <i>Erysiphe cichoracearum</i> <i>Sphaerotheca fuliginea</i> )	10 - 14	4 - 6	0.065 - 0.098

<sup>1</sup>When applied as a dilute spray, maintain the per acre use rate.

<sup>2</sup>Melons (subgroup 9A) including citron melon, muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, true cantaloupe), watermelon

<sup>3</sup>Pepper including bell pepper, chili pepper, cooking pepper, non-bell pepper, pimento, sweet pepper

<sup>4</sup>Stone fruits including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh)

### Specific Use Restrictions:

#### Artichoke

- **Preharvest Interval:** None
- **Maximum Application Rate:** Do not apply more than 32 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

#### Cherry

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or more than 32 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 7 days.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

#### Grape

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6.6 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than five applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide

that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

#### Head and Leaf Lettuce

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

#### Hops

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8.2 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.
- **Minimum Spray Interval:** 14 days.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

#### Melons (Subgroup 9A)

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

#### Pepper

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

#### Stone Fruits

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or 28 fl oz of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

#### **Strawberry**

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after the first application of Quintec, discontinue its use.

#### **Tomatoes**

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 0.098 lbs a.i. (6 fl oz) of Quintec per acre per application or more than 0.390 lbs a.i. (24 fl oz) of Quintec per acre per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each Quintec application and at regular intervals during crop development. Under certain environmental conditions which cause plant stress, Quintec may cause leaf curling, spotting and/or chlorosis. These conditions may include water stress, adverse temperatures, pest infestations, fertilizer deficiency or excess and poor soil conditions. If symptoms occur after application of Quintec, discontinue its use.

#### **Winter Squash, Gourds and Pumpkin**

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 8 fl oz of Quintec per acre per application or 32 fl oz of Quintec per calendar year.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per crop nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec. It is recommended that the total number of group 13 fungicide sprays per crop should not exceed 50% of the total number of powdery mildew sprays.
- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.

---

### **Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### **Warranty Disclaimer**

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

### **Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

EPA accepted \_\_/\_\_/\_\_