

62719-402

8/10/2011

1 of 5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Raymond Brinkmeyer, Ph.D.  
Regulatory Leader – Regulatory Affairs  
Dow AgroSciences, LLC  
9330 Zionsville Road  
Indianapolis, IN 46268-1054

AUG 1 0 2011

Dear Dr. Brinkmeyer:

SUBJECT: Dithane DF Rainshield  
EPA Registration Number 62719-402  
Your submission of April 8, 2011  
Supplemental labels for almonds, broccoli, pepper, lettuce and cabbage  
Decision Number D447511

The supplemental labels referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended are acceptable provided the supplemental labels are incorporated into the main label within 18 months or at the next label printing, whichever comes first and provided you make the following change:

Above the "Directions for Use" header on each label, add "This supplemental label expires March 31, 2014."

Please submit one copy of the final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or [jones.lisa@epa.gov](mailto:jones.lisa@epa.gov).

Sincerely,

<S>

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7504P)

Enclosures: Stamped Supplemental Labels

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Dithane® DF Rainshield® [Alternate brand name: Dithane® 75 DF Rainshield®] EPA Reg No. 62719-402

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dithane® DF Rainshield® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dithane DF Rainshield according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dithane DF Rainshield.

### Disease Control in Lettuce

Crop	Diseases Controlled	Product Rate per Application (lb/acre)	Application Directions	Restrictions
lettuce	downy mildew	2	Begin applications when disease appears and reapply on a 7- to 10-day treatment schedule.	<ul style="list-style-type: none"> <li>• <b>In California</b>, do not apply more than 8.5 lb of product (6.4 lb active ingredient) per acre and do not apply within 14 days of harvest.</li> <li>• <b>In states other than California</b>, do not apply more than 12.8 lb of product (9.6 lb active ingredient) per acre and do not apply within 10 days of harvest.</li> <li>• Do not apply this product with a U-boom device.</li> <li>• <b>Minimum Re-Treatment Interval: 7 days</b></li> </ul>

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
AUG 1 0 2011**

® Trademark of Dow AgroSciences LLC  
R818-025  
EPA accepted: \_\_\_/\_\_\_/\_\_\_  
Initial printing.

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

**62719-402**

30A5

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Dithane® DF Rainshield®

[Alternate brand name: Dithane® 75 DF Rainshield®]

EPA Reg No. 62719-402

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dithane® DF Rainshield® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dithane DF Rainshield according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dithane DF Rainshield.

### Disease Control in Broccoli and Cabbage

Crop	Diseases Controlled	Product Rate per Application (lb/acre)	Application Directions	Restrictions
broccoli	alternaria leaf spot	2	In plant beds or direct-seeded fields, apply 7 to 10 days after planting or earlier if disease is present. If field applications, apply as soon as disease is present and reapply as needed on a 7- to 10-day spray schedule.	<ul style="list-style-type: none"> <li>• Do not apply more than 12.8 lb of product (9.6 lb active ingredient) per acre.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 7 days of harvest.</li> <li>• Do not apply this product with a U-boom device</li> <li>• <b>Minimum Re-Treatment Interval:</b> 7 days</li> </ul>
cabbage			Refer to broccoli	Refer to broccoli

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**AUG 10 2011**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

**62719-402**

® Trademark of Dow AgroSciences LLC

R818-024

EPA accepted: \_\_\_/\_\_\_/\_\_\_

Initial printing.

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Dithane® DF Rainshield®

[Alternate brand name: Dithane® 75 DF Rainshield®]

EPA Reg No. 62719-402

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dithane® DF Rainshield® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dithane DF Rainshield according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dithane DF Rainshield.

### Disease Control in Peppers

Crop	Diseases Controlled	Product Rate per Application (lb/acre)	Application Directions	Restrictions
peppers	anthracnose early blight phomopsis blight or fruit rot	2 (west of the Mississippi River)  3 (east of the Mississippi River)	Begin application when disease appears and reapply on a 7- to 10-day spray schedule.	<ul style="list-style-type: none"> <li>• West of the Mississippi River, do not apply more than 12.8 lb of product (9.6 lb active ingredient) per acre and do not apply within 7 days of harvest.</li> <li>• East of the Mississippi River, do not apply more than 19.2 lb of product (14.4 lb active ingredient) per acre and do not apply within 7 days of harvest.</li> <li>• Do not apply this product with a U-boom device.</li> <li>• <b>Minimum Re-Treatment Interval: 7 days</b></li> </ul>

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
AUG 10 2011**

® Trademark of Dow AgroSciences LLC

R818-026  
EPA accepted: \_\_\_/\_\_\_/\_\_\_  
Initial printing.

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

62719-402

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Dithane® DF Rainshield®

[Alternate brand name: Dithane® 75 DF Rainshield®]

EPA Reg No. 62719-402

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dithane® DF Rainshield® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dithane DF Rainshield according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dithane DF Rainshield.

### Disease Control in Almond

Crop	Diseases Controlled	Product Rate per Application (lb/acre)	Application Directions	Restrictions
almond	blossom blight ( <i>Monilinia</i> spp) shothole ( <i>Stigmina</i> spp)	6.4	Begin application at dormant to popcorn stage, full bloom or petal fall. Reapply every 7 to 10 days if bloom is staggered and weather is rainy. Do not use less than 10 gallons of spray volume per acre if aerially applied.	<ul style="list-style-type: none"> <li>• Do not apply more than 19.2 lb of product (14.4 lb active ingredient) per acre.</li> <li>• Do not make last application later than 5 weeks after petal fall.</li> <li>• Do not graze livestock in treated area.</li> <li>• Do not apply this product with a U-boom device.</li> <li>• <b>Minimum Re-Treatment Interval: 7 days</b></li> </ul>

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
AUG 1 0 2011**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

**62719-402**

®Trademark of Dow AgroSciences LLC

R818-023

EPA accepted: \_\_\_/\_\_\_/\_\_\_

Initial printing.