Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Control of Powdery Mildew on Hops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally[®] 40WSP fungicide before applying. Carefully follow all
 precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Growth Stage	Rally 40WSP (oz/acre)	Spray Interval (days)	Use Precautions
emergence to training	2 - 4	7 - 10	Do not apply less than 2 oz of Rally 40WSP per acre or adequate efficacy may not be achieved.
training to wire (prior to beginning of bloom when vines are rapidly growing)	4 - 6	5 - 10	Do not apply less than 4 oz of Rally 40WSP per acre or adequate efficacy may not be achieved.
wire to 14 day pre- harvest	6 - 10	7 - 10	Do not apply less than 6 oz of Rally 40WSP per acre or adequate efficacy may not be achieved.

Apply Rally 40WSP in sufficient water for thorough coverage using ground equipment or by air in a minimum spray volume of 10 gallons of water per acre. Thorough coverage is essential. Use the shorter spray interval on susceptible varieties or under heavy disease pressure.

Specific Use Restrictions:

- Do not make more than four applications of Rally 40WSP per acre per season.
- Do not apply more than 2.5 lb of Rally 40WSP (1 lb active ingredient) per acre per season.
- Pre-harvest Interval: Do not apply within 14 days of harvest.
- Do not graze livestock in treated areas or harvest crops grown in treated areas for silage or hay.
- Chemigation: Do not apply Rally 40WSP through any type of irrigation system.
- Resistance Management: Rally 40WSP is in fungicide mode of action group 3. Alternation with other modes of action is recommended after each application of Rally 40WSP. Do not make more than 2 consecutive applications of Rally 40WSP before rotating to a product that is effective application of Rally 40WSP. In either case, the alternate product should be applied within 14 days of an application of Rally 43WSP to posture application control.

[®]Trademark of Dow AgroSciences LLC R182-019 EPA accepted __/__/_ Initial printing

NOV 2 7 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg PRGE 1 of 1

62719 - 410