

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission of November 17, 2006 adding peppers (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) and eggplant
OPPIN decision number 372356

The supplemental label for peppers (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) and eggplant, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

283

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

<S>

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Control of Powdery Mildew on Peppers (Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper) and Eggplant

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Leveillula taurica</i>)	2.5 - 5	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. A lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions. The highest rate and shorter re-treatment interval are recommended for large plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 5 oz (0.31 lb) of Rally 40WSP (0.125 lb ai) per acre per application or 20 oz (1.25 lb) of Rally 40WSP (0.5 lb ai) per acre per season. • Do not make more than 4 applications per season. • Pre-harvest Interval: None. • Minimum Re-treatment Interval: 10 to 14 days

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R182-003
EPA accepted ___/___/___
Initial printing

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 2 0 2008**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

62719-410

NEXT

LABEL

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission of November 17, 2006 adding okra
OPPIN decision number 372356

The supplemental label for okra, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Control of Powdery Mildew on Okra

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Sphaerotheca fuliginea</i>)	2.5 - 5	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. A lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions. The highest rate and shorter re-treatment interval are recommended for large plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 5 oz (0.31 lb) of Rally 40WSP (0.125 lb ai) per acre per application or 20 oz (1.25 lb) of Rally 40WSP (0.5 lb ai) per acre per season. • Do not make more than 4 applications per season. • Pre-harvest Interval: None. • Minimum Re-treatment Interval: 10 to 14 days

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R182-002
EPA accepted ___/___/___
Initial printing

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 20 2008**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

NEXT

LABEL

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission of November 17, 2006 adding artichoke
OPPIN decision number 372356

The supplemental label for artichoke, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

<S>

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES

SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

3 8 3

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Rally® 40WSP EPA Reg. No. 62719-410

Control of Powdery Mildew on Artichoke

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Erysiphe cichoracearum</i>)	4	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage. Use a minimum of 30 gallons of spray volume per acre. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 4 oz (0.25 lb) of Rally 40WSP (0.1 lb ai) per acre per application or 24 oz (1.5 lb) of Rally 40WSP (0.6 lb ai) per acre per season. • Do not make more than 6 applications per season. • Pre-harvest Interval: Do not harvest within 3 days of application • Re-treatment Interval: 14 days • Rotational Crop Restriction: Do not plant any crop other than those listed on the main label for 12 months following the last application of Rally 40WSP.

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC
R182-004
EPA accepted ___/___/___
Initial printing

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 20 2008**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

62719-410

NEXT

LABEL

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission of November 17, 2006 adding black sapote, canistel, mamey sapote, mango, papaya, sapodilla, and star apple
OPPIN decision number 372356

The supplemental label for black sapote, canistel, mamey sapote, mango, papaya, sapodilla, and star apple, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

2 8 3

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

< S >

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Rally® 40WSP EPA Reg. No. 62719-410

Disease Control on Black Sapote, Canistel, Mamey Sapote, Mango, Papaya, Sapodilla, and Star Apple

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Oidium caricae</i>)	10	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 10 oz (0.63 lb) of Rally 40WSP (0.25 lb ai) per acre per application or 80 oz (5 lb) of Rally 40WSP (2 lb ai) per acre per season. • Do not make more than 8 applications per season. • Pre-harvest Interval: None. • Re-treatment Interval: 14 days • Rotational Crop Restriction: Do not plant any crop other than those listed on the main label for 12 months following the last application of Rally 40WSP.

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC
R182-001
EPA accepted ___/___/___
Initial printing

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 2 0 2008**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

NEXT

LABEL

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission of November 17, 2006 adding head and leaf lettuce
OPPIN decision number 372356

The supplemental label for head and leaf lettuce, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

< S >

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

3 8 3

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Control of Powdery Mildew on Head and Leaf Lettuce

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Erysiphe cichoracearum</i>)	5	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 5 oz (0.31 lb) of Rally 40WSP (0.125 lb ai) per acre per application or 20 oz (1.25 lb) of Rally 40WSP (0.5 lb ai) per acre per season. • Do not make more than 4 applications per season. • Pre-harvest Interval: Do not harvest within 3 days of application • Re-treatment Interval: 14 days

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R182-005

EPA accepted ___/___/___

Initial printing

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 20 2008**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

62719-410