

62719-410

08-24-2009

1
23



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

**OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES**

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

AUG 24 2009

Dear Dr. Brinkmeyer:

SUBJECT: Rally 40WSP
EPA Registration Number 62719-410
Your submission dated April 3, 2009
Signal word change
OPPIN Decision Number 408347

The amendment referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided you make the following change:

Page 1 and page 6 – Precautionary Statements:

- Reorder the precautionary statements to read as follows: Harmful if swallowed. Causes Moderate Eye Irritation. Harmful if Absorbed Through Skin. Avoid contact with skin eyes or clothing.
- Remove the vertical space above the sentence beginning "Avoid contact with . . ."
- Change the second sentence from: "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco."
- to: "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, tobacco or using the toilet."

Page 1 and page 6 – First Aid Statements:

Move the entire First Aid section so that it goes between the end of the Precautionary Statements and the beginning of the Personal Protective Equipment section.

Page 1 and page 6 – User Safety Requirements:

- Place the text "Follow manufacturer's instructions . . . separately from other laundry" into a box and title the box "User Safety Requirements."

- Change the sentence: "If no such instructions for washables, use detergent and hot water." to: "If no such instructions for washables exist, use detergent and hot water."

Page 1 and page 6 – Engineering Controls:

Move the "Engineering Controls" section so that it follows the "User Safety Recommendations" box.

Page 13 – Apple and Mayhaw In the last two rows of the disease column, change "bloom, postbloom" to "scab – post infection"

Page 20 – Use Restrictions: Label treated seed:

Change:

"This seed is treated with Rally 40WSP. Do not use for food, feed, oil production or any other purpose except planting. Store away from treated foodstuffs."

to:

"This seed is treated with Rally 40WSP, a myclobutanil product.

Do not allow children, pets or livestock to have access to treated seeds.

Wear long pants, long sleeved shirt and protective gloves when handling treated seed.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting.

Dispose of all excess treated seed by burying seed away from bodies of water.

Do not contaminate bodies of water when disposing of planting equipment wash water.

Dispose of seed packaging or containers in accordance with local requirements.

Do not use for food, feed, oil production or any other purpose except planting.

Store away from treated foodstuffs."

Page 20 – Warranty Disclaimer:

Change the sentence: "Dow AgroSciences makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty." to: "To the extent permitted by law, Dow AgroSciences makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty."

Page 20 – Inherent Risks of Use:

Change the text: All such risks shall be assumed by the buyer: . . ." to:

"To the extent permitted by law, all such risk shall be assumed by the buyer."


Lastly, please reformat following pages so that the header and directions for use appear on the same page. This would be applicable to page 12 (Apple and Mayhaw); page 13 (Gooseberry); page 14 (Grape); page 17 (Asparagus); and page 18 (Tomato).

In the future, please format label text so that when tables extend from one page to the next the table headings appear on both pages.

You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label.

A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones of my team at (703) 308-9424 or at jones.lisa@epa.gov.

Sincerely yours,



Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped label
Acute Toxicity review dated July 29, 2009

4
23

(Base label):

Rally® 40WSP

Fungicide

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile	40%
Other Ingredients	60%
Total	100%

**ACCEPTED
with COMMENTS
in EPA Letter Dated:**

AUG 24 2009

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

62719-410

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (Category A in EPA's Chemical-Resistant Category Selection Chart)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry area above freezing. If in a water soluble pouch, the water soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water soluble pouches from the container except for immediate use.

Pesticide Disposal: Wastes resulting in the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

(Storage and Disposal for nonrigid containers any size)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry area above freezing. If in a water soluble pouch, the water soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water soluble pouches from the container except for immediate use.

Pesticide Disposal: Wastes resulting in the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty

6
23

bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

(Storage and Disposal for refillable rigid containers greater than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry area above freezing. If in a water soluble pouch, the water soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water soluble pouches from the container except for immediate use.

Pesticide Disposal: Wastes resulting in the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry area above freezing. If in a water soluble pouch, the water soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water soluble pouches from the container except for immediate use.

Pesticide Disposal: Wastes resulting in the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

7
23

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-410

EPA Est. _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Net Weight _____

8
23

(Label booklet cover):

Rally® 40WSP

Fungicide

Group	3	FUNGICIDE
--------------	----------	------------------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile	40%
Other Ingredients	60%
Total	100%

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-410

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Weight _____

9
23

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin**

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (Category A in EPA's Chemical-Resistant Category Selection Chart)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves (Category A in EPA's chemical-Resistant Category Selection Chart)
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool dry area above freezing. If in a water soluble pouch, the water soluble pouch may become brittle at storage temperatures below 32°F, but the fungicide is not affected. Do not remove the water soluble pouches from the container except for immediate use.

Pesticide Disposal: Wastes resulting in the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

11
23**Refillable rigid containers larger than 5 gal:**

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

General Information

Rally® 40WSP fungicide is a systemic, protectant and curative fungicide for the control of specific diseases mentioned on this label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

General Use Precautions**Fungicide Resistance Management**

Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with another fungicide mode of action. After two consecutive applications of Rally 40WSP, another myclobutanil product or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Rally 40WSP. Apply the alternate products within the intervals specified on the label for Rally 40WSP. Do not apply Rally 40WSP at rates below those specified on the label. If tank mixing, use the full label rate of Rally 40WSP with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Handling Precautions for Water Soluble Packets: Rally 40WSP is packaged in water soluble pouches. **Do not** remove water soluble packet from the overpack except for immediate use. **Do not** allow water soluble pouches to come into contact with water prior to use. **Do not** handle water soluble pouches with wet hands or wet gloves. **Do not** open water soluble packets. **Do not** remove packets from overwrap except to add the intact packet directly into the spray tank. **Do not** split packets. Carefully reseal package containing unopened water soluble packets and protect package from moisture.

12
—
23

Application Guidelines

Carefully read, understand and follow label use rates and restrictions. Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Lower label rates and 14-day application intervals may be used for small plants and under low disease pressure conditions. Maximum label rates and shorter application intervals are recommended for large plants and for severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions.

Ground Application

Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallons per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

Aerial Application

Apply in a minimum of 10 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.

Chemigation Application

Rally 40WSP must be applied on a regular protectant fungicide schedule, **not an irrigation schedule**. If irrigation cycles are less frequent than the application intervals for Rally 40WSP, ground or aerial applications must supplement chemigation applications to achieve adequate disease control.

Special Use Precautions for Sprinkler Irrigation:

- Apply Rally 40WSP only through the following sprinkler irrigation systems: center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialist, equipment manufacturers or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water system are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Before applying Rally 40WSP through sprinkler irrigation equipment, the chemigation system must meet the following specifications:

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Special Use Directions for Sprinkler Irrigation:

Center-Pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment (use only with electric or oil hydraulic drive systems that uniformly distributes water):

- Determine size of area to be treated.
- Determine the time required to apply no more than 1/4 inch water (6,750 gallons of water per acre) over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run system at 80 to 95% of manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Rally 40WSP required for treatment area.
- Add the required amount of Rally 40WSP and sufficient water to meet the injection time requirements of the solution tank.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until Rally 40WSP solution has cleared the sprinkler head.

Solid-Set, Side (Wheel) Roll, and Hand Move Irrigation Equipment:

- Determine acreage covered by sprinkler.
- Fill injector solution tank with water and adjust flow rate to use contents over a 10 to 30 minute interval.
- Determine the amount of Rally 40WSP required for treatment areas.
- Add the required amount of Rally 40WSP into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
- Inject Rally 40WSP at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until Rally 40WSP solution has cleared the last sprinkler head.

Mixing Directions

Determine the number of acres to be treated, the label use rate and the gallons to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

Fill the tank with one-half of the total amount of water required for the load. Start agitation and maintain agitation throughout mixing and application. Add the required number of unopened pouches (as determined by the dosage recommendations) directly into the spray tank. Water soluble packets will float

14
23

on the surface until the water soluble film dissolves and releases the product. Handling packets with hands should be minimized. Always place Rally 40WSP into solution prior to adding co-applied agricultural chemicals registered for use on specific crop or crops being treated. Complete filling the tank. Before spraying, make sure packets have completely disintegrated and product is thoroughly and uniformly mixed. Depending upon the water temperature and the degree of agitation, the pouches and product should dissolve completely within approximately 10 minutes from the time they were added to the water.

Compatibility

Rally 40WSP is compatible with most commonly used agricultural fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of an appropriate Chemical Producers and Distributors Association certified adjuvant.

Note: Rally 40WSP is compatible with boron and spray oils; however, the water-soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

Rotational Crop Restriction

Crops on this label may be planted immediately after the last treatment. Do not plant other crops within 30 days after the last application. **Note:** When using Rally 40WSP with other registered pesticides, always refer to rotational restrictions and precautions on the other product's label and comply with the most restrictive rotational guidelines.

General Use Directions for Tree Fruits

Best control of labeled diseases is achieved when Rally 40WSP is applied on a 7- to 10-day application schedule.

Rally 40WSP is a systemic fungicide and does not redistribute with rainfall after application. Application equipment spray nozzles should be adjusted to apply a uniform spray throughout the entire tree canopy.

Use the following as guidance in determining the amount of Rally 40WSP to be used per 100 gallons of spray or per acre. Refer to specific tree fruit use directions to determine actual use rates for the control of labeled diseases.

Determination of Use Rates on an Acre Basis

The amount of Rally 40WSP required per acre varies with tree size and the volume of fruit and foliage to be treated. The following summary table may be used as additional guidance for the determination of appropriate per acre use rates for Rally 40WSP:

Tree Height (ft)	Rally 40WSP (oz/acre)	
	Apple and Mayhaw ¹	Stone Fruits
≤10	2.5 - 5	2.5 - 4
15	3.75 - 6	4 - 6
≥20	5 - 10	6

¹For apple scab control, always tank mix Rally 40WSP with a multi-site product such as Dithane®.

Concentrate Spray Applications

Rally 40WSP should be used at the specified use rate per acre in either dilute or concentrate sprays. The following formula should be used to determine the equivalent amount of product per acre in 2X, 3X, etc., spray solutions:

$$\frac{\text{Oz of Rally 40WSP per acre} \times 100}{\text{Spray volume per acre (gallons)}} = \text{Oz of Rally 40WSP per 100 gallons}$$

15
23

Example: An apple orchard consisting of apple trees 18 feet in height will require 5 oz of Rally 40WSP for adequate apple scab control. Application equipment has been calibrated to apply 80 gallons spray per acre, therefore:

$$\frac{5 \text{ oz of Rally 40WSP per acre} \times 100}{80 \text{ gallons per acre}} = 6.25 \text{ oz of Rally 40WSP per 100 gallons}$$

Dilute, Thorough Coverage Application

Dilute thorough coverage applications are based on the amount of spray solution required to thoroughly wet trees until spray run-off. The following specific use directions for apple and mayhaw utilizes a 400 gallon per acre dilute basis and the specific use directions for stone fruits utilizes a 250 gallon per acre dilute basis.

Uses

Almond

Best disease control is achieved in thorough coverage sprays applied on a protectant schedule that does not exceed 10 days.

Diseases	Rally 40WSP		Use Directions	Restrictions
	oz/100 gal ¹	oz/acre		
blossom blight (<i>Monilinia</i> spp.) shothole (<i>Stigmina</i> spp.)	1.25 - 2 (0.5 - 0.8 oz ai)	5 - 8 (2 - 3.2 oz ai)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutanil was used for the early application and at full bloom.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 90 days of harvest. • Do not make more than 3 applications or apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per season.
rust (<i>Tranzschelia</i> spp.)			Apply 6 oz (2.4 oz ai) per acre. Begin applications approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on a protectant schedule exceeding 21 days.	
anthracnose (<i>Colletotrichum</i> spp.)			Applying Rally 40WSP to control blossom blight and shothole will suppress anthracnose.	

¹Based upon a standard of 400 gallons of dilute spray per acre or the equivalent amount of product per acre.

Apple and Mayhaw

Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
----------	------------------------	----------------	--------------

26
23

powdery mildew (<i>Podosphaera</i> spp.)	5 - 10 (2 - 4 oz ai)	Begin application at tight cluster and continue through the second cover spray. Additional sprays may be needed on susceptible varieties or under heavy disease pressure. Use high label rate if powdery mildew was present in previous years.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 14 days of harvest. • Do not apply more than 5 lb of Rally 40WSP (2 lb ai) per acre per season.
rust (<i>Gymnosporangium</i> spp.)	5 - 8 (2 - 3.2 oz ai)	Begin applications at pink stage and continue through the second cover spray.	
scab (<i>Venturia</i> spp.) prebloom		Begin applications at green tip or when environmental conditions become favorable for primary scab development. Apply Rally 40WSP in a tank mixture with a protectant fungicide registered for use on apples on a 7- to 10-day schedule.	
scab bloom		Apply Rally 40WSP in a tank mixture with a protectant fungicide registered for use on apple at the specified rate for improved fruit scab and summer disease control.	
post infection	8 (3.2 oz ai)	Rally 40WSP provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

Berries

Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
Blackberry and Raspberry¹			
cane and leaf rust (<i>Kuehneola</i> spp.) orange rust (<i>Arthuriomyces</i> spp.) powdery mildew (<i>Sphaerotheca</i> spp.) yellow rust (<i>Phragmidium</i> spp.)	1.25 - 2.5 (0.5 - 1 oz ai)	Begin applications as early as bud break. Reapply at 10- to 14-day intervals, depending upon the disease(s) to be controlled. Use the shorter spray interval under heavy disease pressure.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 10 oz of Rally 40WSP (0.25 lb ai) per acre per growing season.
Currant			
powdery mildew (<i>Sphaerotheca</i> spp.) white pine blister rust	5 (2 oz ai)	Apply at pre-bloom, full bloom and 2 weeks later.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 40 oz of Rally 40WSP (1 lb ai) per acre per growing season.
Gooseberry			

17
23

anthracnose (<i>Drepanopeziza</i> spp.)	5 (2 oz ai)	Begin applications when the first leaf has completely unfolded. Reapply at 10- to 14-day intervals as long as environmental conditions favor continued disease development.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 40 oz of Rally 40WSP (1 lb ai per acre per growing season).
powdery mildew (<i>Sphaerotheca</i> spp.) white pine blister rust		Apply at pre-bloom, full bloom and 2 weeks later.	
Strawberry			
leaf blight (<i>Phomopsis</i> spp.) leaf spot (<i>Mycosphaerella</i> spp.) powdery mildew (<i>Sphaerotheca</i> spp.)	2.5 - 5 (1 - 2 oz ai)	Begin applications when disease first appears or when conditions favor disease development. Reapply at 14- to 21-day intervals. Comply with fungicide resistance management recommendations in the General Use Precautions section of this label.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 30 oz of Rally 40WSP (0.75 lb ai) per acre per year.

¹Includes use on the following members of the caneberry group including varieties and/or hybrids:

- | | | |
|---------------------|-------------------------|------------------------|
| bingleberry | dirksen thornless berry | olallieberry |
| black satin berry | Himalayaberry | Oregon evergreen berry |
| boysenberry | hullberry | phenomenalberry |
| Cherokee blackberry | lavacaberry | rangeberry |
| chesterberry | lowberry | ravenberry |
| Cheyenne blackberry | lucretiaberry | rossberry |
| coryberry | mammoth blackberry | Shawnee blackberry |
| darrowberry | marionberry | youngberry |
| dewberry | nectarberry | |

Grape

Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate.

Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
anthracnose (<i>Elsinoe</i> spp.)	3 - 5 (1.2 - 2 oz ai)	Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 14 days of harvest.

18
23

black rot (<i>Guignardia</i> spp.)		Preventative Schedule: Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days. Use a higher rate under heavy disease pressure. Post Infection Schedule: Apply within 72 hours after the beginning of an infection period.	<ul style="list-style-type: none"> Do not apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per year.
powdery mildew (<i>Uncinula</i> spp.)		It is recommended to begin application before bloom (12- to 18-inch shoot growth). Do not extend application intervals beyond 21 days. Use higher rates and/or shorter spray intervals on susceptible varieties or under heavy disease pressure.	

Mint

Crops	Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
peppermint spearmint	powdery mildew (<i>Erysiphe</i> spp.) rust (<i>Puccinia</i> spp.)	4 - 5 (1.6 - 2 oz ai)	Begin application in early spring when plants break dormancy. Reapply on a 14- to 21-day protectant schedule.	<ul style="list-style-type: none"> Preharvest Interval: Do not apply within 30 days of harvest. Do not apply more than 15 oz of Rally 40WSP (0.375 lb ai) per acre per growing season.

Stone Fruits

For stone fruits not specifically listed, follow use directions for the appropriate disease listed below.

Diseases	Rally 40WSP	Use Directions	Restrictions
	oz/acre ¹		
Apricot			
brown rot blossom blight (<i>Monilinia</i> spp.)	2.5 - 6 (1 - 2.4 oz ai)	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutanil was used for the early application and at full bloom.	<ul style="list-style-type: none"> Preharvest Interval: Applications may be made up to the day of harvest. Do not apply more than 2.75 lb of Rally 40WSP (1.1 lb ai) per acre per season.
brown rot (<i>Monilinia</i> spp.)		Apply 6 oz (2.4 oz ai) per acre on a 7- to 14-day protectant schedule. Apply when environmental conditions favor disease development during the month before harvest.	
powdery mildew (<i>Podosphaera</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 10- to 14-day intervals until terminal growth ceases.	

19
23

shothole (<i>Stigmia</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals as long as needed.	
Cherries			
brown rot blossom blight (<i>Monilinia</i> spp.)	2.5 - 6 (1 - 2.4 oz ai)	Begin application at early popcorn stage, before infection occurs. If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutail was used for the early application and at full bloom.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 3.25 lb of Rally 40WSP (1.3 lb ai) per acre per season.
brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)			
leaf spot (<i>Blumeriella</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals. Additional applications after harvest are recommended.	
Nectarine			
brown rot blossom blight (<i>Monilinia</i> spp.)	2.5 - 6 (1 - 2.4 oz ai)	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutanil was used for the early application and at full bloom.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 3.25 lb of Rally 40WSP (1.3 lb ai) per acre per season.
brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)			
shothole (<i>Stigmia</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals as long as needed.	
Peach			
brown rot blossom blight (<i>Monilinia</i> spp.)	2.5 - 6 (1 - 2.4 oz ai)	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutanil was used for the early application and at full bloom.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 3.25 lb of Rally 40WSP (1.3 lb ai) per acre per season.
brown rot (<i>Monilinia</i> spp.)		Refer to apricot	

20
23

powdery mildew (<i>Podosphaera</i> spp.)			
rust (<i>Tranzschelia</i> spp.)		Apply 6 oz (2.4 oz ai) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on a protectant schedule exceeding 21 days.	
Plum, Prune			
brown rot blossom blight (<i>Monilinia</i> spp.)	2.5 - 6 (1 - 2.4 oz ai)	Begin application at green tip before infection occurs. If conditions are favorable for disease development, reapply at full bloom. If an application at petal fall also is needed, alternate to another effective fungicide mode of action if myclobutanil was used for the early application and at full bloom.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to the day of harvest. • Do not apply more than 2.75 lb of Rally 40WSP (1.1 lb ai) per acre per season.
brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
powdery mildew (<i>Podosphaera</i> spp.)			
rust (<i>Tranzschelia</i> spp.)		Refer to peach	

¹250 gallon dilute spray per acre basis.

Vegetables

Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
Asparagus			

rust (<i>Puccinia</i> spp.)	5 (2 oz ai)	Begin applications to the developing ferns after harvest has taken place. Reapply on a protectant schedule not to exceed 14 days. Apply with a spray adjuvant recommended and registered for this specific use pattern.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 180 days of harvest in all states except California. In California, either a 30- or 180-day preharvest interval may be used. • Do not apply to harvestable spears. • 180-day preharvest interval: Do not make more than 6 applications (0.125 lb ai) per growing season. This is equivalent to 30 oz of product (0.75 lb ai) per acre per year. • 30-day preharvest interval: Do not make more than 4 applications (0.125 lb ai) or 20 oz of product (0.5 lb ai) per acre per year.
Cucurbits			
<p>Including: acorn squash, balsam apple, balsam pear, bitter melon, butternut squash, calabaza, cantaloupe, casaba, chayote, Chinese cucumber, Chinese waxgourd, citron melon, crenshaw melon, crookneck squash, cucumber, edible gourd, gherkin, golden pershaw melon, honey balls, honeydew melon, hubbard squash, mango melon, melon, muskmelon, ornamental gourd, Persian melon, pineapple melon, pumpkin, Santa Claus melon, scallop squash, snake melon, spaghetti squash, straightneck squash, summer squash, true cantaloupe, vegetable marrow, watermelon, winter squash, zucchini</p>			
powdery mildew (<i>Erysiphe</i> and <i>Sphaerotheca</i> spp.)	2.5 - 5 (1 - 2 oz ai)	Begin application at first sign of disease development. Reapply on a 7- to 10-day protectant schedule. For the control of other foliar cucurbit diseases, co-applications of registered protectant fungicides should be made according to label use directions.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to and including the day of harvest. • Do not apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per crop. • Observe a 30-day plantback interval between the last application and planting new crops at the treatment site.
Snap Bean			
Asian soybean rust (<i>Phakopsora pachyrhizi</i>) pod tip rot (<i>Rhizoctonia</i> spp.) rust (<i>Uromyces</i> spp.)	4 - 5 (1.6 - 2 oz ai)	Begin applications when rust is first observed. For pod tip rot, begin applications when pods begin to develop. Reapply on a 7- to 10-day protectant schedule if conditions remain favorable for disease development.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be made up to and including the day of harvest. • Do not apply more than 1.25 lb of Rally 40WSP (0.5 lb ai) per acre per crop. • Observe a 30-day plantback interval between the last application and planting new crops at the treatment site.
Tomato			

22
23

powdery mildew (<i>Leveillula</i> spp.)	2.5 - 4 (1 - 1.6 oz ai)	Begin applications at the first sign of disease or when environmental conditions are favorable for disease development. Do not exceed 21 days between applications. Apply using a minimum of 20 gallons of water per acre by ground or a minimum of 10 gallons of water by air.	<ul style="list-style-type: none"> • Preharvest Interval: Applications may be up to and including the day of harvest. • Do not apply more than 1.25 lb of Rally 40WSP (0.5 lb ai) per acre per crop. • Observe a 30-day plantback interval between the last application and planting new crops at the treatment site.
---	----------------------------	---	---

Non-Food Use

Diseases	Rally 40WSP oz/acre	Use Directions	Restrictions
Hybrid Poplar (For use in nurseries or forested areas used for wood pulp production)			
rust (<i>Melampsora</i> spp.)	4 - 6 (1.6 - 2.4 oz ai)	Begin applications at the first sign of disease. Reapply at 10- to 14-day intervals.	<ul style="list-style-type: none"> • Do not apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per year.
Douglas Fir (Nursery Use Only)			
needle rust (<i>Melampsora</i> spp.)	5 - 10 (2 - 4 oz ai)	Begin applications in early spring. Reapply at 2- to 3-week intervals until the threat of infection is past. Apply with a spray adjuvant recommended and registered for this specific use pattern to obtain good spray coverage and disease control.	<ul style="list-style-type: none"> • Do not apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per year.
Loblolly Pine (Nursery Use Only)			
fusiform rust (<i>Cronartium quercuum</i>)	5 - 10 (2 - 4 oz ai)	Begin applications in early spring. Reapply at 2- to 3-week intervals until the threat of infection is past. Apply with a spray adjuvant recommended and registered for this specific use pattern to obtain good spray coverage and disease control.	<ul style="list-style-type: none"> • Do not apply more than 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per year.

Commercial Cotton Seed Treatment

Use Rally 40WSP at time of planting as a seed treatment to protect against sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*) which impairs good cotton seed germination and seedling development. Rally 40WSP may be applied to dry seed with conventional slurry or mist seed treating equipment. For best results, the seed must be completely and uniformly covered with fungicide. Rally 40WSP can be tank mixed with other properly labeled cotton seed protectants such as Apron TL for Pythium control. Do not tank mix with products whose labels contain a prohibition against tank mixing.

Use Rate

Use 0.05 to 0.0675 lb (0.8 to 1 oz) of Rally 40WSP (0.02 to 0.027 lb ai) per 100 lb of cotton seed to reduce seedling damage caused by sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*).

23
23**Use Restrictions:**

Label treated seed "This seed is treated with Rally 40WSP. Do not use for food, feed, oil production or any other purpose except planting." Do not graze cotton plants grown from treated seed. Store away from feed and foodstuffs.

In California, this product must be applied through a closed system such as WECO-RED (Returnable Evacuation Device).

For cotton seed treatment, always use Rally 40WSP with an EPA approved dye or colorant that imparts an unnatural color to the seed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC
EPA accepted ___/___/___