

62719-410

2/17/2012

10f3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

FEB 17 2012

Darin Lickfeldt, Ph.D.  
Regulatory Leader  
Dow AgroSciences, LLC  
9330 Zionsville Road  
Indianapolis, IN 46268-1054

Dear Dr. Lickfeldt:

SUBJECT: Rally 40WSP  
EPA Registration Number 62719-410  
Your submission of May 26, 2011  
OPPIN decision number 449796

The supplemental label for artichoke, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

1. This supplemental label expires on February 17, 2015 and must not be used or distributed after this date. This statement should be prominently located towards the top of label.
2. Move the heading "Directions for Use" to where the "Control of Rust" is now, and move the heading "Control of Rust" to where "Directions for Use" was.
3. Move the sentence: "It is a violation of Federal law to use this product in a manner inconsistent with its labeling" immediately below the "Directions for Use" heading.
4. Remove the word "Attention" from the box.
5. Add the following text after "Resistance Management":  
Rotational Crops: Fields treated with Laredo EW can be rotated at any time to crops that are listed on a registered myclobutanil label (Laredo EG, Nova® and Rally®) immediately after the last treatment. Do not plant other crops within 30 days after the last application of a product containing myclobutanil


6. Add the following text after "Resistance Management":  
Rotational Crops: Do not allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment.

A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or [jones.lisa@epa.gov](mailto:jones.lisa@epa.gov).

Sincerely yours,



Shaja B. Joyner  
Product Manager (20)  
Fungicide Branch  
Registration Division (7505P)

Enclosure: Stamped supplemental label  
Residue chemistry review dated January 19, 2012

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Rally® 40WSP EPA Reg. No. 62719-410

### Control of Rust (*Phakopsora pachyrhizi*) in Soybean

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

#### Directions for Use

Refer to product label for General Information, Mixing and Application instructions.

Disease	Rally 40WSP oz/acre	Use Directions	Restrictions
rust ( <i>Phakopsora pachyrhizi</i> )	2.5 - 5 (1 - 2 oz ai)	Apply using ground or aerial equipment in an adequate spray volume to achieve good coverage and canopy penetration. For best results, apply as early as possible, prior to infection or at first sign of disease, and make subsequent applications at 10- to 14-day intervals. For established disease, use a higher rate in the rate range. For aerial application, apply Rally 40WSP in a minimum spray volume of 5 gallons per acre.	<ul style="list-style-type: none"> <li>• <b>Preharvest Interval:</b> Do not apply within 28 days of harvest.</li> <li>• Do not make more than two applications or apply more than 10 oz of Rally 40WSP (4 oz ai) per acre per season.</li> </ul>

**Resistance Management:** Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. After two consecutive applications of Rally 40WSP, another myclobutanil product or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Rally 40WSP. Apply the alternate products within the intervals specified on the label for Rally 40WSP. Do not apply Rally 40WSP at rates below those specified on the label. If tank mixing, use the full label rate of Rally 40WSP with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC  
R182-038  
EPA accepted \_\_\_/\_\_\_/\_\_\_  
Initial printing

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**FEB 17 2012**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

62719-410