62719-412 (Dop/2007 C) Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Laredo[®] EC

EPA Reg. No. 62719-412

Control of Rust (*Phakopsora pachyrhizi*) in Soybean

ATTENTION


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Laredo[®] EC fungicide before applying. Carefully follow all
 precautionary statements and applicable use directions.
- Use of Laredo EC according to this supplemental labeling is subject to all use precautions and
- limitations imposed by the label affixed to the product container.

Directions for Use

Apply 4 to 8 fl oz of Laredo EC (0.0625 to 0.125 lb myclobutanil) per acre, using ground or aerial equipment, in an adequate spray volume to achieve good coverage and canopy penetration. For aerial application, apply Laredo EC in a minimum spray volume of 5 gallons per acre. For best results, apply as early as possible, prior to infection or at first sign of disease, and make subsequent applications at 10- to 14-day intervals. For established disease, use the higher rate in the rate range.

Specific Use Restrictions:

- Do not allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.
- Preharvest Interval: Do not make applications within 28 days of harvest.
- **Do not** make more than two applications or apply more than 16 fl oz of Laredo EC (0.25 lb ai myclobutanil) per acre per year.
- Rotation Crops: Fields treated with Laredo EC can be rotated at any time to crops that are listed on a registered myclobutanil label (Laredo EW, Nova[®] and Rally[®]) immediately after the last treatment. Do not plant other crops within 30 days after the last application of a product containing myclobutanil.



[®]Trademark of Dow AgroSciences LLC

R153-085 Accepted: __/__/ Initial printing.