

62719-414

12/20/2005

1/3

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Kelthane® 50WSP

EPA Reg. No. 62719-414

For Control of Mites in Stone Fruits†

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Kelthane® 50WSP miticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Kelthane 50WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Kelthane 50WSP.

Directions for Use

†Stone fruits including, but not limited to: Apricot, Chickasaw Plum, Damson Plum, Japanese Plum, Nectarine, Peach, Plum, Plumcot, Prune (Fresh), Sweet Cherry, Tart Cherry

Kelthane 50WSP is a contact miticide, not a systemic pesticide. Mites must come in contact with Kelthane 50WSP as it is sprayed or be exposed to residues on the leaf and/or fruit surface. Since mites often infest the under surfaces of leaves and fruits, effective control requires an application that thoroughly and uniformly coats all foliar plant surfaces. For maximum effectiveness, apply Kelthane 50WSP to low populations of mites.

Refer to product label for Kelthane 50WSP miticide for General Use Precautions, Mixing and Application Instructions.

Pests and Application Rates:

Pests	Kelthane 50WSP (lb/acre)
European red mite	2 - 3
McDaniel mite	(1 -1.5 lb ai per acre)
pacific spider mite	
plum nursery mite	
strawberry spider mite	
two-spotted spider mite	

ACCEPTED
DEC 20 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-414

Specific Use Directions:

Application Timing: Apply when mites reach threshold levels as specified by Agricultural Experiment Station or State Extension Service Specialists.

Application Rate: Apply as a dilute or concentrate ground spray with a properly calibrated and maintained sprayer in sufficient water to assure thorough and uniform coverage of foliage and fruit surfaces.

Spray Adjuvants: Kelthane 50WSP may be used with or without any additional adjuvants. A spreader-sticker is beneficial in improving the spray coverage on difficult-to-wet foliage and to smooth out spray deposits. Kelthane 50WSP should only be combined with a spreader-sticker that has been proven safe to crops.

Specific Use Restrictions:

- Do not exceed 1.5 lb ai dicofol per acre per application.
- Do not make more than 1 application per year.
- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Chemigation:** Do not apply through any type of irrigation equipment.


Note: This product is toxic to fish and wildlife. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R849-002

EPA accepted: / /

Initial printing



H3B / Kelthane 50WSP / Prop Sec 3 Supp Stone Fruits / 05-28-04
File: R849-002 Kelthane 50WSP Sec 3 Mite Control in Stone Fruits 28May04d.doc

Kelthane[®] 50WSP

EPA Reg. No. 62719-414

Proposed Section 3 supplemental labeling for use of Kelthane 50WSP for control of mites in stone fruits.

®Trademark of Dow AgroSciences LLC

