

62719-416

1/19/2007

10238



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Dr. Raymond S. Brinkmeyer
Dow Agro Sciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268

JAN 19 2007

Subject: Enable® 2F Fungicide
EPA Registration Number 62719-416
Your amendment dated November 19, 2004 and its seven related re-submissions for decision 351452, and related submissions dated April 6, 2005, December 3, 1999, and May 1, 1996, and related petitions.

We have reviewed the subject amended labeling, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended. The amended labeling is acceptable provided that:

1. You make the following changes to the labeling.
 - a. In the second paragraph in the "General Information" section of the master label near the bottom of page 6 change "...approved for use in registered pesticide products..." to "...approved for use with registered pesticide products...".
 - b. In the "Almond" specific use directions section in the master label and the "Almond" supplemental label add a statement specifying that the minimum retreatment interval is 10 days.
 - c. In the "Banana" specific use directions section in the master label specify whether application can be made by air or ground or both and if the amount of water required varies from the amounts mentioned in the "General Information" section, specify that, as well.
 - d. In the "Cranberry" specific use directions section in the master label and the "Cranberry" supplemental label add the following statement: "Do not allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last

application."

e. In the "Pecan" specific use directions section in the master label add the statement "Do not apply after shuck split."

f. At the start of the first sentence in the first paragraph in the "Limitation of Remedies" section in the master label delete "To the extent permitted by law" and replace it with "To the extent consistent with applicable law".

g. At the beginning of each of the two sentences in the second paragraph (it directly follows the two numbered items) in the "Limitation of Remedies" section in the master label add "To the extent consistent with applicable law,".

2. The GC/NPD analytical method (TR 34-94-142) for determining residues in livestock commodities passes a Tolerance Method Validation initiated by Health Effects Division.

3. You submit to the Agency each of the following studies or materials by its stated due date.

a. Provide, by March 31, 2007, to the Agency's Analytical Chemistry Branch repository at Ft. Meade, Maryland one gram each of the reference standards for the two Fenbuconazole metabolites RH 9129 and RH 9130.

b. Submit an acceptable citrus processing study of Fenbuconazole residues by March 31, 2008. As a part of this study samples must be analyzed within the interval during which Fenbuconazole and its metabolite residues have been shown to be stable (30 months for fruit and 12 months for oil and dried pulp). However, this study does not need to include analysis of the juice samples because current data for juice are adequate.

c. Submit an acceptable Apple processing study of Fenbuconazole residues by June 30, 2008.

d. Submit acceptable studies that report two additional residue field trials for Blueberries; one each from regions 2 and 5, by June 30, 2008.

e. Submit acceptable studies that report a complete set of 20 Fenbuconazole residue field trials on Wheat forage by June 30, 2008. These field trials must be conducted using the currently proposed use pattern,

which reflects one application of Fenbuconazole on a Wheat forage crop at 0.063 lb active ingredient per acre. Although conditions will vary between test sites, the sampling intervals for Wheat forage must be consistent enough for the Agency to be able to compare residues and determine an appropriate pre-harvest interval (PHI).

f. Submit acceptable studies that report an additional 12 Fenbuconazole residue field trials on Wheat hay by June 30, 2008. These field trials must be conducted using the currently proposed use pattern, which reflects up to two applications of Fenbuconazole on a Wheat hay crop at 0.063 lb active ingredient per acre per application. Although conditions will vary between test sites, the sampling intervals for Wheat hay must be consistent enough for the Agency to be able to compare residues and determine an appropriate pre-harvest interval (PHI).

g. Submit an acceptable study(ies) of Fenbuconazole toxicity to aquatic vascular plants and estuarine/marine nonvascular plants by June 30, 2008. This involves conducting tier one studies under GLN 122-1.

h. Submit an acceptable Fenbuconazole aquatic field dissipation study by December 31, 2008.

i. Submit an acceptable Fenbuconazole mysid life cycle study by December 31, 2008.

j. Submit a Tier II seedling emergence (of oat, cabbage, cucumber, and turnip) and vegetative vigor (of onion, lettuce, corn, oat, and soybean) study of Fenbuconazole by June 30, 2008.

k. Submit a Confidential Statement of Formula (or its equivalent) for the Fenbuconazole product Indar 5EC (which is not registered in the U.S.) by March 31, 2007.

4. Submit one copy of your final printed master and supplemental labels before you release product bearing this amended labeling for shipment.

"In the Agency's risk assessment for 1,2,4-triazole and its metabolites triazole alanine, and triazole acetic acid, dated February 7, 2006 (see docket number 2005-0497 at <http://www.regulations.gov>), the Agency identified additional data that are needed for triazole-derivative active ingredients

which have the 1,2,4,-triazole ring. The Agency plans to issue a Data Call-In for all triazole-derivative active ingredients which will identify all data needs for these active ingredients listed in the February 7, 2006 risk assessment. The Data Call-In will not address data needs that are related to parent-only data requirements."

The supplemental labels must be incorporated into your master label at the next printing or within 18 months from the date of this letter.

If you have any questions about this letter, please contact John Bazuin at (703)305-7381 or bazuin.john@epa.gov.

Sincerely yours,

5
Cynthia Giles-Parker, Chief
Fungicide Branch
Registration Division (7505P)

- Attachments: Master label stamped "ACCEPTED with COMMENTS"
- Almond supplemental label stamped "ACCEPTED with COMMENTS"
- Apple supplemental label stamped "ACCEPTED"
- Blueberry supplemental label stamped "ACCEPTED"
- Citrus Fruits supplemental label stamped "ACCEPTED"
- Cranberry supplemental label stamped "ACCEPTED with COMMENTS"
- Peanut supplemental label stamped "ACCEPTED"
- Plum and Prune supplemental label stamped "ACCEPTED"
- Sugarbeets supplemental label stamped "ACCEPTED"
- Wheat supplemental label stamped "ACCEPTED"

50238

ACCEPTED
with COMMENTS
In EPA Letter Dated

JAN 19 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

62719-416

(Base Label)

(Logo) Dow AgroSciences

Enable[®] 2F

Fungicide

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile 23.5%
Other Ingredients 76.5%
Total 100.0%

Contains 2 lb of active ingredient per gallon
U.S. Patent No. 4366165

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Inhaled

Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistant category selection sheet.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber.
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards
 This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-416

EPA Est. _____

®Trademark of Dow AgroSciences LLC
 Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents __

7 of 38

(Label Booklet)

(Logo) Dow AgroSciences LLC

Enable[®] 2F

Fungicide

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile	23.5%
Other Ingredients	76.5%
Total	100.0%

Contains 2 lb of active ingredient per gallon
U.S. Patent No. 4366165

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for Precautionary Statements and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-416

EPA Est. _____

®Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Contents ____

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
Engineering Controls	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Resistance Management	-
Use Rate Determination	-
Mixing	-
Application	-
Orchard Use	-
Aerial Spray Drift Management	-
Compatibility	-
Plantback Intervals	-
Uses	-
Almond	-
Apple	-
Banana	-
Blueberry	-
Citrus Fruits	-
Cranberry	-
Peanut	-
Pecan	-
Plum and Prune	-
Stone Fruits	-
Sugarbeets	-
Wheat	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Harmful If Inhaled**

Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistant category selection sheet.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber.
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool dry well-ventilated area, but not below 32°F (0°C).

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Steps to be Taken in Case Material is Released or Spilled: : Appropriate protective equipment must be worn when handling a spill of this material. Remove contaminated clothing promptly. Wash exposed skin areas with soap and water immediately after exposure. Contain spills immediately with inert materials (e.g., sand, earth). Transfer spilled liquid and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Material Safety Data Sheet.

General Information

Enable® 2F fungicide is a systemic fungicide with preventative and curative properties used for the control of specific diseases listed on this label. For optimal disease control with Enable 2F, apply sprays on a preventative spray program.

A non-polymer containing spray adjuvant approved for use in registered pesticide products may be added to spray solutions according to manufacturer's use instructions to achieve optimum disease control.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

110f 38

Use Rate Determination

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum suggested use rates can be used, while maximum suggested label rates and shortened spray schedules are recommended for severe or threatening disease conditions. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

Mixing

Thoroughly shake container before using contents. Triple-rinse container used to measure Enable 2F and add rinsate to spray tank. Add a registered spray adjuvant after Enable 2F is fully dispersed in the spray solution. **Note:** Reduced product efficacy may occur if water containing suspended soil particles such as water from ponds, streams or unlined ditches is used.

Application

Do not make ground or aerial applications within 75 feet of streams, rivers, ponds, lakes or reservoirs.

Ground: Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. For tree fruits and nuts the same amount of Enable 2F should be applied per acre in either dilute or concentrate sprays.

Aerial: Apply in a minimum of 5 gallons of water per acre on annual crops and 10 gallons of water per acre on perennial tree fruits and nuts. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Flagmen and loaders should wash thoroughly before eating, drinking, or smoking and at the end of each day's spray operation.

Orchard Use

Apply Enable 2F to orchards using a dilute or concentrate spray.

Dilute Spray: Dilute or thorough coverage applications are based upon the amount of spray solution necessary to thoroughly wet the trees until spray run-off. To determine the number of gallons of dilute spray required per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance.

Concentrate Spray: Concentrate spray applications assume all of the tree is uniformly covered with spray solution but not to the point of run-off. A lower spray volume is used to deliver the same application rate per acre as used for a dilute spray. A minimum of 50 gallons of spray solution per acre is recommended for concentrate sprays with Enable 2F.

Aerial Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

120238

- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Chemigation: Do not apply this product through any type of irrigation system.

Compatibility

Enable 2F is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. For any registered adjuvant, follow the manufacturer's use instructions to achieve proper spray concentrations.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

Uses

Almond

Disease	Enable 2F		Use Directions	Restrictions
	(fl oz/100 gal) [†]	(fl oz/acre)		
blossom blight (<i>Monilinia</i> spp.)	2	4 - 6 (0.06 - 0.094 lb active)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 3 applications or apply more than 18 fl oz of Enable 2F (0.28 lb active) per acre per year. Preharvest Interval: Do not apply within 160 days of harvest.

[†] Based on a standard of 300 gallons of dilute spray per acre or the equivalent amount of product per acre as a concentrate spray.

Apple

Disease	Enable 2F (fl oz/acre) [†]	Use Directions	Restrictions
---------	--	----------------	--------------

flyspeck (<i>Scytothorium pomi</i>) powdery mildew (<i>Podosphaera leucotricha</i>) rusts (<i>Gymnosporangium</i> spp.) scab (<i>Venturia inaequalis</i>) sooty blotch (<i>Gloeodes pomigena</i>)	6 - 8 (0.094 - 0.12 lb active)	Begin application at green tip or when environmental conditions become favorable for disease development. Apply on a 7- to 10-day schedule through the first cover spray; thereafter, apply on a 10- to 14-day schedule. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 4 applications or apply more than 32 fl oz of Enable 2F (0.5 lb active) per acre per year. Preharvest Interval: Do not apply within 14 days of harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
--	-----------------------------------	---	---

† Maintain per acre use rate regardless of spray volume per acre.

Banana

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
sigatoka (<i>Mycosphaerella</i> spp.)	6 (0.09 lb active)	Apply when leaves first appear and repeat every 14 to 21 days or as required. Best control of diseases is achieved on a protectant application schedule and when an emulsifiable spray oil is added to the spray solution.	<ul style="list-style-type: none"> Do not make more than 8 applications of Enable 2F (0.75 lb active) per acre per year. Preharvest Interval: Up to day of harvest. Do not graze livestock in treated areas. Do not cut cover crops for livestock food.

Blueberry

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
alternaria (<i>Alternaria</i> spp.) anthracnose (<i>Colletotrichum gloeosporioides</i>) leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) mummy berry disease (<i>Monilinia vaccinii-corymbosi</i>) phomopsis twig blight and fruit rot (<i>Phomopsis vaccinii</i>) powdery mildew (<i>Microsphaera vaccinii</i>)	6 (0.094 lb active)	For mummy berry and phomopsis twig blight, begin applications at early green tip and make subsequent applications at intervals of 8 to 14 days. For control of powdery mildew and other diseases listed, begin applications prior to or at onset of disease development and repeat at 8- to 14-day intervals. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 4 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. Preharvest Interval: Do not apply within 30 days of harvest. Do not graze livestock in treated fields or offer treated material as a livestock feed item.

140e38

rusts (<i>Pucciniastrum</i> spp.)			
--	--	--	--

Citrus Fruits

Including, but not limited to: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange (Sweet, Sour and Mandarin), Pummello, and Tangerine

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Enable 2F with a nonionic surfactant or spray oil at the manufacturer's recommended rate and making thorough coverage applications..

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
greasy spot (<i>Mycosphaerella</i> <i>citri</i>)	8 (0.12 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application or for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> Do not make more than 3 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. Preharvest Interval: Up to day of harvest. Minimum Re-Treatment Interval: 21 days. Do not allow livestock to graze in treated areas. Do not apply Enable 2F with polymer based spray adjuvants.
scab (<i>Elsinoe</i> <i>fawcettii</i>)		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold (<i>Capnodium</i> spp.)		Applications made for control of greasy spot or scab will also control sooty mold.	

Cranberry

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, it is recommended to use an agriculturally registered non-polymer non-ionic surfactant at the manufacturer's recommended rate.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
cottonball tip blight (<i>Monilinia</i> <i>oxycocci</i>)	6 - 12 (0.094 - 0.188 lb active)	Begin applications when 50% of the shoots have begun to elongate. Refer to your local spray recommendation guides for more specific application timing information.	<ul style="list-style-type: none"> Do not make more than 4 applications or apply more than 48 fl oz of Enable 2F (0.75 lb active) per acre per year. Preharvest Interval: Do not apply within 30 days of harvest.
cranberry fruit rot complex (1)		Begin applications prior to bloom at the onset of	

		disease. Continue on a 10- to 14-day spray schedule depending upon local conditions. Refer to your local recommendation guides for more specific application information.	
--	--	---	--

1. Cranberry fruit rot disease complex may contain one or more of the following fungal pathogens: *Allantophomopsis lycopodina*, *Botryosphaeria vaccinii*, *Coleophoma empetri*, *Diaporthe vaccinii*, *Glomerella cingulata-vaccinii*, *Godronia cassandrae*, *Phyllosticta vaccinii*, *Physalospora vaccinii*, *Colletotrichum* spp. , *Phomopsis vaccinii* and *Strasseria geniculata*.

Peanut

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
early leaf spot (<i>Cercospora arachidicola</i>)	4 - 6 (0.062 - 0.094 lb active)	Begin foliar applications approximately 4 weeks after planting or at first sign of disease development and continue application on a 10- to 14-day schedule.	<ul style="list-style-type: none"> Do not make more than 4 applications or apply more than 32 fl oz of Enable 2F (0.5 lb active) per acre per year. Preharvest Interval: Do not apply within 14 days of harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.
late leaf spot (<i>Cercosporidium personatum</i>)	6 - 8 (0.094 - 0.12 lb active)		

Pecan

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
downy spot (<i>Mycosphaerella caryigena</i>) gnomonias leafspot (<i>Gnomonia dispersa</i>) leaf blotch (<i>Mycosphaerella dendroides</i>) powdery mildew (<i>Microsphaera penicillata</i>) scab (<i>Cladosporium caryigenum</i>) vein spot (<i>Gnomonia nerviseda</i>)	8 (0.125 lb active)	Begin applications at bud break and continue at 10- to 14-day intervals through pollination. Apply at 14- to 21-day intervals through the cover sprays. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 6 applications or apply more than 48 fl oz of Enable 2F (0.75 lb active) per acre per season. Preharvest Interval: Do not apply after shuck split or within 28 days of harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

Plum and Prune

1100238

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, a non-polymer spray adjuvant/wetting agent labeled for use on food crops should be added to spray solutions at the manufacturers' recommended rate.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
blossom blight (<i>Monilinia</i> spp.)	6 (0.094 lb active)	Begin applications at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	<ul style="list-style-type: none"> Do not make more than 4 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. Preharvest Interval: Up to day of harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
fruit brown rot (<i>Monilinia</i> spp.)		Begin applications 2 to 3 weeks before harvest and reapply on a 7- to 14-day spray schedule.	

Stone Fruits

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.

Crop	Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
apricot	blossom blight (<i>Monilinia</i> spp.)	6 (0.094 lb active)	Begin applications at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and at petal-fall.	<ul style="list-style-type: none"> Do not make more than 8 applications or apply more than 48 fl oz of Enable 2F (0.75 lb active) per acre per season. Preharvest Interval: Up to day of harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
	fruit brown rot (<i>Monilinia</i> spp.)		Begin applications 2 to 3 weeks before harvest using a 7- to 10-day spray interval.	
cherries	blossom blight (<i>Monilinia</i> spp.) fruit brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
	leaf spot (<i>Blumeriella</i> spp.)		Follow blossom blight schedule and continue application at 10- to 14-day intervals. Additional foliar applications may be made after harvest.	
nectarine	blossom blight (<i>Monilinia</i> spp.) fruit brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
peach	blossom blight		Refer to apricot	

	(<i>Monilinia</i> spp.) fruit brown rot (<i>Monilinia</i> spp.)			
	scab (<i>Cladosporium</i> spp.)			Begin applications at shuck split. Make 2 to 3 subsequent thorough coverage applications at 10- to 14-day intervals.

Sugarbeets

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
cercospora leaf spot (<i>Cercospora beticola</i>) powdery mildew (<i>Erysiphe</i> spp.)	8 (0.125 lb active)	Apply Enable 2F at intervals of 14 to 21 days beginning when conditions are favorable for the development of disease. For resistance management, alternate applications of Enable 2F with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant such as crop oil concentrate that is labeled for crop use should be added to the spray mixture at the manufacturer's recommended rate. Add an adjuvant to the spray solution last. For best spray coverage of foliage, apply aerially in a recommended minimum of 5 to 7 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 2 applications or apply more than 16 fl oz of Enable 2F (0.25 lb active) per acre per year. Preharvest Interval: Do not apply within 14 days of harvest.

Wheat

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
glume blotch (<i>Septoria nodorum</i>) leaf blight, spot blotch, barley scald (<i>Rhynchosporium</i> spp.) leaf blotch (<i>Septoria tritici</i>)	4 (0.062 lb active)	Apply anytime through heading (Feekes growth stage 10.3 to 10.5) on a 7- to 21-day schedule. Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 15 gallons of water per	<ul style="list-style-type: none"> Do not make more than 2 applications or apply more than 8 fl oz of Enable 2F (0.12 lb active) per acre per year. Preharvest Interval: Do not apply after completion of flowering (Feekes growth stage 10.5) or within 35 days of harvest.

powdery mildew <i>(Erysiphe graminis)</i> rusts (<i>Puccinia</i> spp.) tan spot, net blotch <i>(Pyrenophora</i> spp.)		acre.	<ul style="list-style-type: none"> • Preharvest Interval: Do not apply within 14 days of harvest for forage and hay.
--	--	-------	--

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted: __/__/__

190838

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Almond

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F		Use Directions	Restrictions
	(fl oz/100 gal) †	(fl oz/acre)		
blossom blight (<i>Monilinia</i> spp.)	2	4 - 6 (0.06 - 0.094 lb active)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 3 applications or apply more than 18 fl oz of Enable 2F (0.28 lb active) per acre per year. • Preharvest Interval: Do not apply within 160 days of harvest.

† Based on a standard of 300 gallons of dilute spray per acre or the equivalent amount of product per acre as a concentrate spray.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-015

EPA accepted: ___/___/___

Initial printing.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN 19 2007

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

62719-416

200638

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F EPA Reg. No. 62719-416

Disease Control in Apple

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F (fl oz/acre) [†]	Use Directions	Restrictions
flyspeck (<i>Scytothyrrium pomi</i>)	6 - 8 (0.094 - 0.12 lb active)	Begin application at green tip or when environmental conditions become favorable for disease development. Apply on a 7- to 10-day schedule through the first cover spray; thereafter, apply on a 10- to 14-day schedule. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 32 fl oz of Enable 2F (0.5 lb active) per acre per year. • Preharvest Interval: Do not apply within 14 days of harvest. • Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
powdery mildew (<i>Podosphaera leucotricha</i>)			
rusts (<i>Gymnosporangium</i> spp.)			
scab (<i>Venturia inaequalis</i>)			
sooty blotch (<i>Gloeodes pomigena</i>)			


[†] Maintain per acre use rate regardless of spray volume per acre.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-016
EPA accepted: / /
Initial printing.



210438

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Apple / 11-13-06
file: R833-016 Enable 2F Sec 3 Disease Control in Apple 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Blueberry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
alternaria (<i>Alternaria</i> spp.) anthracnose (<i>Colletotrichum gloeosporioides</i>) leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) mummy berry disease (<i>Monilinia vaccinii-corymbosi</i>) phomopsis twig blight and fruit rot (<i>Phomopsis vaccinii</i>) powdery mildew (<i>Microsphaera vaccinii</i>) rusts (<i>Pucciniastrum</i> spp.)	6 (0.094 lb active)	For mummy berry and phomopsis twig blight, begin applications at early green tip and make subsequent applications at intervals of 8 to 14 days. For control of powdery mildew and other diseases listed, begin applications prior to or at onset of disease development and repeat at 8- to 14-day intervals. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. • Preharvest Interval: Do not apply within 30 days of harvest. • Do not graze livestock in treated fields or offer treated material as a livestock feed item.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Plantback Intervals


Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R833-017

EPA accepted: ___/___/___

Initial printing.



230e38

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Blueberry / 11-13-06
file: R833-017 Enable 2F Sec 3 Mummy Disease in Blueberry 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Citrus Fruits†

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

†Including, but not limited to: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange (Sweet, Sour and Mandarin), Pummello, and Tangerine

Refer to product label for Enable 2F for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Enable 2F with a nonionic surfactant or spray oil at the manufacturer's recommended rate and making thorough coverage applications.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
greasy spot (<i>Mycosphaerella citri</i>)	8 (0.12 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application or for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> • Do not make more than 3 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. • Preharvest Interval: Up to day of harvest. • Minimum Re-Treatment Interval: 21 days. • Do not allow livestock to graze in treated areas. • Do not apply Enable 2F with polymer based spray adjuvants.
scab (<i>Elsinoe fawcettii</i>)		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold (<i>Capnodium spp.</i>)		Applications made for control of greasy spot or scab will also control sooty mold.	

ACCEPTED

JAN 19 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act, the applicant for the pesticide

62719-416

250638

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Citrus Fruits / 11-13-06
file: R833-018 Enable 2F Sec 3 Disease Control in Citrus Fruits 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-018

EPA accepted: __/__/__

Initial printing.

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Enable[®] 2F

EPA Reg. No. 62719-416

Disease Control in Cranberry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, it is recommended to use an agriculturally registered non-polymer non-ionic surfactant at the manufacturer's recommended rate.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
cottonball tip blight (<i>Monilinia oxycocci</i>)	6 - 12 (0.094 – 0.188 lb active)	Begin applications when 50% of the shoots have begun to elongate. Refer to your local spray recommendation guides for more specific application timing information.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 48 fl oz of Enable 2F (0.75 lb active) per acre per year. • Preharvest Interval: Do not apply within 30 days of harvest.
cranberry fruit rot complex (1)		Begin applications prior to bloom at the onset of disease. Continue on a 10- to 14-day spray schedule depending upon local conditions. Refer to your local recommendation guides for more specific application information.	

1. Cranberry fruit rot disease complex may contain one or more of the following fungal pathogens: *Allantophomopsis lycopodina*, *Botryosphaeria vaccinii*, *Coleophoma empetri*, *Diaporthe vaccinii*, *Glomerella cingulata-vaccinii*, *Godronia cassandrae*, *Phyllosticta vaccinii*, *Physalospora vaccinii*, *Colletotrichum* spp., *Phomopsis vaccinii* and *Strasseria geniculata*.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN 19 2007

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

62719-416

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Cranberry / 11-13-06
file: R833-019 Enable 2F Sec 3 Disease Control in Cranberry 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-019

EPA accepted: ___/___/___

Initial printing.

300238

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Peanut

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
early leaf spot (<i>Cercospora arachidicola</i>)	4 - 6 (0.062 - 0.094 lb active)	Begin foliar applications approximately 4 weeks after planting or at first sign of disease development and continue application on a 10- to 14-day schedule.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 32 fl oz of Enable 2F (0.5 lb active) per acre per year. • Preharvest Interval: Do not apply within 14 days of harvest. • Do not feed hay or threshings or allow livestock to graze in treated areas.
late leaf spot (<i>Cercosporidium personatum</i>)	6 - 8 (0.094 - 0.12 lb active)		

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R833-020

EPA accepted: / /

Initial printing.

ACCEPTED

JAN 19 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered with EPA

62719-416

3/06/38

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Peanut / 11-13-06
file: R833-020 Enable 2F Sec 3 Disease Control in Peanut 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

32 of 38

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Plum and Prune

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, a non-polymer spray adjuvant/wetting agent labeled for use on food crops should be added to spray solutions at the manufacturers' recommended rate.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
blossom blight (<i>Monilinia</i> spp.)	6 (0.094 lb active)	Begin applications at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	<ul style="list-style-type: none"> • Do not make more than 4 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. • Preharvest Interval: Up to day of harvest. • Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
fruit brown rot (<i>Monilinia</i> spp.)		Begin applications 2 to 3 weeks before harvest and reapply on a 7- to 14-day spray schedule.	

Resistance Management


Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-021

EPA accepted: / /

Initial printing.



330238

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Plum-Prune / 11-13-06
file: R833-021 Enable 2F Sec 3 Disease Control in Plum-Prune 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

340f38

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Sugarbeets

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
cercospora leaf spot (<i>Cercospora beticola</i>) powdery mildew (<i>Erysiphe</i> spp.)	8 (0.125 lb active)	Apply Enable 2F at intervals of 14 to 21 days beginning when conditions are favorable for the development of disease. For resistance management, alternate applications of Enable 2F with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant such as crop oil concentrate that is labeled for crop use should be added to the spray mixture at the manufacturer's recommended rate. Add an adjuvant to the spray solution last. For best spray coverage of foliage, apply aerially in a recommended minimum of 5 to 7 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 2 applications or apply more than 16 fl oz of Enable 2F (0.25 lb active) per acre per year. • Preharvest Interval: Do not apply within 14 days of harvest.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

ACCEPTED
 JAN 19 2007
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, use is limited to the pesticide label only.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R833-022

EPA accepted: / /

Initial printing.

3600F 38

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Sugarbeets / 11-13-06
file: R833-022 Enable 2F Sec 3 Disease Control in Sugarbeets 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC

37 of 38

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable® 2F

EPA Reg. No. 62719-416

Disease Control in Wheat

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable® 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
glume blotch (<i>Septoria nodorum</i>) leaf blight, spot blotch, barley scald (<i>Rhynchosporium</i> spp.) leaf blotch (<i>Septoria tritici</i>) powdery mildew (<i>Erysiphe graminis</i>) rusts (<i>Puccinia</i> spp.) tan spot, net blotch (<i>Pyrenophora</i> spp.)	4 (0.062 lb active)	Apply anytime through heading (Feekes growth stage 10.3 to 10.5) on a 7- to 21-day schedule. Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 15 gallons of water per acre.	<ul style="list-style-type: none"> • Do not make more than 2 applications or apply more than 8 fl oz of Enable 2F (0.12 lb active) per acre per year. • Preharvest Interval: Do not apply after completion of flowering (Feekes growth stage 10.5) or within 35 days of harvest. • Preharvest Interval: Do not apply within 14 days of harvest for forage and hay.

Resistance Management


Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R833-023
EPA accepted: ___/___/___
Initial printing.



38 of 38

A3B / Enable 2F / Prop Sec 3 Supp Disease Control in Wheat / 11-13-06
file: R833-023 Enable 2F Sec 3 Disease Control in Wheat 13Nov06d.doc

Enable[®] 2F

EPA Reg. No. 62719-416

Registration Notes:

Proposed Section 3 supplemental labeling.

[®]Trademark of Dow AgroSciences LLC