

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Confirm\* 2F

EPA Reg. No. 62719-420

### For Control of Insects Infesting Canola

#### ATTENTION


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Confirm\* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Confirm 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Confirm 2F.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

#### Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
beet armyworm ( <i>Spodoptera exigua</i> ) bertha armyworm ( <i>Mamestra configurata</i> )	8 (0.12 lb ai/acre)	For early season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	<ul style="list-style-type: none"> <li>• Do not apply more than 16 fl oz of Confirm 2F per application or exceed 64 fl oz of Confirm 2F per season.</li> <li>• <b>Pre-harvest Interval:</b> Do not harvest within 14 days of application.</li> <li>• Refer to <b>Rotational Crop Restrictions</b> in the body of the product label.</li> </ul>
	8 - 16 (0.12 to 0.25 lb ai/acre)	For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, re-apply in 10 to 14 days to protect new growth until moth flights and/or hits subside.	



#### Specific Use Directions

Equipment and spray volume should be carefully adjusted to assure thorough uniform coverage of infested parts of the crop.

**Ground Application:** Apply a minimum of 8 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 10 gallons per acre to densely foliated or difficult to cover crop to ensure thorough coverage.

**Aerial Application:** Apply in a minimum of 5 gallons per acre.

**Note:** This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

\*Trademark of Dow AgroSciences LLC

D3B / Confirm 2F / Prop Sec 3 Supp Canola / 06-23-03

D06-809-XXX

EPA-accepted   /  /  

Initial printing.